

Amyotrophic Lateral Sclerosis–Frontotemporal Lobar Dementia in 3 Families With p.Ala382Thr *TARDBP* Mutations

Adriano Chiò, MD; Andrea Calvo, MD, PhD; Cristina Moglia, MD; Gabriella Restagno, MD; Irene Ossola, BS; Maura Brunetti, BS; Anna Montuschi, PsyD; Angelina Cistaro, MD; Anna Ticca, MD; Bryan J. Traynor, MD; Jennifer C. Schymick, PhD; Roberto Mutani, MD; Maria Giovanna Marrosu, MD; Maria Rita Murru, BS; Giuseppe Borghero, MD

Background: TAR DNA-binding protein 43, encoded by the *TARDBP* gene, has been identified as the major pathological protein of frontotemporal lobar dementia (FTLD) with or without amyotrophic lateral sclerosis (ALS) and sporadic ALS. Subsequently, mutations in the *TARDBP* gene have been detected in 2% to 3% of patients with ALS (both familial and sporadic ALS). However, to our knowledge, there is only 1 description of 2 patients with FTLD and *TARDBP* gene mutations who later developed motor neuron disease.

Objective: To describe cognitive abnormalities in 3 Italian families with familial ALS and *TARDBP* gene mutations.

Design, Setting, and Participants: Genetic, neuropsychological, and neuroimaging analyses in 36 patients with familial non–superoxide dismutase 1 gene (*SOD1*) ALS and 280 healthy controls.

Main Outcome Measure: We identified 3 index cases

of familial ALS carrying the p.Ala382Thr missense mutation of the *TARDBP* gene and with clinical, neuroimaging, and neuropsychological features of FTLD.

Results: The p.Ala382Thr missense mutation of the *TARDBP* gene was absent in the 280 controls. It was present in all affected members of the 3 families for whom DNA was available. All affected members of the 3 families developed FTLD after the onset of ALS, confirmed by neuropsychological testing and hypometabolism in frontal associative areas assessed with fludeoxyglucose F 18 positron emission tomography and computed tomography.

Conclusions: Three apparently unrelated families with familial ALS carrying the p.Ala382Thr *TARDBP* missense mutation developed FTLD. In these families, FTLD cosegregates with ALS. Patients with ALS carrying *TARDBP* mutations may develop FTLD.

Arch Neurol. 2010;67(8):1002-1009

AMYOTROPHIC LATERAL SCLEROSIS (ALS) is a neurodegenerative disorder of adult life, characterized by a progressive impairment of motor function at spinal and bulbar levels. The disorder is sporadic in approximately 95% of cases, while it is inherited in the remaining cases (familial ALS [FALS]), typically with autosomal dominant transmission. To date, mutations in 3 genes have been described as causative for ALS, namely *SOD1* (OMIM 147450), *TARDBP* (OMIM 605078), *FUS/TLS* (OMIM 137070), and *FIG4* (OMIM 612577).¹⁻⁵ It is now well recognized that about 10% of patients with ALS also develop florid symptoms of frontotemporal lobar dementia (FTLD) and a further 30% to 50% of patients have subthreshold disturbances of frontal lobe dysfunction.⁶ Patients carrying *TARDBP* mutations can present with FTLD and subsequently develop

motor neuron dysfunction later in the course of their illness (ie, FTLD–motor neuron disease [FTLD–MND]).⁷ However, to our knowledge, there are no reports of *TARDBP* mutations being associated with the related, although distinct, phenotype of ALS–FTLD (ie, a patient presenting with ALS who subsequently develops FTLD).

Here we describe 3 apparently unrelated families carrying p.Ala382Thr *TARDBP* missense mutations in whom FTLD developed after ALS in all affected members.

METHODS

SUBJECTS

A total of 36 non-*SOD1*- and non-*FUS*-carrying proband cases of FALS seen at the Turin ALS center were screened for *TARDBP* gene mutations. A case was considered to be

Author Affiliations are listed at the end of this article.

Table. Summary of Patients' Clinical Characteristics

| Patient No./Sex | Age at Onset of ALS, y | Age at Onset of FTLT, y | Type of Onset | FTLD Classification ^a | TARDBP Mutation Detected |
|-----------------|------------------------|-------------------------|---------------|----------------------------------|--------------------------|
| Family A | | | | | |
| II-7/F | 58 | 58.5 | Bulbar | ALS-FTLD | Yes |
| I-4/M | 78 | 79 | Spinal | ALS-FTLD | NA ^b |
| III-1/M | 33 | 35 | Spinal | ALS-FTLD | Yes |
| Family B | | | | | |
| III-2/M | 44 | 44.5 | Bulbar | ALSbi | Yes |
| III-1/M | 33 | 35 | Spinal | ALS-FTLD | NA ^b |
| II-7/F | 69 | 70 | Spinal | ALS-FTLD | Yes |
| Family C | | | | | |
| III-1/F | 25 | 29 | Spinal | ALSbi | Yes |
| II-3/M | 43 | 44.5 | Spinal | ALS-FTLD | NA ^b |
| II-6/M | 64 | 65 | Spinal | ALS-FTLD | Yes |

Abbreviations: ALS, amyotrophic lateral sclerosis; FTLT, frontotemporal lobar dementia; NA, not applicable.

^aA classification of ALS-FTLD indicates ALS with FTLT meeting the criteria by Neary et al⁹; a classification of ALSbi indicates ALS with behavioral impairment.¹⁰

^bFor these patients, DNA was not available.

familial if a first- or second-degree relative had been diagnosed with ALS. Amyotrophic lateral sclerosis was diagnosed according to revised El Escorial criteria,⁸ and FTLT was diagnosed according to the criteria by Neary et al.⁹

TARDBP MUTATIONAL SCREENING

Genomic DNA was extracted from 1 mL of peripheral blood with the Biorobot MDx DSP system (Qiagen Inc, Valencia, California). All the coding exons and 100 base pairs of the flanking intron-exon boundaries of *TARDBP* were amplified by polymerase chain reaction using specific primers,² sequenced using the BigDye Terminator v3.1 sequencing kit (Applied Biosystems, Inc, Carlsbad, California), and run on an ABI Prism 3100-Avant genetic analyzer (Applied Biosystems, Inc).

NEUROPSYCHOLOGICAL BATTERY

The patients underwent neuropsychological testing according to the consensus criteria for the diagnosis of frontotemporal cognitive and behavioral syndromes in ALS.¹⁰ The following tests were performed: Frontal Systems Behavior Scale (FrSBe), using the family form evaluated by a close relative (scores of ≤ 59 indicated normal; 60-64, borderline; and ≥ 65 , pathological); Mini-Mental State Examination; Wisconsin Card Sorting Test; Trail Making Test A and B; Stroop Color-Word Interference Test; letter and category fluency test; Wechsler Memory Scale-Revised (form 2); Rey-Osterrieth Complex Figure Test; Token Test; Wechsler Adult Intelligence Scale-Revised; and Raven's Coloured Progressive Matrices. Anxiety and depression were assessed with the Hospital Anxiety and Depression Scale. When applicable, test results were corrected for age and educational level.

POSITRON EMISSION TOMOGRAPHY AND COMPUTED TOMOGRAPHY

Patients fasted for at least 6 hours prior to examination. Each patient was given a dose of 185 MBq of fludeoxyglucose F 18 (FDG) (to convert to megacurie, multiply by 2.7×10^{-11}) by intravenous injection 60 minutes before the scan. Between FDG injection and the brain scans, the patients remained still and relaxed in a quiet room. The positron emission tomographic (PET) and computed tomographic (CT) scans were acquired by a Discovery ST-E PET/CT system (GE Healthcare, Wauke-

sha, Wisconsin), combining a helical multislice CT scanner and a designed bismuth germanate block detector PET tomograph, in 3-dimensional modality. The FDG-PET/CT images were acquired through 2 sequential scans: CT brain scan (thickness, 3.75 mm; 140 kV; 60-80 mA/s) and PET brain scan (1 field of view of 30 cm transaxially). The PET scan was initiated immediately after the CT examination in order to use the CT data for the attenuation correction of the PET data. Data were collected in 128×128 matrices.

The PET/CT images have been evaluated by visual analysis and statistical mapping techniques to compare a patient's brain PET/CT scan with those in a database of age-matched healthy subjects (Cortex ID; GE Healthcare). The resulting data with 3-dimensional stereotactic surface projection maps of the brain allow for quantitative visualization of patterns and severity of such abnormalities.

STANDARD PROTOCOL APPROVALS AND PATIENT CONSENTS

The study has been approved by the ethical committee of our institution. The patients provided written informed consent.

RESULTS

While performing mutational screening of *TARDBP* in 36 index cases of FALS, we found 3 apparently unrelated FALS patients carrying a c.1144G>A (p.Ala382Thr) missense mutation. This mutation was not found in 280 healthy subjects from our center and was not found in 956 healthy controls in 2 previous studies on *TARDBP* mutations in ALS.^{11,12} Each proband belonged to a large, apparently unrelated pedigree. Because the 3 cases showed evident clinical symptoms of FTLT, they underwent an in-depth neuropsychological battery and a brain PET/CT scan. A summary of patients' clinical characteristics is reported in the **Table**.

FAMILY A

In family A (**Figure 1A**), the proband (II-7) is a 60-year-old woman who developed dysphagia, dysarthria, tongue hypotrophy, and emotional lability at age 58 years,




Figure 1. Pedigrees of family A (A), family B (B), and family C (C) with chromatograms of part of exon 6 of the *TARDBP* gene showing c.1144G>A (p.Ala382Thr) mutations in all of the examined patients. Squares indicate males; circles, females; diagonal lines, deceased; open symbols, unaffected; solid symbols, affected; P, proband; and WT, wild type.

followed by wasting and weakness of the upper limbs, mainly distally. She had generalized hyperreflexia, and Babinski and Hoffmann signs were present. At neurophysiological examination, she showed chronic and active denervation, confirmed by a muscle biopsy. Brain and spinal cord magnetic resonance imaging (MRI) results were negative. The diagnosis was definite ALS. One year after the onset of motor signs, she developed apathy, mental rigidity, perseverative behavior, distractibility, aspon-taneity, and speech stereotypies. Her FrSBe total score was 70 (premorbid score of 49), with increased scores in all domains (apathy, disinhibition, and executive dys-functions). Neuropsychological testing demonstrated a global frontal cognitive deterioration. The PET/CT brain study showed a reduction of glucose metabolism on the associative frontal cortex, associative parietal cortex, anterior cingulus, and left head of caudate nuclei. The reduction was more marked on the left side. The PET/CT images from a healthy member of the same family (**Figure 2**) compared with the PET/CT images from this patient (**Figure 3**) are shown.

The proband's father (I-4) developed wasting and weakness of the distal upper limbs at age 78 years, rapidly followed by dysphagia and dysarthria. At neurological examination, he also showed diffuse hyperreflexia. At neurophysiological examination, he showed chronic and active denervation. He was diagnosed as having definite ALS. According to clinical records, he showed marked apathy and executive dysfunction, confirmed by the FrSBe total score of 75 (premorbid score of 37). He died 2 years later of respiratory failure. No DNA was available from this subject.

The eldest son of the proband (III-1) developed distal wasting and weakness of the upper limbs at age 33 years, followed rapidly by wasting of the lower limbs and respiratory impairment. At neurophysiological examination, he showed chronic and active denervation of all limbs. His diagnosis was definite ALS. He was tracheos-tomized at age 35 years. One year after the onset of ALS, the patient showed restless pacing, distractibility, im-pulsivity, motor imperistence, mental rigidity, and in-flexibility. His FrSBe total score was 69 (premorbid score of 32). He is still alive 48 months after the onset of ALS and carries the p.Ala382Thr missense mutation of the *TARDBP* gene.

FAMILY B

In family B (**Figure 1B**), the proband (III-2) is a 44-year-old man who developed dysarthria and dysphagia with tongue hypotrophy and emotional lability at age 43 years. He also presented with pyramidal signs of the upper and lower limbs, with Babinski and Hoffman signs. He was diagnosed as having possible ALS. Head and spinal cord MRI results were normal, with the exception of a type I Chiari malformation. One year later, he developed weakness and wasting of both hands and of the shoulder girdle. He subsequently developed behavior symptoms including apathy, irritability, distractibility, stereotyped behavior, perseveration, and avolition. The total FrSBe score of 89 was pathological, with an increase from the pre-morbid score of 48. Neuropsychological testing results




Figure 2. Right lateral (A), left lateral (B), right medial (C), left medial (D), anterior (E), posterior (F), superior (G), and inferior (H) positron emission tomography/computed tomography statistical parametric maps of a healthy member of family A. P indicates posterior; A, anterior; R, right; and L, left.

were pathological for spatial abilities and for letter fluency, with deficits in cognitive inhibitory abilities and impulsivity. The PET/CT scan showed a reduction of metabolism in the frontal associative and frontal mesial areas, right temporal associative areas, and head of right caudatus nucleus (**Figure 4**).

The proband's mother (II-7) carries the p.Ala382Thr missense mutation of the *TARDBP* gene; she developed hypotrophy and loss of strength in both hands at age 69 years. She had pyramidal signs in the upper and lower limbs, and the neurophysiological examination showed active and chronic denervation involving the upper and lower limbs. Head and spinal cord MRI results as well as cerebrospinal fluid examination results were normal. She was diagnosed as having probable, laboratory-supported ALS. A few months after the onset of motor impairment, she developed mental rigidity, persevera-




Figure 3. Right lateral (A), left lateral (B), right medial (C), left medial (D), anterior (E), posterior (F), superior (G), and inferior (H) positron emission tomography/computed tomography statistical parametric maps of patient II-7 from family A. The images show the reduction in fludeoxyglucose F 18 uptake on the frontal and parietal associative cortices, the anterior cingulus, and the caudate nuclei. The metabolism is reduced on the left side (white arrows). A slight reduction in fludeoxyglucose F 18 uptake on the right cerebellar cortex (crossed cerebellar diaschisis) is also evident (yellow arrow). P indicates posterior; A, anterior; R, right; and L, left.

tive behavior, and distractibility. Neuropsychological testing demonstrated a mild global frontal cognitive deterioration. She did not undergo PET/CT brain study. She is still alive 18 months after the onset of ALS.

The proband's father (II-8) developed hand wasting, weakness, and generalized pyramidal signs at age 64 years, followed by spreading of muscle hypotrophy to the lower limbs. At neurophysiological examination, he showed chronic and active denervation of all limbs. Head and spinal cord MRI results and cerebrospinal fluid examination results were normal. He was diagnosed as having definite ALS. At age 67 years, he underwent noninvasive ventilation and later tracheotomy. One year after the onset of muscle deficit, he developed anxiety, depression,




Figure 4. Right lateral (A), left lateral (B), right medial (C), left medial (D), anterior (E), posterior (F), superior (G), and inferior (H) positron emission tomography/computed tomography statistical parametric maps of patient III-2 from family B. The images show the hypometabolic areas on the frontal and temporal associative cortices, the frontal mesial cortex, and the caudate nuclei (arrows). P indicates posterior; A, anterior; R, right; and L, left.

and agitation of psychotic character. He is still alive 55 months after the onset of ALS. He does not carry *SOD1*, *FUS*, or *TARDBP* mutations.

The proband's brother (III-1) developed hypotrophy and weakness of left hand muscles at age 33 years. The disturbances progressed to involvement of proximal muscles, with the development of a clinical picture of flail arm syndrome. Pyramidal signs were evident 2 years after the onset of symptoms. The neurophysiological examination showed active and chronic denervation involving the upper and lower limbs, and cerebrospinal fluid examination results were normal. Head and spinal cord MRI results were normal, with the exception of a type I Chiari malformation. He was diagnosed as having definite ALS with a flail arm phenotype. Two years after the onset of ALS, he showed neurobehavioral disturbances

and his neuropsychological testing demonstrated a global frontal cognitive deterioration. He died 58 months after the onset of ALS at age 39 years. No DNA was available from this subject.

FAMILY C

In family C (Figure 1C), the proband (III-1) is a 31-year-old woman who developed distal hypotrophy and weakness of the lower limbs at age 25 years, followed by a diffusion of the weakness of both hands 1 year later. She also manifested pyramidal signs of the upper and lower limbs, with Babinski and Hoffman signs. She was diagnosed as having definite ALS. At neurophysiological examination, she showed chronic and active denervation of all limbs.

Brain and spinal cord MRI results were normal. She had no bulbar signs or symptoms. Three years after the onset of motor symptoms, she developed disinhibition, emotional lability, distractibility, asponaneity, and speech stereotypy. Her total FrSBe score of 62 was borderline pathological, with an increase from the premorbid score of 34. Neuropsychological testing showed deficits in visuo-spatial abilities and impulsivity. She is alive 72 months after the onset of symptoms. Her PET/CT scan is shown in **Figure 5**.

The proband's mother (II-2) is aged 65 years and is neurologically normal. She carries the p.Ala382Thr missense mutation of the *TARDBP* gene. She refused to undergo neuropsychological assessment and PET/CT scan.

One of the proband's maternal uncles (II-3) developed wasting and weakness of the hands, rapidly spreading to involve his lower limbs, with pyramidal signs at age 43 years. The diagnosis of definite ALS was confirmed by neurophysiological examination and by head and cervical MRI scan. Bulbar signs developed 1 year later, followed by an overt FTLT. The patient died 30 months after the onset of ALS. No DNA from this subject was available.

Another maternal uncle (II-6) developed weakness of the lower limbs with slight distal atrophy and hyperreflexia and spastic tetraparesis at age 64 years. Neurophysiological examination showed diffuse chronic and active denervation. He was diagnosed as having probable ALS. One year later, he presented with dysphagia and dysarthria. At that time, symptoms of FTLT were present with apathy and executive dysfunction. He is still alive 62 months after the onset of ALS. He carries the p.Ala382Thr missense mutation of the *TARDBP* gene.

COMMENT

We have described 3 families with ALS carrying the p.Ala382Thr missense mutation of the *TARDBP* gene in whom symptoms of ALS were followed by the onset of overt FTLT symptoms meeting the criteria for FTLT by Neary et al.⁹ Detailed questioning of the informants within each family failed to reveal a genealogical link between the 3 families. However, it is still possible that the families share a common haplotype and that the p.Ala382Thr mutation arose from a single mutational event many gen-




Figure 5. Right lateral (A), left lateral (B), right medial (C), left medial (D), anterior (E), posterior (F), superior (G), and inferior (H) positron emission tomography/computed tomography statistical parametric maps of patient III-1 from family C. The images show the hypometabolic areas on the left frontal and temporal associative cortices (arrows). The lesions are less prominent than those of the patients shown in Figure 3 and Figure 4. P indicates posterior; A, anterior; R, right; and L, left.

erations ago. The 3 index cases showed both behavioral and cognitive changes of FTLT as demonstrated by the neuropsychological examination and had PET/CT abnormalities, mainly in the frontal associative areas. This report expands the clinical phenotype associated with mutations in the *TARDBP* gene to include ALS-FTLT, which is considered to be a separate clinical entity from FTLT with MND.

It has been shown that patients with FTLT with and without ALS show neuronal ubiquitin-positive inclusions containing a hyperphosphorylated, ubiquitinated, and cleaved form of TAR DNA-binding protein 43.¹³ More recently, mutations in the *TARDBP* gene coding for TAR DNA-binding protein 43 have been found in individuals with ALS.^{14,15} However, no ALS cases with *TARDBP*

mutations described in these and other articles had personal or familial histories of FTLN.^{11,12,14-21} A patient with FALS and the p.Gly294Val missense mutation of the *TARDBP* gene was affected by Alzheimer-type dementia presenting with anxiety, depression, and progressive memory loss 3 years before the onset of bulbar ALS.¹² Both his siblings, affected by ALS, had no signs of cognitive impairment.

Interestingly, *TARDBP* mutations have not been found in a large series of Italian patients with familial and sporadic FTLN, including 4 with FTLN and MND.²² This finding could support the concept that ALS-FTLN and FTLN are different disorders, but it could simply be related to the fact that *TARDBP* mutations are rare in this subset of patients. In fact, recently, 2 unrelated cases carrying the c.Gly295Ser missense mutation of the *TARDBP* gene with behavioral FTLN and semantic dementia later developing MND have been described.⁷ However, in the case with a positive family history for MND (family F389), FTLN did not segregate with MND because both the affected parent and 1 sibling had MND without clinical symptoms of dementia. The other patient (family F242) was an apparently sporadic case.

The p.Ala382Thr missense mutation has been previously described in ALS, in 4 familial and 4 sporadic cases of French and Italian origin.^{11,12} In Italy, the p.Ala382Thr mutation seems to be the most frequent mutation of the *TARDBP* gene.¹² However, none of the 8 p.Ala382Thr mutation carriers described in previous articles had cognitive or behavioral impairment.^{11,12} Therefore, apparently this specific mutation of the *TARDBP* gene is not always associated with cognitive impairment. The underlying reasons for the variable phenotypic presentations of the p.Ala382Thr mutation are unclear.

In conclusion, although rare, an association between ALS and FTLN in patients carrying a mutation of the *TARDBP* gene can be found. In our families, cognitive and behavioral symptoms cosegregated with ALS.

Accepted for Publication: December 23, 2009.

Author Affiliations: Department of Neuroscience, University of Turin (Drs Chiò, Calvo, Moglia, Montuschi, and Mutani), Laboratory of Molecular Genetics, Azienda Sanitaria Ospedaliera Ospedale Infantile Regina Margherita-Sant'Anna (Dr Restagno and Mss Ossola and Brunetti), and Positron Emission Tomography Center, IRMET Diagnostic Imaging Torino (Dr Cistaro), Turin, Italy; Department of Neurology, Azienda Ospedaliera San Francesco, Nuoro, Italy (Dr Ticca); Neuromuscular Diseases Research Group, Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, Maryland (Drs Traynor and Schymick); and Department of Neurology, Azienda Universitaria-Ospedaliera di Cagliari and University of Cagliari, Cagliari, Italy (Drs Marrosu and Borghero and Ms Murru).

Correspondence: Adriano Chiò, MD, Department of Neuroscience, University of Turin, via Cherasco 15, 10126 Turin, Italy (achio@usa.net).

Author Contributions: Dr Chiò had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. *Study concept and design:* Chiò and Borghero. *Acquisi-*

tion of data: Calvo, Moglia, Ossola, Brunetti, Montuschi, Cistaro, Ticca, Murru, and Borghero. *Analysis and interpretation of data:* Chiò, Calvo, Restagno, Montuschi, Cistaro, Traynor, Schymick, Mutani, Marrosu, and Borghero. *Drafting of the manuscript:* Chiò and Traynor. *Critical revision of the manuscript for important intellectual content:* Chiò, Calvo, Moglia, Restagno, Ossola, Brunetti, Montuschi, Cistaro, Ticca, Traynor, Schymick, Mutani, Marrosu, Murru, and Borghero. *Statistical analysis:* Traynor and Schymick. *Obtained funding:* Chiò and Restagno. *Administrative, technical, and material support:* Chiò. *Study supervision:* Chiò and Borghero.

Financial Disclosure: None reported.

Funding/Support: This work was supported by grants from the Fondazione Vialli e Mauro for ALS Research Onlus, Federazione Italiana Giuoco Calcio, and Ministero della Salute (Ricerca Sanitaria Finalizzata 2007) (Dr Chiò); by the Ministero della Salute (Ricerca Sanitaria Finalizzata 2007) (Dr Restagno); and by grant Z01 AG000949-02 from the Intramural Research Program and National Institute on Aging, National Institutes of Health.

Role of the Sponsors: The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, or approval of the manuscript.

Additional Contributions: We thank the patients and their families for their participation.

REFERENCES

1. Rosen DR, Siddique T, Patterson D, et al. Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature*. 1993;362(6415):59-62.
2. Sreedharan J, Blair IP, Tripathi VB, et al. TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science*. 2008;319(5870):1668-1672.
3. Kwiatkowski TJ Jr, Bosco DA, Leclerc AL, et al. Mutations in the *FUS/TLS* gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science*. 2009;323(5918):1205-1208.
4. Vance C, Rogelj B, Hortobágyi T, et al. Mutations in *FUS*, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science*. 2009;323(5918):1208-1211.
5. Chow CY, Landers JE, Bergren SK, et al. Deleterious variants of *FIG4*, a phosphoinositide phosphatase, in patients with ALS. *Am J Hum Genet*. 2009;84(1):85-88.
6. Lomen-Hoerth C, Murphy J, Langmore S, Kramer JH, Olney RK, Miller B. Are amyotrophic lateral sclerosis patients cognitively normal? *Neurology*. 2003;60(7):1094-1097.
7. Benajiba L, Le Ber I, Camuzat A, et al; French Clinical and Genetic Research Network on Frontotemporal Lobar Degeneration/Frontotemporal Lobar Degeneration with Motoneuron Disease. *TARDBP* mutations in motoneuron disease with frontotemporal lobar degeneration. *Ann Neurol*. 2009;65(4):470-473.
8. Brooks BR, Miller RG, Swash M, Munsat TL; World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord*. 2000;1(5):293-299.
9. Neary D, Snowden JS, Gustafson L, et al. Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology*. 1998;51(6):1546-1554.
10. Strong MJ, Grace GM, Freedman M, et al. Consensus criteria for the diagnosis of frontotemporal cognitive and behavioural syndromes in amyotrophic lateral sclerosis. *Amyotroph Lateral Scler*. 2009;10(3):131-146. doi:10.1080/17482960802654364.
11. Kabashi E, Valdmanis PN, Dion P, et al. *TARDBP* mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet*. 2008;40(5):572-574.
12. Corrado L, Ratti A, Gellera C, et al. High frequency of *TARDBP* gene mutations in Italian patients with amyotrophic lateral sclerosis. *Hum Mutat*. 2009;30(4):688-694.
13. Neumann M, Sampathu DM, Kwong LK, et al. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science*. 2006;

- 314(5796):130-133.
14. Yokoseki A, Shiga A, Tan CF, et al. *TDP-43* mutation in familial amyotrophic lateral sclerosis. *Ann Neurol*. 2008;63(4):538-542.
 15. Gitcho MA, Baloh RH, Chakraverty S, et al. *TDP-43* A315T mutation in familial motor neuron disease. *Ann Neurol*. 2008;63(4):535-538.
 16. Rutherford NJ, Zhang YJ, Baker M, et al. Novel mutations in *TARDBP* (TDP-43) in patients with familial amyotrophic lateral sclerosis. *PLoS Genet*. 2008;4(9):e1000193. doi:10.1371/journal.pgen.1000193.
 17. Van Deerlin VM, Leverenz JB, Bekris LM, et al. *TARDBP* mutations in amyotrophic lateral sclerosis with TDP-43 neuropathology: a genetic and histopathological analysis. *Lancet Neurol*. 2008;7(5):409-416.
 18. Del Bo R, Ghezzi S, Corti S, et al. *TARDBP* (TDP-43) sequence analysis in patients with familial and sporadic ALS: identification of two novel mutations. *Eur J Neurol*. 2009;16(6):727-732.
 19. Daoud H, Valdmanis PN, Kabashi E, et al. Contribution of *TARDBP* mutations to sporadic amyotrophic lateral sclerosis. *J Med Genet*. 2009;46(2):112-114.
 20. Kühnlein P, Sperfeld AD, Vanmassenhove B, et al. Two German kindreds with familial amyotrophic lateral sclerosis due to *TARDBP* mutations. *Arch Neurol*. 2008;65(9):1185-1189.
 21. Kamada M, Maruyama H, Tanaka E, et al. Screening for *TARDBP* mutations in Japanese familial amyotrophic lateral sclerosis. *J Neurol Sci*. 2009;284(1-2):69-71.
 22. Gallone S, Giordana MT, Scarpini E, et al. Absence of *TARDBP* gene mutations in an Italian series of patients with frontotemporal lobar degeneration. *Dement Geriatr Cogn Disord*. 2009;28(3):239-243.

Call for Papers

Archives of Neurology will publish a special theme issue in April 2011 on infectious disease and immunology. We invite the submission of papers as Neurological Reviews, Clinical Trials, Original Communications, Case Reports, Images in Neurology, and Research Letters. Papers submitted by October 1, 2010, will have the best opportunity to be considered for this theme issue.

Announcement

Visit www.archneurology.com. As an individual subscriber, you may elect to be contacted when a specific article is cited. Receive an e-mail alert when the article you are viewing is cited by any of the journals hosted by HighWire. You will be asked to enter the volume, issue, and page number of the article you wish to track. Your e-mail address will be shared with other journals in this feature; other journals' privacy policies may differ from *JAMA & Archives Journals*. You may also sign up to receive an e-mail alert when articles on particular topics are published.