Predisplacement and Postdisplacement Factors Associated With Mental Health of Refugees and Internally Displaced Persons A Meta-analysis

Matthew Porter, PhD

Nick Haslam, PhD

HE GLOBAL REFUGEE CRISIS HAS reached dire proportions. The world's population of people forced to abandon their homes and communities because of war, political violence, or related threats includes not only those displaced outside their country of residence but also internally displaced persons, asylum seekers, stateless persons, and recently returned refugees. Although estimates of the total numbers of these persons vary, data from several international agencies indicate that these groups comprise a population that exceeded 42 million at the end of 2004.1-3 Severe and lasting psychological aftereffects have been extensively documented among them. 4-6 Additionally, the United Nations High Commissioner for Refugees alone returned approximately 15 million refugees to their countries of origin as recently as 1994 through 2003. Including all former refugees, the number of people who have been directly affected by forced displacement is estimated to exceed 1% of the world's population.8 Analyses presented in this article specify relationships between subcategories of displaced persons and mental health outcomes. Because of important common-

See also Patient Page.

CME available online at www.jama.com

Context The global refugee crisis requires that researchers, policymakers, and clinicians comprehend the magnitude of the psychological consequences of forced displacement and the factors that moderate them. To date, no empirical synthesis of research on these issues has been undertaken.

Objective To meta-analytically establish the extent of compromised mental health among refugees (including internally displaced persons, asylum seekers, and stateless persons) using a worldwide study sample. Potential moderators of mental health outcomes were examined, including enduring contextual variables (eg, postdisplacement accommodation and economic opportunity) and refugee characteristics.

Data Sources Published studies (1959-2002) were obtained using broad searches of computerized databases (PsycINFO and PILOTS), manual searches of reference lists, and interviews with prominent authors.

Study Selection Studies were selected if they investigated a refugee group and at least 1 nonrefugee comparison group and reported 1 or more quantitative group comparison on measures of psychopathology. Fifty-six reports met inclusion criteria (4.4% of identified reports), yielding 59 independent comparisons and including 67 294 participants (22 221 refugees and 45 073 nonrefugees).

Data Extraction Data on study and report characteristics, study participant characteristics, and statistical outcomes were extracted using a coding manual and subjected to blind recoding, which indicated high reliability. Methodological quality information was coded to assess potential sources of bias.

Data Synthesis Effect size estimates for the refugee-nonrefugee comparisons were averaged across psychopathology measures within studies and weighted by sample size. The weighted mean effect size was 0.41 (SD, 0.02; range, -1.36 to 2.91 [SE, 0.01]), indicating that refugees had moderately poorer outcomes. Postdisplacement conditions moderated mental health outcomes. Worse outcomes were observed for refugees living in institutional accommodation, experiencing restricted economic opportunity, displaced internally within their own country, repatriated to a country they had previously fled, or whose initiating conflict was unresolved. Refugees who were older, more educated, and female and who had higher predisplacement socioeconomic status and rural residence also had worse outcomes. Methodological differences between studies affected effect sizes.

Conclusions The sociopolitical context of the refugee experience is associated with refugee mental health. Humanitarian efforts that improve these conditions are likely to have positive impacts.

JAMA. 2005;294:602-612

www.jama.com

Author Affiliations: Columbia Presbyterian Medical Center, New York, NY (Dr Porter); University of Melbourne, Melbourne, Australia (Dr Haslam).

Corresponding Author: Matthew Porter, PhD,

Columbia Presbyterian Medical Center, Department of Pediatric Psychiatry, 622 W 165th St, Sixth Floor N, New York, NY 10032 (mattporteremail@yahoo.com).

602 JAMA, August 3, 2005—Vol 294, No. 5 (Reprinted)

 $\hbox{$@2005$ American Medical Association. All rights reserved.}\\$

alities, the various subcategories have herein been subsumed under the general term *refugee* for simplicity.

Research on refugee mental health has often proceeded from a life events model^{9,10} in which psychopathology is understood as a posttraumatic reaction to the acute stressor of war.11 Research in this tradition has established high rates of psychological disturbance among persons exposed to traumas through war and disaster. 12-14 Theorists have recently argued that this model fails to capture important dimensions of the refugee experience, 15,16 which is characterized by multiple events occurring in multiple contexts that persist over time. In addition, refugees are vulnerable to multiple dimensions of psychopathology beyond those that are narrowly posttraumatic. 16-18

Unlike survivors of most discrete traumatic events, refugees experience diverse stressors that accumulate over the preflight, flight, exile, and resettlement/repatriation periods. ¹⁹ Despite the historical focus on the acute stressors of war, the enduring contextual postmigration stress that refugees face—including marginalization, socioeconomic disadvantage, acculturation difficulties, loss of social support, and "cultural bereavement" —must be recognized. Such persisting forms of adversity reflect the broader sociopolitical context of the refugee experience. ^{15,21}

Research on refugee mental health is fraught with practical obstacles. Populations are often physically, linguistically, and culturally inaccessible to researchers,22 and humanitarian aid usually has higher priority than scientific investigation. Studies are often exploratory and methodologically compromised, and the specificity of local circumstances makes it difficult to draw generalized conclusions.23 As a result, research synthesis is needed to establish the magnitude of the mental health consequences of forced displacement and the variables that moderate these consequences. We have previously reported a meta-analytic investigation of refugee mental health,24 but it was restricted to the former Yugoslavian Republic. We therefore conducted a metaanalysis that comprehensively examined studies of populations displaced in conflicts distributed throughout the world and across 5 decades. The metaanalysis tested several potential moderators of refugee mental health, including contextual variables subsequent to the acute stress of war.

Six postdisplacement condition moderators were investigated. Type of accommodation was examined in view of evidence that long-term institutional housing promotes dependency and demoralization.^{25,26} Postdisplacement economic opportunity was studied as a factor expected to be associated with better refugee outcomes.27 We predicted that greater cultural access (ie, the extent to which individuals are free to practice or engage in cultural, social, and religious customs and have access to the cultural institutions necessary for such practice), internal displacement (ie, being displaced within country of origin vs external displacement), and being repatriated would be associated with better outcomes because acculturative stress and cultural dislocation should have negative effects on mental health.^{20,24,28} We also predicted that the status of the conflict would be another contextual condition affecting refugees' mental health, with ongoing (vs resolved) conflicts being associated with worse outcomes.

In addition to these contextual variables, several refugee characteristics were investigated as potential moderators. The association between age and severity of posttraumatic response has been controversial, 13,29 with some evidence indicating worse outcomes for older people,30-33 some indicating greater vulnerability in children, 17,34-35 and some mixed or null. 38-40 Refugee sex was also examined, 17,41 with some evidence suggesting worse mental health outcomes for female refugees^{38,42} and some indicating complex and culturally variable associations.²³ Predisplacement urban vs rural residence, region of origin (ie, Africa, Asia, Europe, Latin America, or Middle East), educational level, and predisplacement socioeconomic status were also investigated.

We also inquired about the moderating role of methodological characteristics of studies that might serve as confounders. We examined whether methodologically stronger studies⁴¹ and those whose control groups had greater exposure to adversity estimated less negative relative mental health outcomes for refugees. Finally, we assessed whether the time elapsed since displacement was associated with smaller effects, consistent with the decay of stress responses. ⁴³⁻⁴⁶

METHODS

Selection of Primary Studies

A search of 2 databases was undertaken at the end of 2002 for all published English-language research reports that (1) investigated a refugee group (including all categories within UNHCR's population of concern¹) and at least 1 nonrefugee comparison group and (2) reported 1 or more quantitative estimates of the relationship between forced displacement and indices of mental health (eg, symptom questionnaires, interviewbased psychiatric diagnoses, maladaptive behavior checklists, neuropsychological tests, personality inventories, and self-esteem and well-being scales).

Searches were conducted using PsycINFO, a broad psychological database, and PILOTS, a comprehensive trauma-specific database maintained by the US Department of Veterans Affairs National Center for PTSD.⁴⁷ PILOTS was queried using a Boolean operator (refugee* OR displaced) and PsycINFO with narrower criteria: key word= {[(refugee*) OR (displaced person*) OR (displaced people)] AND [(control*) OR (compar*) OR ("nonrefugees") OR ("nonrefugee") OR (nonrefugee*) OR ("non-displaced") OR (nondisplaced)]}. No restrictions were placed on the time period covered by the search, so all indexed literature from 1871-2002 for PILOTS and from 1840-2002 for PsycInfo was included.

The search of PILOTS produced 1010 citations, and PsycINFO yielded 273 additional nonoverlapping records. All abstracts were read and those that unambiguously failed to meet inclusion criteria

were eliminated. Full texts were obtained for the 141 remaining articles. Manual searches through reference lists of identified articles were conducted and

prominent authors were contacted informally but no additional published references were obtained. After eliminating all redundant research reports, 56

reports⁴⁸⁻¹⁰³ (published from 1959 through 2002; median year of publication, 1996) were identified as meeting inclusion criteria (TABLE 1 and TABLE 2).

Author	No. of Participants	Sample Composition	Measures
Adams et al, ⁴⁸ 1980	201	Cuban refugee boys vs nonrefugee American black boys	Louisville Aggression Survey Schedule
Ahmad et al, ⁴⁹ 2000	64	Kurdish refugee children vs nonrefugee Swedish children	CBC Child Report, CBC Parent Report
Al-Eissa,50 1995	226	Kuwaiti refugee children vs nonrefugee Saudi children	Symptom questionnaire, interview
Bauer et al, ⁵¹ 1994	168	East German refugees vs nonrefugee West Germans	Clinical interview
Bayard-Burfield et al, ⁵² 2001	644 1139 3198	Iranian refugees vs nonrefugee Turkish migrant laborers Chilean refugees vs nonrefugee Polish migrant laborers Kurdish refugees vs nonrefugee Swedes	Clinical interview Clinical interview Clinical interview
Beiser, ⁵³ 1987	1644	Southeast Asian refugees vs nonrefugee Canadians	Ad hoc questionnaire
Carballo et al,54 1996	5242	Bosnian refugees vs nonrefugee Bosnians	Clinical interview
Cervantes et al, ⁵⁵ 1989	587	Central American refugees vs nonrefugee Central American immigrants vs nonrefugee Mexican immigrants vs nonrefugee US residents	Symptom questionnaire
Clark et al, ⁵⁶ 1996	472	Central American refugees vs nonrefugee Mexican immigrants	Minnesota Multiphasic Personality Inventory
Dube, ⁵⁷ 1968	29 468	Punjabi refugees vs nonrefugee Indians	Clinical interview
Dunnigan et al,58 1993	1019	Hmong refugees vs nonrefugee US residents	Youth Development Study Questionnaire
Eitinger, ⁵⁹ 1959	3758	Central and Eastern European refugees vs nonrefugee Norwegians	Clinical interview and psychiatric history
El Habir et al,60 1994	355	Lebanese refugee children vs nonrefugee Middle Eastern children vs nonrefugee US children	CBC Teacher Report
Fazel et al,61 1988	125	Tibetan refugees vs nonrefugee Indians	Quality of Life Scale
Folnegovic-Smalc et al, ⁶² 1997	1076	Croatian IDPs vs nonrefugee Croatians	Neuropsychological battery and clinical interview
Goenjian et al,63 1994	150	Armenian refugees vs nonrefugee Armenians whose homes were destroyed by earthquake vs nonrefugee Armenians with intact homes	PTSD Reaction Index
Herceg et al, ⁶⁴ 1996	56	Croatian and Bosnian IDP children vs nonrefugee Croatian children	Life Orientation Test, IES
Hourani et al,65 1986	5788	Lebanese IDPs in camps vs Lebanese IDPs in empty apartments vs Lebanese IDPs displaced 5 years earlier vs nonrefugee Lebanese under siege	Symptom questionnaire
Howard and Hodes, 66 2000	135	Refugees from diverse backgrounds vs nonrefugee matched immigrants vs nonrefugee British	Clinical interview
Jensen et al, ⁶⁷ 1989	93	Middle Eastern refugees vs nonrefugee Middle Eastern immigrants	Clinical interview
Klimidis et al,68 1994	156	Vietnamese refugee boys vs nonrefugee immigrant boys vs nonrefugee Australian boys	Symptom questionnaire based on HSCL
	289	Vietnamese refugee girls vs nonrefugee immigrant girls vs nonrefugee Australian girls	Symptom questionnaire based on HSCL
Kocijan-Hercigonja et al, ⁶⁹ 1996	321	Wounded Croatian IDP children vs wounded nonrefugee Croatian children	Clinical interview
Kocijan-Hercigonja et al, ⁷⁰ 1998	105	Bosnian refugee children vs Croatian IDP children vs nonrefugee Croatian children	Schoolagers' Coping Strategies Questionnaire, Questionnaire of Psychosomatic Symptoms, Anxiety Scale, Depression Scale, Psychosocia Adjustment Questionnaire
Kocijan-Hercigonja et al, ⁷¹ 1996	62	Croatian IDPs vs nonrefugee Croatians	Cornell Index
Kondic and Mavar, ⁷² 1992	142	Croatian IDPs vs nonrefugee Croatians	State-Trait Anxiety Inventory
Kutorovoo et al 73	124	Creation IDD obildren va poprofugeo Creation obildren	IEQ

Abbreviations: CBC, Child Behavior Checklist; HSCL, Hopkins Symptom Check List; IDP, internally displaced person; IES, Impact of Event Scale; PTSD, posttraumatic stress disorder.

Croatian IDP children vs nonrefugee Croatian children

134

IES

Kuterovac et al,73

Author	No. of Participants	Sample Composition	Measures
Laor et al, ⁷⁴ 1996	454	Israeli IDP women and children vs nonrefugee Israeli women and children threatened by war events vs nonrefugee, nonexposed Israeli women and children	CBC, Symptom Checklist-90-R
Lopes Cardozo et al, ⁷⁵ 2000	1272	Kosovar IDP Albanians vs repatriated Albanian refugees vs nonrefugee Kosovar Albanians	General Health Questionnaire-28, Medica Outcomes Study Short-Form, HTQ
oughry and Flouri, ⁷⁶ 2001	455	Repatriated Vietnamese refugees vs nonrefugee Vietnamese	Youth Self-Report, Perceived Self-Efficac Scale
Macksoud and Aber, ⁷⁷ 1996	115	Lebanese IDP children vs repatriated Lebanese refugee children vs nonrefugee Lebanese children	Child Behavior Inventory, Post-Traumatic Stress Reaction Checklist
McCloskey et al,78 1995	70	Central American refugee women vs nonrefugee Mexican immigrant women	PTSD symptom inventory
	70	Central American refugee children vs nonrefugee Mexican immigrant children	PTSD symptom inventory, CBC
Myers et al, ⁷⁹ 1987	88	Vietnamese refugees vs nonrefugee Chinese immigrants and US citizens	Adult Fear Schedule
Onwumere et al,80 2002	112	African refugee women with human immunodeficiency virus infection vs nonrefugees with chronic illness	Short Form Health Survey
Paardekooper et al, ⁸¹ 1999	396	Sudanese refugee children vs nonrefugee Ugandan children	Kidcope, Reporting Questionnaire for Children–Parent Form, General Health Questionnaire, cartoon-based interview
Pernice and Brook,82 1994	186	Southeast Asian refugees vs nonrefugee Pacific Islander immigrants	HSCL-25
Punamaki,83 1990	239	Palestinian refugee women vs nonrefugee Palestinian women in the Occupied Territories vs nonrefugee Palestinian women in Israel	Structured interview, Multiple Affect Adjective Checklist
Quta and El Sarraj,84 1993	160	Palestinian refugees in camps vs resettled Palestinian refugees vs Palestinian town refugees vs nonrefugee Palestinians	Psychosomatic Symptoms Test, Taylor Manifest Anxiety Scale, Rigidity Test
Rasanen,85 1989	523	Repatriated Finnish refugees vs nonrefugee Swedes	Symptom questionnaire
Rasekh et al, ⁸⁶ 1998	160	Afghan refugee women vs nonrefugee Afghan women under the Taliban	HSCL-25, symptom questionnaire
Reeler, ⁸⁷ 1994	208	Mozambican refugees vs nonrefugee Zimbabweans	Self-Reporting Questionnaire, clinical interview
Riolli et al, ⁸⁸ 2002	112	Kosovar Albanian refugees vs nonrefugee Albanian immigrants vs nonrefugee Albanians	BSI
Roglic et al,89 1993	88	Croatian IDPs with diabetes vs nonrefugee Croatians with diabetes	Structured interview, BDI
Roncevic-Grzeta et al,90 2001	109	Tortured Croatian IDP war veterans vs nontortured Croatian IDP war veterans vs nonrefugee, nontortured, noncombat Croatians	Symptom questionnaire, clinical interview BDI, Hamilton Depression Inventory
Rousseau et al, ⁹¹ 2000	225	Cambodian adolescent refugees vs Central American adolescent refugees vs nonrefugee Canadian adolescents	CBC Parent Report, Youth Self-Report
Sabioncello et al, ⁹² 2000	34	Croatian IDP women vs nonrefugee Croatian women	Cornell Index
Silove et al,93 1998	196	Asylum-seeking Tamil refugees vs permanently resettled Tamil refugees vs nonrefugee Tamil immigrants	HSCL-25, HTQ
Smither and Rodriguez-Giegling, ⁹⁴ 1979	89	Vietnamese refugees vs Laotian refugees vs nonrefugee US residents	Marginality Scale, State-Trait Anxiety Inventory
Gundelin-Wahlsten et al, ⁹⁵ 2001	85	Kurdish refugees vs nonrefugee Swedes	HTQ
Sundquist, ⁹⁶ 1993	1891	Latin American refugees vs nonrefugee European migrant laborers vs nonrefugee Swedes	Swedish Annual Level-of-Living Survey
Sundquist and Johansson,98 1996	1219	Repatriated Latin American refugees vs nonrefugee Swedes	Swedish Annual Level-of-Living Survey
hulesius and Hakansson, ⁹⁹ 1999	593	Bosnian refugees vs nonrefugee Swedes	Posttraumatic Symptom Scale
ousignant et al,100 1999	406	Adolescent refugees from 35 countries vs nonrefugee Canadian adolescents	Diagnostic Interview Schedule for Childre version 2.25
Vong-Rieger and Quintana, ¹⁰¹ 1987	164	Southeast Asian refugees vs nonrefugee Anglo-American US residents	Multicultural Acculturation Scale
oung and Evans, 102 1997	120	Salvadorean refugees vs nonrefugee Canadians	BSI, Quality of Life Scale, Satisfaction Wit Life Scale
Zivcic, ¹⁰³ 1993	638	Croatian IDP children vs nonrefugee Croatian children vs prewar archival records of nonrefugee Croatian children	Emotion Scale Child Depression Inventor

Two of these reports provided complementary information about the same comparison and 3 reported multiple independent comparisons, bringing the total meta-analytic sample to 59 independent comparisons (studies), detailed in Table 1 and Table 2. Of these, 20 used multiple comparison groups,

yielding 85 stochastically dependent comparisons from which subsets of statistically independent comparisons were selected for analysis. The studies contained 67 294 participants (22 221 refugees and 45 073 nonrefugees), with a range of 34 to 29 468 (median, 164) participants per study. Together, the stud-

ies yielded 190 effect sizes, using primarily self- and parent-report questionnaires (83.2%), including standard104-106 and ad hoc instruments and structured (3.7%) and unstructured (12.6%) interviews.

Study Coding

Coding of the studies (M.P.) was conducted in accordance with established procedures, 107,108 encompassing report and study characteristics (eg, methods), participant characteristics (eg. attributes and circumstances of each independent sample), and statistical outcome information. Information pertaining to all study groups and mental health measures was coded. A coding manual was developed a priori and revised iteratively. 108 Coded variables of interest are presented in TABLE 3. After extensive spot checking, a randomly selected sample of 6 reports was blindly recoded (M.P.), yielding perfect coding agreement, indicating strong reliability.

Effect Size Computation and Statistical Analysis

Two layers of dependencies existed in the coded database. When multiple measures were used on a single sample, their effect sizes were averaged. 107,108 When multiple comparison groups were used, only 1 effect size from each study was allowed in any given analysis, as described below.

All statistical analyses followed established procedures, 107,108 with the parameter of interest, δ , quantifying the degree of overlap between refugee and nonrefugee distributions on mental health measures. 109 The δ was estimated by the Cohen d, the effect size equal to the difference between 2 group means divided by the pooled standard deviation. Direct effect size calculation from published means and standard deviations was possible only in roughly half of the studies. For the 24 studies (41%) reporting categorical data (eg, group differences in prevalence of a diagnosis), effect sizes were calculated as the difference between the probits, or cut points on the nor-

Variables	Measurement Level	Coding Categories
stdisplacement conditions		ocaming categories
Accommodation	Ordinal	Institutional, private (temporary), private (permanent/semipermanent)
Cultural access	Ordinal	Freedom to practice culture or origin: highly restricted, moderately restricted, unrestricted
Economic opportunity	Ordinal	Highly restricted, moderately restricted unrestricted (loss of status), unrestricted (no substantial loss of status)
Locus of displacement	Nominal-dichotomous	External or internal to refugee country of origin
Repatriation status	Nominal-dichotomous	Refugees repatriated or not repatriated
Stage of conflict	Nominal-dichotomous	Ongoing, postconflict
fugee characteristics		01.11/ 1.1
Age	Interval	Child/adolescent, adult, senior (≥65 y)
Sex	Interval	% Female
Community type	Nominal-dichotomous	Rural (including villages, small towns), urban
Region of origin	Nominal-polychotomous	Africa, Asia, Europe, Latin America, Middle East
Education	Ordinal	Mostly primary only, mostly secondary only, substantial postsecondary
Predisplacement socioeconomic status	Ordinal	Primarily lower/working class, primarily middle class
ethodological characteristics Time since displacement	Interval	Mean number of years between displacement and data collection
Nonrefugee displacement	Nominal-dichotomous	Nondisplaced nonrefugees, nonrefuge immigrants
Nonrefugee violence	Ordinal	No exposure to organized violence, some direct exposure, tortured
Nonrefugee war experience	Ordinal	No, past, or current war experience
Research setting	Nominal-dichotomous	Academic, nonacademic
Publication date	Ordinal	Decade of publication (1950s-2000s)
Effect size primacy	Nominal-dichotomous	Mental health differences between groups as primary or secondary research focus
Primary researcher profession	Nominal-polychotomous	Psychiatrist, psychologist, other menta health professional, student, layperson
Primary researcher qualification	Dichotomous	Professional researcher, student or layperson
Research setting	Dichotomous	Academic (university or university-affiliated hospital), nonacademic
Measure quality ¹⁶	Dichotomous	High, low
Sample randomization	Dichotomous	Randomized, not randomized
Study blinding	Dichotomous	Blind, not blind
Researcher ethnicity	Nominal-polychotomous	Same as refugees, same broad cultura category, different ethnicity, different broad cultural category


606 JAMA, August 3, 2005-Vol 294, No. 5 (Reprinted)

mal distribution below which the 2 proportions fell. For studies failing to report pertinent statistical data, d was calculated using the other standard procedures for effect size estimation. 108 Each independent effect size estimate was weighted with the inverse of its variance (roughly in proportion to its study's sample size). 107 Effect sizes derived from studies with less than 20 participants were corrected for smallsample bias according to a standard algorithm.110 Effect sizes were windsorized (artificially adjusted) to 2 SD units from the mean and inverse variance weights to the nearest cluster so that extreme outliers would not eclipse observed variance in the rest of the distribution. 108,111

Homogeneity analysis using the Q statistic112 was performed to determine if effect size variability exceeded that due to sampling error, implying the systematic effect of moderator variables. Analysis indicated significant variability ($Q_{\text{total}} = 1329.73$; P < .001): 96% of the variance in effect size point estimates was due to heterogeneity rather than sampling error $(I^2=0.96)$. ¹¹³ Proposed moderator variables were examined by testing the probability of their systematic association with observed variance in the effect size distribution. Variables differing only between studies were investigated by averaging effect sizes across multiple comparison groups within individual studies. Some variables also varied between groups within multiple comparison group studies. For analyses of these variables, data sets consisted of effect sizes from all codable single comparison group studies plus 1 effect size from each codable multiple comparison group study, selected using a standard algorithm to yield maximally equal representation of all levels of the grouping variable, a requirement to optimize statistical power. 108 When only 1 group from a multiple comparison group study contained the required level of the variable, its effect size was included in the analysis and the remaining groups were excluded. When more than 1 group from such a study contained the required level of the variable, effect sizes from those groups were combined in a weighted average.

Once the appropriate data sets were constructed, effect sizes were grouped according to the levels of each variable. Variability within and between levels was analyzed with homogeneity tests. Q statistics were calculated for each level and were summed for Q_{within} . The difference between Q_{within} and Qtotal tests the significance of a potential moderator variable's effect on the overall variance of effect sizes.

Figure 1. Effect Sizes of Primary Studies



Error bars indicate 95% confidence intervals; dashed line, overall weighted mean effect size.

Publication bias in the meta-analytic sample was assessed as a potential threat to validity¹¹⁴ using funnel plots.^{115,116} Publication bias implies that studies with less statistical power are less likely to be published than larger, more powerful studies and, hence, may be underrepre-

Table 4. Moderating Role of Postdisplacement Conditions on Refugee Mental Health

Moderator Variables	No. of Studies	Effect Size (95% Confidence Interval)
Accommodation Institutional	13	0.65 (0.58-0.71)
Private, temporary	9	0.67 (0.60-0.75)
Private, permanent	27	0.37 (0.33-0.40)
Cultural access Moderately restricted	27	0.50 (0.45-0.54)
Unrestricted	32	0.46 (0.42-0.50)
Economic opportunity Highly restricted	8	1.06 (0.97-1.14)
Moderately restricted	13	0.45 (0.37-0.53)
Unrestricted, status loss	32	0.43 (0.39-0.46)
Unrestricted, no status loss	6	0.27 (0.19-0.34)
Locus of displacement External	44	0.39 (0.35-0.42)
Internal	15	0.65 (0.60-0.71)
Repatriation status Not repatriated	54	0.44 (0.41-0.47)
Repatriated	5	0.82 (0.73-0.92)
Stage of conflict Ongoing	41	0.60 (0.56-0.64)
Postconflict	15	0.32 (0.28-0.36)

sented. Smaller samples should yield more variable estimates of the population parameter than larger studies, and if publication bias exists, their estimates should also tend to be larger. Funnel plots indicated that a relatively large number of small studies were published and revealed no relationship between effect size and sample size. The Begg test 117,118 revealed no association between effect size estimates and their variances (ρ =0.19; P>.05), indicating that there was no significant publication bias.

RESULTS

The weighted mean effect size *d* across the 59 studies was 0.41 (range, –1.36 to 2.91; SE, 0.01) (FIGURE 1). Across studies, despite large variations in effect sizes, refugees scored 0.41 SDs lower on indices of mental health than nonrefugees.

Systematic associations between the proposed moderator variables and effect size variance were investigated and are presented separately for postdisplacement conditions, refugee characteristics, and study methodological characteristics.

Postdisplacement Conditions

Effect size variation as a function of postdisplacement conditions is summarized in TABLE 4, which demonstrates several significant moderators of the mental health of refugees relative to nonrefugees. As expected, postdisplacement accommodation was asso-


ciated with mental health (Q=86.82; P<.001; R^2 =0.06) (FIGURE 2), with refugees resettled in permanent, private accommodations having significantly better mental health than those resettled in institutional and temporary private accommodations. Economic opportunity (right to work, access to employment, maintenance of socioeconomic status) had a linear relationship with better mental health (Q=217.96; P<.001; R^2 =0.13) (Figure 2). Access to cultural practices was not associated with mental health (Q=1.45; P=.23; R^2 =0.00).

The position of refugees vis-à-vis their initiating conflicts was also significantly associated with mental health. Internally displaced persons scored lower on mental health indices than externally displaced refugees (Q=65.47; P<.001; $R^2=0.05$) (Figure 2). Repatriated refugees scored lower than those in exile, many of whom had been permanently resettled (Q=52.79; P<.001; $R^2=0.04$). Refugees from conflicts that remained ongoing scored lower than those from conflicts that had been resolved (Q=92.00; P<.001; $R^2=0.07$).

Predisplacement Refugee Characteristics

All of the coded refugee characteristics were significantly associated with the mental health of refugees relative to nonrefugees (TABLE 5). Age had a strong association (Q = 227.28; P < .001; $R^2 = 0.13$). Child and adolescent refugees had better relative outcomes than adults. Those aged 65 years or older scored lower on mental health than adults younger than 65 years, although only 2 studies assessed this population. Refugee sex also had a weak but significant moderator effect (β =0.15; R^2 =0.02; P<.001). Studies containing a higher proportion of female refugees indicated poorer mental health outcomes. Refugees displaced from rural areas had poorer outcomes than those from urban areas (Q = 17.64; P < .001; $R^2 = 0.03$). Region of origin was also significantly associated with mental health (Q = 505.96; P < .001) and was probably confounded by comparison group region of origin. Refugees from

Figure 2. Effect Size Means Associated With Type of Accomodation, Degree of Economic Opportunity, and Locus of Displacement at the Time of Measurement



Error bars indicate 95% confidence intervals.

608 JAMA, August 3, 2005—Vol 294, No. 5 (Reprinted)

Europe (most often compared with nonrefugee residents of peaceful European countries) had the poorest relative outcomes, and those from Asia and the Middle East had mental health outcomes that were the least different from their comparison groups, which were most often nonrefugee residents of the respective regions, often developing countries with histories of civil unrest. More-educated refugees scored lower on mental health indices than lesseducated refugees (Q=319.68; P<.001; R^2 =0.28), as did those with higher predisplacement socioeconomic status $(Q=177.71; P<.001; R^2=0.16).$

Methodological Characteristics

Several study characteristics were associated with effect size (TABLE 6). Greater time between date of study and time of displacement was associated with better mental health for refugees relative to nonrefugees ($\beta = -0.28$; P < .001; R^2 =0.08), although durations were often short (median, 2 years; range, 1 month to 41 years after displacement) and more research is needed on longterm effects. Refugees had less negative outcomes compared with nonrefugees who had undergone displacement (ie, immigrants) vs nonrefugees who had not been displaced (Q = 23.14; P < .001; R^2 =0.01). This expected effect also held true for nonrefugee exposure to violence (Q=106.82; P<.001; $R^2=0.03$) and nonrefugee war experience (Q=14.24; P < .001; $R^2 = 0.02$). For these variables, the highest levels of contamination in the comparison groups were clearly associated with smaller effect sizes. Refugees showed mental health comparable with nonrefugees who were tortured or living in active war zones.

As expected, higher methodological quality was associated with slightly smaller effect sizes. 41 An index based on the sum of 5 dichotomously coded indicators of quality: study, blindness, sample randomization, researcher qualifications (psychiatrist, psychologist, or other mental health professional vs student or layperson), type of setting (academic [universities or universityaffiliated hospitals] vs nonacademic), and

quality of study measurement instruments (surveys and scales)16 revealed a weak negative association between methodological quality and effect size $(\beta = -0.10; P < .001; R^2 = 0.01).$

Publication decade was weakly associated with effect sizes (Q=302.31; P < .001; $R^2 = 0.01$) but lacked a discernible pattern. Studies for which comparing mental health in refugees and nonrefugees was not a primary focus yielded larger effect sizes (Q=12.98; P<.001; R^2 =0.01), providing additional evidence against the influence of publication bias in the meta-analytic sample. 107 Proximity of researcher ethnicity to refugee ethnicity was not significantly associated with effect size (Q=5.3718; P=.07; $R^2=0.00$), although the data demonstrated a marginal trend toward researchers finding larger effects when examining refugees of ethnicities dissimilar to their own.

Given the collinearity of some coded variables in the meta-analysis, the 10 substantive variables (ie, postdisplacement conditions and refugee characteristics) measured dichotomously or ordinally were factor analyzed to yield statistically independent predictors of effect size. Three factors emerged: material welfare (postdisplacement private accommodation and economic opportunity, high predisplacement socioeconomic status, conflict resolved), predisplacement cultural capital (greater education and age, predisplacement urban residence), and distance from original conflict (external displacement, not being repatriated, greater cultural freedom). All 3 factors independently predicted effect size in a least-squares regression analysis (β =-0.16, 0.22, and -0.14, respectively; all P < .05; $R^2 = 0.33$). Better relative mental health among refugees was associated with greater material welfare, lesser predisplacement cultural capital, and greater distance from the original conflict.

COMMENT

This meta-analysis of refugee mental health clarifies the magnitude and determinants of the psychological consequences of the refugee experience. The magnitude of these consequences, relative to those experienced by control populations, is within the range considered medium-sized. 119 It remains substantial even when the comparison group has been displaced or directly exposed to war and violence, attesting to the depth of adversity that refugees experience.

	naracteristics on Refugee Mental Health		
Moderator Variables	No. of Studies	Effect Size (95% Confidence Interval)	
Age	110. 01 0144100	(co /c commonico miorval)	
Children/adolescents	17	0.28 (0.21-0.35)	
Adults	35	0.53 (0.49-0.57)	
Seniors (≥65 y)	2	1.21 (1.08-1.33)	
Community type Rural	39	0.50 (0.47-0.54)	
Urban	10	0.32 (0.25-0.40)	
Region of origin Africa	3	0.61 (0.44-0.78)	
Latin America	10	0.46 (0.39-0.53)	
Middle East	12	0.18 (0.12-0.24)	
Asia	14	0.17 (0.12-0.21)	
Europe	18	0.82 (0.78-0.86)	
Education Mostly primary only	20	0.25 (0.19-0.31)	
Mostly secondary only	11	0.34 (0.27-0.42)	
Substantial postsecondary	15	0.87 (0.82-0.92)	
Predisplacement socioeconomic status Primarily working class	26	0.34 (0.30-0.38)	
Primarily middle class	13	0.91 (0.83-0.99)	

As predicted, postdisplacement conditions were associated with mental health outcomes. Materially secure conditions, indexed by economic opportunities and permanent private accommodation, were associated with superior outcomes. By implication, psychopathology among refugees is not an inevi-

table posttraumatic consequence of acute wartime stress but reflects contextual factors that can be significantly remediated by generous material support on the part of governments and agencies. Similarly, the status of the initiating conflict appears to be a significant feature of the sociopolitical context, because

ology among refugees is not an inevi- of the sociop

Moderator Variables	No. of Studies	Effect Size (95% Confidence Interval)
Nonrefugee displacement		(0070 00111101100 1111011011)
Nondisplaced	42	0.50 (0.47 to 0.53)
Displaced	16	0.33 (0.26 to 0.39)
Nonrefugee violence No direct exposure	45	0.52 (0.49 to 0.55)
Some direct exposure	12	0.62 (0.53 to 0.72)
Tortured	1	0.01 (-0.09 to 0.10)
Nonrefugee war experience None	31	0.43 (0.39 to 0.48)
Past	22	0.61 (0.56 to 0.65)
Current	6	0.11 (0.02 to 0.19)
Research setting Academic	44	0.35 (0.31 to 0.38)
Nonacademic	15	0.80 (0.74 to 0.85)
Publication decade 1950s	1	0.67 (0.60 to 0.73)
1960s	1	0.23 (0.18 to 0.29)
1970s	1	0.72 (0.19 to 10.24)
1980s	8	0.05 (-0.01 to 0.11)
1990s	35	0.71 (0.67 to 0.74)
2000s	13	0.41 (0.35 to 0.47)
Effect size primacy Primary focus	46	0.45 (0.42 to 0.48)
Secondary focus	13	0.58 (0.52 to 0.64)
Researcher profession Psychiatrist	25	0.53 (0.50 to 0.56)
Psychologist	14	0.43 (0.33 to 0.54)
Mental health professional	12	0.40 (0.36 to 0.45)
Student	4	0.00 (-0.09 to 0.10)
Layperson	2	1.44 (1.27 to 1.62)
Researcher qualification Professional researcher	50	0.49 (0.46 to 0.52)
Student or layperson	9	0.35 (0.28 to 0.42)
Research setting Academic	44	,
	15	0.35 (0.31 to 0.38)
Nonacademic Sample randomization		0.80 (0.74 to 0.85)
Randomized	30	0.52 (0.46 to 0.57)
Not randomized	29	0.46 (0.43 to 0.48)
Study blinding Blind	57	0.49 (0.46 to 0.52)
Not blind	2	-0.33 (-0.62 to 0.05)
Measure quality ¹⁶ High	27	0.66 (0.59 to 0.72)
Low	32	0.44 (0.41 to 0.46)
Researcher ethnicity Same as refugees	18	0.46 (0.40 to 0.51)
Close to refugees	7	0.43 (0.39 to 0.47)
Dissimilar to refugees	34	0.50 (0.46 to 0.53)

refugees had much better outcomes when the conflict that displaced them had been resolved.

Other postdisplacement conditions had unexpected effects. Acculturative stress, cultural dislocation, and bereavement^{20,24,28} were predicted to yield better outcomes for refugees who had unrestricted cultural access and were internally displaced and repatriated, but none of these predictions were supported. Externally displaced and unrepatriated refugees had relatively positive mental health outcomes, a finding that runs counter to the emphasis on repatriation in the humanitarian aid community. Repatriation, internal displacement, and cultural continuity are likely to have positive implications, but these may be overshadowed by the harmful effects of unstable political and economic circumstances within the nation of origin.

Refugee characteristics also moderated mental health outcomes. Consistent with some past work on posttraumatic responses, 38,42 female refugees had slightly worse mental health outcomes than male refugees. The apparently greater resilience of younger refugees supports previous findings,30-33 implying that children and adolescents are less affected by the enduring stresses of displacement.²⁹ However, higher levels of education and socioeconomic status before displacement, considered by some to have buffering functions,19 were associated with worse mental health outcomes in the analysis. Greater predisplacement intellectual and economic resources may imply a greater subsequent loss of status rather than a protective effect on refugees against their predicament. Refugee mental health outcomes were associated with region of origin, although no predictions were tested. European refugees had the poorest relative outcomes, perhaps in part because their comparison groups tended to be residents of peaceful and economically privileged countries. Refugee outcomes were also relatively poor in Africa, but because only 2% of the metaanalytic sample was drawn from Africa compared with 23.6% of the UNHCR's

610 JAMA, August 3, 2005—Vol 294, No. 5 (Reprinted)

population of concern, the continent's refugee crises demand more investigation. At present, the global distribution of refugees is not adequately represented in the mental health literature.

In sum, this study supports the role of enduring contextual factors before and after displacement as moderators of mental health among the world's refugees. The psychological aftereffects of displacement by war cannot be understood simply as the product of an acute and discrete stressor, but depend crucially on the economic, social, and cultural conditions from which refugees are displaced and in which refugees are placed. 15,22

Author Contributions: Dr Porter had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Porter, Haslam.

Acquisition of data: Porter.

Analysis and interpretation of data: Porter, Haslam. Drafting of the manuscript: Porter, Haslam.

Critical revision of the manuscript for important intellectual content: Porter, Haslam.

Statistical analysis: Porter.

Administrative, technical, or material support: Porter, Haslam

Study supervision: Haslam.

Financial Disclosures: None reported.

REFERENCES

- 1. United Nations High Commissioner for Refugees. The 2004 Global Report. Geneva, Switzerland: United Nations High Commissioner for Refugees; 2005.
- 2. Norwegian Refugee Council. Internal Displacement: Global Overview of Trends and Developments in 2004. Geneva, Switzerland: Global IDP Project; 2005. 3. United Nations Relief and Works Agency for Pal-
- estinian Refugees in the Near East. Annual Report of the Department of Health 2004. Amman, Jordan: United Nations Relief and Works Agency for Palestinian Refugees in the Near East; 2005.
- 4. Garcia-Peltoniemi R. Epidemiological perspectives. In: Westermeyer J, Williams CL, Nguyen AN, eds. Mental Health Services for Refugees. Washington, DC: US Government Printing Office; 1991:24-41. DHHS Publication ADM 91-1824.
- 5. Kinzie D. Posttraumatic effects and their treatment among Southeast Asian refugees. In: Wilson J, Raphael B, eds. International Handbook of Traumatic Stress Syndromes. New York, NY: Plenum Press; 1993:311-320
- 6. Weisaeth L, Eitinger L. Posttraumatic stress phenomena: common themes across wars, disasters and traumatic events. In: Wilson J, Raphael B, eds. International Handbook of Traumatic Stress Syndromes. New York, NY: Plenum Press; 1993:69-78.
- 7. United Nations High Commissioner for Refugees. Statistical Yearbook 2003: Trends in Displacement, Protection and Solutions, Geneva, Switzerland: United Nations High Commissioner for Refugees; 2004.
- 8. Summerfield D. The psychological legacy of war and atrocity: the question of long-term and transgenerational effects and the need for a broad view. J Nerv Ment Dis. 1996;184:375-377.
- 9. Dohrenwend BP, Dohrenwend BS. Social Status and Psychological Disorder: A Causal Inquiry. New York, NÝ: Wiley; 1969.

- 10. Brown GW. Life events and measurement. In: Brown GW, Harris TO, eds. Life Events and Illness. New York, NY: Guilford Press; 1989:3-48.
- 11. De Girolamo G, McFarlane AC. The epidemiology of PTSD: a comprehensive review of the international literature. In: Marsella AJ, Friedman MJ, Gerrity ET, Scurfield RM, eds. Ethnocultural Aspects of Posttraumatic Stress Disorder: Issues, Research, and Clinical Applications. Washington, DC: American Psychological Association; 1996.
- 12. Kulka RA, Schlenger WE, Fairbank JA, et al. Trauma and the Vietnam War Generation. New York, NY: Brunner & Mazel; 1990.
- 13. Pynoos RS, Frederick C, Nader K, et al. Life threat and posttraumatic stress in school-age children. Arch Gen Psychiatry. 1987;44:1057-1063.
- 14. Rubonis AV, Bickman L. Psychological impairment in the wake of disaster: the disasterpsychopathology relationship. Psychol Bull. 1991;109: 384-399.
- 15. Bracken PJ, Giller JE, Summerfield D. Psychological responses to war and atrocity: the limitations of current concepts. Soc Sci Med. 1995;40:1073-1082.
- 16. Hollifield M, Warner TD, Lian N, et al. Measuring trauma and health status in refugees: a critical review. JAMA. 2002;288:611-621.
- 17. Jablensky A, Marsella AJ, Ekblad S, Jansson B, Levi L, Bornemann TH. Refugee mental health and wellbeing: conclusions and recommendations. In: Marsella AJ, Bornemann TH, Ekblad S, Orley J, eds. Amidst Peril and Pain: The Mental Health and Well-Being of the World's Refugees. Washington, DC: American Psychological Association; 1994:327-339.
- 18. Marsella AJ, Friedman MJ, Gerrity ET, Scurfield RM, eds. Ethnocultural Aspects of Posttraumatic Stress Disorder: Issues, Research and Clinical Applications. Washington, DC: American Psychological Association: 1996.
- 19. Martin SF. A policy perspective on the mental health and psychosocial needs of refugees. In: Marsella AJ, Bornemann TH, Ekblad S, Orley J, eds. Amidst Peril and Pain: The Mental Health and Well-Being of the World's Refugees. Washington, DC: American Psychological Association; 1994:69-80.
- 20. Eisenbruch M. From post-traumatic stress disorder to cultural bereavement: diagnosis of Southeast Asian refugees. Soc Sci Med. 1991;33:673.
- 21. Higginbotham N, Marsella AJ. International consultation and the homogenization of psychiatry in Southeast Asia. Soc Sci Med. 1988;27:553.
- 22. Hjern A, Angel B, Jeppson O. Political violence, family stress and mental health of refugee children in exile. Scand J Soc Med. 1998;26:18-25.
- 23. De Jong JT, Komproe IH, Van Ommeren M, et al. Lifetime events and posttraumatic stress disorder in 4 postconflict settings. *JAMA*. 2001;286:555-
- 24. Porter M, Haslam N. Forced displacement in Yugoslavia: a meta-analysis of psychological consequences and their moderators. *J Trauma Stress*. 2001; 14:817-834
- 25. Rangaraj A. The health status of refugees in South East Asia. In: Miserez D, ed. Refugees: The Trauma of Exile. Dordrecht, the Netherlands: Marinus Nijhoff;
- 26. Von Buchwald U. Refugee dependency: origins and consequences. In: Marsella AJ, Bornemann TH, Ekblad S, Orley J, eds. Amidst Peril and Pain: The Mental Health and Well-Being of the World's Refugees. Washington, DC: American Psychological Association: 1994:229-237.
- 27. Brody E. The mental health and well-being of refugees: issues and directions. In: Marsella AJ, Bornemann TH, Ekblad S, Orley J, eds. Amidst Peril and Pain: The Mental Health and Well-being of the World's Refugees. Washington, DC: American Psychological Association; 1994:57-68.
- 28. Marsella AJ. Ethnocultural diversity and international refugees: challenges for the global community. In: Marsella AJ, Bornemann TH, Ekblad S, Orley eds. Amidst Peril and Pain: The Mental Health

- and Well-Being of the World's Refugees. Washington, DC: American Psychological Association; 1994: 341-364.
- 29. Green BL, Korol M, Grace M, et al. Children and disaster: age, gender and parental effects on PTSD symptoms. J Am Acad Child Adolesc Psychiatry. 1991;
- 30. Green BL, Kramer TL, Grace MC, et al. Traumatic events over the lifespan: survivors of the Buffalo Creek disaster. In: Miller TW, ed. Clinical Disorders and Stressful Life Events. Madison, Wis: International Universities Press; 1997:283-306. Stress and Health Series monograph 7.
- 31. Green BL, Lindy J, Grace M, et al. Buffalo Creek survivors in the second decade: stability of stress symptoms. Am J Orthopsychiatry. 1990;60:43-54.
- 32. Mollica R, Cui X, McInnes K, Massagli M. Sciencebased policy for psychosocial interventions in refugee camps: a Cambodian example. J Nerv Ment Dis. 2002; 190:158-166.
- 33. Wilkinson CB. Aftermath of a disaster: the collapse of the Hyatt Regency Hotel skywalk. Am J Psychiatry. 1983;140:1134-1139.
- 34. Boothby N, Upton P, Sultan A. Children of Mozambique: The Cost of Survival. Washington, DC: US Committee for Refugees; 1991.
- 35. Garbarino J, Kostelny K, Dubrow N. No Place to Be a Child: Growing up in a War Zone. Lexington, Mass: Lexington Books; 1991.
- 36. Huerta F, Horton R. Coping behavior of elderly flood victims. Gerontologist. 1978;18:541-546.
- 37. Kinzie D, Sack W, Angell R, Manson S, Rath B. The psychic effects of massive trauma on Cambodian children. *J Am Acad Child Psychiatry*. 1986;25:370. **38.** Allden K, Poole C, Chantavanich S, Ohmar K, Aung
- NN, Mollica RF. Burmese political dissidents in Thailand: trauma and survival among young adults in exile. Am J Public Health. 1996;86:1561-1569.
- 39. Freedy J, Shaw D, Jarrell M, Masters C. Towards an understanding of the psychological impact of natural disasters: an application of the conservation resources stress model. J Trauma Stress. 1992;5:441-454.
- 40. Van Willigen LHM, Hondius AJK, Van Der Ploeg HM. Health problems of refugees in the Netherlands. Trop Geogr Med. 1995;47:118-124.
- 41. Silove D. Trauma and forced relocation. Curr Opin Psychiatry. 2000;13:231-236.
- 42. Chung RC, Bemak F, Kagawa-Singer M. Gender differences in psychological distress among Southeast Asian refugees. J Nerv Ment Dis. 1998;186:112-119.
- 43. Conrad M, Hammen C. Role of maternal depression in perceptions of child maladjustment. J Consult Clin Psychol. 1989;57:663-667.
- 44. Locke C, Southwick K, McCloskey L, Fernandez-Esquer ME. The psychological and medical sequelae of war in Central American refugee mothers and children. Arch Pediatr Adolesc Med. 1996;150:822-832.
- 45. Surtees PG. Adversity and psychiatric disorder: a decay model. In: Brown GW, Harris TO, eds. Life Events and Illness. New York, NY: Guilford Press; 1989:160-
- 46. Surtees PG, Miller PM, Ingham JG, Kreitman NB, Rennie D, Sashidharan SP. Life events and the onset of affective disorder: a longitudinal general population study. J Affect Disord. 1986;10:37-50.
- 47. Department of Veterans Affairs. The PILOTS Database: An Electronic Index to the Traumatic Stress Literature. Available at: http://www.ncptsd.org /publications/pilots/index.html. Accessed December 2002
- 48. Adams PL, Horovitz JH. Psychopathology and fatherlessness in poor boys. *Child Psychiatry Hum Dev.* 1980:10:135-143.
- 49. Ahmad A, Sundelin-Wahlsten V, Sofi MA, Qahar JA, Von Knorring A. Reliability and validity of a childspecific cross-cultural instrument for assessing posttraumatic stress disorder. Eur Child Adolesc Psychiatry. 2000;9:285-294.
- 50. Al-Eissa YA. The impact of the Gulf armed conflict on the health and behaviour of Kuwaiti children. Soc Sci Med. 1995;41:1033-1037.

- **51.** Bauer M, Priebe S, Kurten I, Graf K, Baumgartner A. Psychological and endocrine abnormalities in refugees from East Germany, I: prolonged stress, psychopathology, and hypothalamic-pituitary-thyroid axis activity. Psychiatry Res. 1994;51:61-73.
- 52. Bayard-Burfield L, Sundquist J, Johansson S. Ethnicity, self reported psychiatric illness, and intake of psychotropic drugs in five ethnic groups in Sweden. J Epidemiol Community Health. 2001;55:657-664.
- **53.** Beiser M. Changing time perspective and mental health among Southeast Asian refugees. *Cult Med* Psychiatry. 1987;11:437-464.
- 54. Carballo M, Zeric D, Smajkic A. Health and Social Status of Displaced People in Bosnia and Herzegovina. Geneva, Switzerland: International Centre for Migration and Health Report; 1996.
- 55. Cervantes RC, Salgado de Snyder VN, Padilla AM. Posttraumatic stress in immigrants from Central America and Mexico. Hosp Community Psychiatry. 1989;40:615-619
- 56. Clark S, Callahan WJ, Lichtszajn J, Velasquez RJ. MMPI performance of Central American refugees and Mexican immigrants. *Psychol Rep.* 1996;79:819-824. **57.** Dube KC. Mental disorder in Agra. Soc Psychiatry. 1968;3:139-143.
- 58. Dunnigan T, McNall M, Mortimer JT. The problem of metaphorical nonequivalence in cross-cultural survey research: comparing the mental health statuses of Hmong refugee and general population adolescents. *J Cross Cult Psychol*. 1993;24:344-365.
- 59. Eitinger L. The incidence of mental disease among refugees in Norway. J Ment Sci. 1959;105:326-338. **60.** El Habir E, Marriage K, Littlefield L, Pratt K. Teachers' perceptions of maladaptive behaviour in Lebanese refugee children. Aust N Z J Psychiatry. 1994;28:
- 61. Fazel MK, Young DM. Life quality of Tibetans and Hindus: a function of religion. J Sci Study Relig. 1988; 27:229-242.
- 62. Folnegovic-Smalc V, Folnegovic Z, Uzun S, Filibic M, Dujmic S, Makaric G. Psychotrauma related to war and exile as a risk factor for the development of dementia of Alheimer's type in refugees. Croat Med J. 1997:38:273-276
- 63. Goenjian AK, Najarian LM, Pynoos RS, et al. Posttraumatic stress reactions after single and double trauma. Acta Psychiatr Scand. 1994;90:214-221.
- 64. Herceg M, Melamed B, Pregrad J. Effects of war on refugee and non-refugee children from Croatia and Bosnia-Hercegovina. Croat Med J. 1996;37:111-114. 65. Hourani LL, Armenian HK, Zurayk H, Afifi L. A population-based survey of loss and psychological distress during war. Soc Sci Med. 1986;23:269-275.
- 66. Howard M, Hodes M. Psychopathology, adversity, and service utilization of young refugees. *J Am Acad Child Adolesc Psychiatry*. 2000;39:368-377.
- 67. Jensen SB, Schaumburg E, Leroy B, Larsen BO. Psychiatric care of refugees exposed to organized violence: a comparative study of refugees and immigrants in Frederiksborg County, Denmark. Acta Psychiatr Scand. 1989;80:125-131.
- 68. Klimidis S, Stuart G, Minas IH, Ata AW. Immigrant status and gender effects on psychopathology and self-concept in adolescents: a test of the migration morbidity hypothesis. Compr Psychiatry. 1994;35:393-404. **69.** Kocijan-Hercigonja D, Rijavec J, Parry-Jones W, Remeta D. Psychologic problems of children wounded during the war in Croatia. Nord Psykiatr Tidsskr. 1996; 50:451-456.
- 70. Kocijan-Hercigonja D, Rijavec M, Marusic A, Hercigonja V. Coping strategies of refugee, displaced, and non-displaced children in a war area. Nord J Psychiatry. 1998:52:45-50.
- 71. Kocijan-Hercigonja D, Sabioncello A, Rijavec M, et al. Psychological condition hormone levels in war trauma. *J Psychiatr Res.* 1996;30:391-399. **72.** Kondic L, Mavar M. Anxiety and depressive reac-
- tions in refugees. Psychol Beitr. 1992;34:179-183.
- 73. Kuterovac G, Dyregrov A, Stuvland R. Children in war: a silent majority under stress. Br J Med Psychol. 1994;67:363-375

- 74. Laor N, Wolmer L, Mayes LC, et al. Israeli preschoolers under Scud missile attacks: a developmental perspective on risk-modifying factors. Arch Gen Psychiatry. 1996;53:416-423.
- 75. Lopes Cardozo B, Vergara A, Agani F, Gotway CA. Mental health, social functioning, and attitudes of Kosovar Albanians following the war in Kosovo. JAMA. 2000:284:569-577
- 76. Loughry M, Flouri E. The behavioral and emotional problems of former unaccompanied refugee children 3-4 years after their return to Vietnam. Child Abuse Negl. 2001;25:249-263.
- 77. Macksoud MS, Aber JL. The war experiences and psychosocial development of children in Lebanon. Child Dév. 1996;67:70-88
- 78. McCloskey LA, Southwick K, Fernandez-Esquer ME, Locke C. The psychological effects of political and domestic violence on Central American and Mexican immigrant mothers and children. J Community Psychol. 1995.23.95-116
- 79. Myers KM, Croake JW, Singh A. Adult fears of four ethnic groups: whites, Chinese, Japanese and "boat people." *Int J Soc Psychiatry*. 1987;33:56-67.
- 80. Onwumere J, Holttum S, Hirst F. Determinants of quality of life in black African women with HIV living in London. Psychol Health Med. 2002;7:61-74.
- 81. Paardekooper B, de Jong JTVM, Hermanns JMA. The psychological impact of war and the refugee situation on South Sudanese children in refugee camps in northern Uganda: an exploratory study. J Child Psychol Psychiatry. 1999;40:529-536.
- 82. Pernice R, Brook J. Relationship of migrant status (refugee or immigrant) to mental health. Int J Soc Psychiatry. 1994;40:177-188.
- 83. Punamaki RL. Relationships between political violence and psychological responses among Palestinian women. J Peace Res. 1990;27:75-85.
- 84. Quta S, El Sarraj E. Level of anxiety in Gaza before and after the Intifada. Derasat Nafseyah. 1993;3:
- 85. Rasanen E. Change in culture and language environment in childhood and its effect on adult life. Acta Psychiatr Scand. 1989;80:280-286. 86. Rasekh Z, Bauer HM, Manos MM, Iacopino V.
- Women's health and human rights in Afghanistan. JAMA. 1998;280:449-455.
- 87. Reeler AP. A preliminary investigation into psychological disorders among Mozambican refugees: prevalence and clinical features. Cent Afr J Med. 1994; 40:309-315.
- 88. Riolli L, Savicki V, Cepani A. Resilience in the face of catastrophe: optimism, personality and coping in the Kosovo crisis. J Appl Soc Psychol. 2002;32: 1604-1627.
- 89. Roglic G, Pibernik-Okanovic M, Prasek M, Metelko Z. Effect of war-induced prolonged stress on cortisol of persons with type II diabetes mellitus, Behav Med. 1993.19.53-59
- 90. Roncevic-Grzeta I, Franciskovic T, Moro L, Kastelan A. Depression and torture. Mil Med. 2001;166:530-
- 91. Rousseau C, Drapeau A, Platt R. Living conditions and emotional profiles of Cambodian, Central American and Quebecois youth. Can J Psychiatry. 2000;45: 905-911.
- 92. Sabioncello A, Kocijan-Hercigonja D, Rabatic S, et al. Immune, endocrine, and psychological responses in civilians displaced by war. Psychosom Med. 2000;62:502-
- 93. Silove D, Steel Z, McGorry PD, Mohan P. Trauma exposure, postmigration stressors, and symptoms of anxiety, depression and post-traumatic stress in Tamil asylum-seekers: comparison with refugees and immigrants. Acta Psychiatr Scand. 1998;97:175-181. 94. Smither R, Rodriguez-Giegling M. Marginality, modernity, and anxiety in Indochinese refugees. J Cross Cult Psychol. 1979;10:469-478.
- 95. Sundelin-Wahlsten V. Ahmad A. Von Knorring A-L. Traumatic experiences and post-traumatic stress reactions in children from Kurdistan and Sweden. Acta Paediatr. 2001;90:563-568.

- 96. Sundquist J. Ethnicity as a risk factor for mental illness: a population-based study of 338 Latin American refugees and 996 age-, sex- and education-matched Swedish controls. Acta Psychiatr Scand. 1993;87:208-
- 97. Sundquist J. Refugees, labour migrants and psychological distress: a population-based study of 338 Latin-American refugees, 161 South European and 396 Finnish labour migrants, and 996 Swedish age-, sexand education-matched controls. Soc Psychiatry Psychiatr Epidemiol. 1994;29:20-24.
- 98. Sundquist J, Johansson SE. The influence of exile and repatriation on mental and physical health: a population-based study. Soc Psychiatry Psychiatr Epidemiol. 1996:31:21-28.
- 99. Thulesius H, Hakansson A. Screening for posttraumatic stress disorder symptoms among Bosnian refugees. J Trauma Stress. 1999;12:167-174.
- 100. Tousignant M, Habimana E, Biron C, Malo C. Sidoli-LeBlanc E, Bendris N. The Quebec Adolescent Refugee Project: psychopathology and family variables in a sample from 35 nations. J Am Acad Child Adolesc Psychiatry. 1999;38:1426-1432.
- 101. Wong-Rieger D, Quintana D. Comparative acculturation of Southeast Asian and Hispanic immigrants and sojourners. J Cross Cult Psychol. 1987;18: 345-362.
- 102. Young MY, Evans DR. The well-being of Salvadoran refugees. Int J Psychol. 1997;32:289-300.
- 103. Zivcic I. Emotional reactions of children to war stress in Croatia. J Am Acad Child Adolesc Psychiatry. 1993:32:709-713.
- 104. Beck AT, Steer RA. The Beck Depression Inventory. New York, NY: Psychological Corp; 1987.
- 105. Mollica RF, Caspi-Yavin Y, Bollini P, Truong T, Tor S, Lavelle J. The Harvard Trauma Questionnaire: validating a cross-cultural instrument for measuring torture, trauma, and posttraumatic stress disorder in Indochinese refugees. J Nerv Ment Dis. 1992;180:111-116.
- 106. Achenbach T. Manual for the Child Behavior Checklist. Burlington: University of Vermont; 1991.
- 107. Cooper H. Synthesizing Research: A Guide for Literature Reviews. 3rd ed. New York, NY: Russell Sage Foundation; 1998.
- 108. Lipsey MW, Wilson DB. Practical Meta-analysis. New York, NY: Russell Sage Foundation; 2001. Applied Social Research Methods Series, Volume 49.
- 109. Glass GV. Primary, secondary, and metaanalysis of research. Educ Res. 1976;5:3-8.
- 110. Hedges LV. Unbiased Estimation of Effect Size. Oxford, England: Pergamon Press; 1980:25-27. Evaluation in Education: An International Review Series.
- 111. Huffcutt Al. Arthur W. Development of a new outlier statistic for meta-analytic data. J Appl Psychol. 1995; 80.327-334
- 112. Hedges LV, Olkin I. Statistical Methods for Meta-analysis. Orlando, Fla: Academic Press; 1985.
- 113. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002;21:
- 114. Rosenthal R. The "file drawer problem" and tolerance for null results. Psychol Bull. 1979;86:638-641. 115. Elvik R. Evaluating the statistical conclusion validity of weighted mean results in meta-analysis by analyzing funnel graph diagrams. Accid Anal Prev. 1998; 30:255-266.
- 116. Light RJ, Singer JD, Willett JB. The visual presentation and interpretation of meta-analysis. In: Cooper H, Hedges LV, eds. The Handbook of Research Synthesis. New York, NY: Russell Sage Foundation; 1994: 439-453
- 117. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. Riometrics 1994:50:1088-1101
- 118. Song F, Khan KS, Dinnes J, Sutton AJ. Asymmetric funnel plots and publication bias in meta-analyses of diagnostic accuracy. Int J Epidemiol. 2002;31:88-95. 119. Cohen J. Statistical Power Analysis for the Behavior Sciences. 2nd ed. Hillsdale, NJ: Erlbaum; 1988.