

Cost-effectiveness of Intensive Psychiatric Community Care for High Users of Inpatient Services

Robert A. Rosenheck, MD; Michael S. Neale, PhD

Background: This 2-year experimental study evaluated the effectiveness and cost of 10 intensive psychiatric community care (IPCC) programs at Department of Veterans Affairs medical centers in the northeastern United States.

Methods: High users of inpatient services were randomly assigned to either IPCC or standard Department of Veterans Affairs care at 6 general medical and surgical hospitals (n=271 vs 257) and 4 neuropsychiatric hospitals (n=183 vs 162). Patient interviews every 6 months and national computerized data were used to assess clinical outcomes, health service use, health care costs, and non-health care costs.

Results: There was only 1 significant clinical difference between groups across follow-up periods: IPCC patients at general medical and surgical sites had higher community living skills. However, at the final interview, IPCC patients at general medical and surgical sites showed significantly lower symptoms, higher functioning, and greater

satisfaction with services. Treatment with IPCC significantly reduced hospital use only at neuropsychiatric sites (320 vs 513 days, $P<.001$). Total societal costs, including the cost of IPCC, were lower for IPCC at neuropsychiatric sites (\$82 454 vs \$116 651, $P<.001$), but greater at general medical and surgical sites (\$51 537 vs \$46 491, $P<.01$). When 2 sites that incompletely implemented the model were dropped from the analysis, costs at general medical and surgical sites were \$38 lower for IPCC ($P=.26$).

Conclusions: At acute care hospitals, IPCC treatment is associated with greater long-term clinical improvement and, when fully implemented, is cost-neutral. At long-stay hospitals treating older, less-functional patients, it is not associated with clinical or functional improvement but generates substantial cost savings. Intensive psychiatric community care thus has beneficial, but somewhat different, outcome profiles at different types of hospitals.

Arch Gen Psychiatry. 1998;55:459-466

EXPERIMENTAL studies published during the past 15 years have reported that concentrating treatment resources in community-based assertive community treatment (ACT) teams or intensive case-management programs can result in improved clinical status of severely mentally ill patients at no additional cost.¹⁻⁶ Some studies, however, have found case management to be associated with no clinical change and/or increased service utilization and cost.⁷⁻¹¹ Three recent reviews¹²⁻¹⁴ concluded that intensive community treatment frequently reduces hospital use but does not always achieve net cost savings, and far less consistently achieves clinical improvement.

Most published controlled trials of ACT were conducted at single sites as specially funded research or demonstration programs implemented by leaders in the

field. Although these demonstrations illustrate the potential value of intensive community treatment under optimal operating circumstances, the applicability of their findings to ordinary conditions is uncertain.¹⁵ There is, in addition, a need to identify clinical settings in which such programs are consistently cost-effective.

In this article we evaluate the effects of 10 intensive psychiatric community care (IPCC) programs on the clinical status, community adjustment, and total costs to society (health care costs and non-health care costs) of high users of Department of Veterans Affairs (VA) psychiatric inpatient care. The study is distinctive because it involves multiple sites comprising 2 distinct types of facility, neuropsychiatric hospitals and general medical and surgical hospitals, and was implemented under routine circumstances, as would be the case with dissemination across large public health care systems.

From the Northeast Program Evaluation Center, Veterans Affairs Connecticut Healthcare System (Drs Rosenheck and Neale), and the Department of Psychiatry, Yale Medical School (Dr Rosenheck), West Haven, Conn.

SUBJECTS AND METHODS

SUBJECT ELIGIBILITY AND PROGRAM ENTRY

Veterans were eligible for the study if they: (1) were currently hospitalized on a VA psychiatric inpatient unit, (2) had a primary psychiatric diagnosis (as determined by the primary clinician) other than substance abuse or organic brain disorder, and (3) met criteria for high recent psychiatric inpatient utilization (see below). Patients who met these criteria were referred to the research team by their inpatient clinicians. Recruitment occurred from 1987 to 1990 and the last follow-up interview was completed in mid 1992. Virtually all patients referred for screening met the eligibility criteria and 85% agreed to participate in the study. After giving written informed consent to participate, patients were randomized to IPCC or standard VA treatment by the coin-flip of an independent research coordinator. Those assigned to IPCC were introduced to their clinicians shortly after randomization. Control patients continued their prior inpatient treatment until they were determined on clinical grounds to be ready for discharge.

SITE TYPES

Two contrasting types of VA facilities were involved. Neuropsychiatric sites, typically located in suburban or rural settings, are large facilities that provide long-term mental health care to severely disabled patients. General medical and surgical sites are located in urban centers and provide shorter-term, crisis-oriented inpatient care.

Utilization criteria for program entry were site type-specific because of very different utilization patterns.¹⁶ Entry criteria required 180 days of hospitalization during the previous year or 4 or more prior admissions at long-stay neuropsychiatric facilities; and 40 or more days of psychiatric hospitalization or 2 prior admissions at general medical and surgical sites.

IPCC PROGRAM

The IPCC program was designed through a comprehensive literature review supplemented by consultation from national experts in the Wisconsin ACT program. The IPCC program was designed to operate under 4 core principles: (1) Intensity. Patients were to be seen as frequently as clinically indicated and case loads were to be low (7-15

patients per clinician). (2) Flexibility and community orientation. Clinicians were urged to provide the majority of contacts in community settings where maximal clinical leverage could be obtained. (3) Rehabilitation focus. A broad range of rehabilitation services was emphasized, such as training in practical problem solving and adaptive skill building. (4) Continuity of care. The IPCC teams were to be a "fixed point of continuing responsibility,"¹⁷ assertively maintaining contact with even the most reluctant patients, although parallel involvement in other VA services was not discouraged.

Most team members were masters-level social workers and nurses, and each team was staffed with a part-time psychiatrist. Teams met to review cases at least weekly (in some instances daily), although team focus was not emphasized in the IPCC program as much as it has been emphasized in recent writing about the Wisconsin ACT model.¹⁸ Staff were available to other VA clinicians and to clients after hours and on weekends.

STANDARD CARE

Standard VA care provided to controls included inpatient psychiatric and psychopharmacologic treatment, outpatient psychiatric treatment, and rehabilitation services, such as work therapy. These services did not differ programmatically between general medical and surgical and neuropsychiatric sites. There was substantial fiscal and administrative pressure for both IPCC and standard-care patients to be discharged from the hospital as soon as possible.

ASSESSMENT OF PROGRAM IMPLEMENTATION

Program implementation was monitored using weekly clinician logs, structured clinical summaries completed on each case every 6 months, and a standardized rating instrument of team operation based on a checklist of 23 ACT program characteristics, previously shown to predict program effectiveness.¹⁸⁻²⁰ These data allow evaluation of program implementation at each site with respect to: (1) the frequency, intensity, and location of clinical contacts; (2) the types of services provided; (3) duration of involvement; and (4) overall fidelity to the ACT model.¹⁶

CLINICAL OUTCOME MEASURES

Data on clinical status and community adjustment were obtained through face-to-face patient interviews conducted

RESULTS

STUDY SAMPLE

Table 1 shows that patients at neuropsychiatric sites were older, less likely to have been married or working, and less likely to have substance abuse problems (which were relatively infrequent at both types of site). The cohort had substantial hospitalization histories, averaging 452 days during the 2 years prior to program entry at neuropsychiatric sites and 123 days at general medical and surgical sites.

Comparison of sociodemographic and baseline clinical characteristics of experimental and control patients showed no significant differences on any of 21 measures

at general medical and surgical sites, but significantly different scores on 2 measures at neuropsychiatric sites: the BPRS (mean=16.0 for IPCC patients vs 19.6 for controls; $t=2.97$, $df=273$, $P<.01$) and the GAS (mean=42.1 for IPCC patients vs 37.9 for controls; $t=3.05$, $df=340$, $P<.01$).

IMPLEMENTATION

Across the entire IPCC program, caseloads averaged 13.2 patients per clinician. The IPCC patients had face-to-face clinical contacts with their IPCC clinicians during 60 of the 104 weeks (58%) of the study, and averaged 1.9 contacts per patient per week and 41 minutes per contact during those weeks. More than half (64%) of face-to-face clinical

by independent interviewers every 6 months. Interviewers received 3 days of intensive training in the use of all instruments at the beginning of the project and periodically thereafter.

Psychiatric symptoms were assessed with both an observer-rated instrument, the Brief Psychiatric Rating Scale (BPRS),²¹ and a self-report symptom measure, the Brief Symptom Inventory.²² Substance abuse was evaluated with the alcohol and drug composite indices of the Addiction Severity Index.²³ Functional status was assessed with the Global Assessment Scale (GAS) (although scores on this measure are somewhat contaminated by symptoms)²⁴ and with a self-report measure of community functioning that documented capability to shop, socialize, and perform other basic activities independently. Satisfaction with services was evaluated with a series of 5-point Likert-type questions (ranging from 0=very dissatisfied to 4=very satisfied) that addressed satisfaction with IPCC treatment and other VA and non-VA mental health services.

Community tenure was assessed by subtracting all known days of hospitalization or other days residing in controlled environments (such as nursing homes and jails) from 730 days (2 years).

ASSESSMENT OF COSTS

Health care costs were estimated by multiplying the number of units of service by their estimated unit costs. Veterans Affairs service use was determined from national computerized workload databases and both medication and non-VA service use from interviews. Use of non-VA services may have been undercounted owing to recall difficulties during a 6-month period. Unit costs for VA general psychiatry, substance abuse, medical and surgical, inpatient, and outpatient care (including group treatment and day hospital care) were estimated for each medical center from site-specific cost data available from the local cost distribution report and computerized workload data, using methods described elsewhere.^{16,25,26} Reported non-VA costs were minimal (<3% of all costs) and unit costs were estimated on the basis of published data from private hospitals.

Interview data were also used to estimate non-health care service use and costs including: (1) the administrative costs of transferring payments (eg, disability and welfare)^{27,28}; (2) criminal justice system costs (police contacts, arrests, court appearances, nights in jail, and probation visits)^{29,30}; and (3) productivity (estimated by employment earnings, included as negative costs).

cal time took place in community settings. Seventy-eight percent of patients were receiving at least 1 type of rehabilitative service after 1 year (the midpoint of the study) and 75% after 2 years. Only 12% of IPCC patients dropped out before the study period was over.

The fidelity of IPCC treatment to the ACT model was compared with 14 model ACT teams implemented under closely supervised conditions and a set of standard mental health clinics. Intensive psychiatric community care programs (mean score=3.7, SD=0.55 [potential range, 0-5]) were substantially more ACT-like than the control programs (mean score=2.4, SD=0.36), but slightly less ACT-like than the model programs (mean score=4.0, SD= 0.39).²⁰

Cost data were summarized and analyzed from the perspectives of the health care system (VA and non-VA, inpatient and outpatient, psychiatric and medical and surgical, plus drug costs) and society (health care costs plus all non-health care costs, minus productivity).

ANALYSES

Analyses were based on intention-to-treat principles including all subjects as randomized. Although 83% of planned interviews were completed at general medical and surgical facilities (88% among IPCC patients vs 77% among controls) and 77% at neuropsychiatric facilities (77% for IPCC vs 76% for controls), primary longitudinal outcomes were analyzed using random effects regression models³¹ to maximize statistical power for testing hypotheses with longitudinal data. These analyses were conducted using the PROC MIXED function from the SAS statistical computer software package, version 6.09 (SAS Institute Inc, Cary, NC). These models accommodate correlations among the repeated observations and missing data at some assessment periods. The significance of differences between treatment conditions was tested using the likelihood ratio χ^2 statistic. Specifically, we compared a model that included the effects of time, time squared, and treatment condition to a model that adds group \times time interactions. The hypothesis of interest is the group \times time interaction. A secondary analysis of outcomes was conducted using analysis of covariance (ANCOVA) comparing groups at the final follow-up interview, controlling for baseline status on the follow-up measure. Follow-up rates for this interview at general medical and surgical facilities were 82% for IPCC patients and 73% for control patients and 71% for IPCC patients and 67% for control patients at neuropsychiatric facilities. At general medical and surgical sites, patients who were followed up had higher GAS scores ($P<.009$) and were more likely to be married ($P<.03$) than patients who were not. At neuropsychiatric sites, those who were followed up had lower BPRS scores ($P<.02$) and were also more likely to be married ($P<.01$).

Cost data were analyzed in their raw form and with natural log transformation using ANCOVA, controlling for utilization and cost during the 6 months before study entry.

Findings were also summarized as incremental cost-effectiveness ratios, the ratio of the incremental cost of the treatment (the average cost of IPCC minus the average cost of standard VA care) to the incremental effectiveness of the treatment (the average effectiveness of IPCC minus the average effectiveness of standard VA care).^{32,33}

Site-by-site implementation data presented elsewhere¹⁶ showed that 2 general medical and surgical sites saw patients only once every 3 weeks during the 2-year period, as compared with about twice every 3 weeks at the other sites. These 2 sites also had the lowest proportion of time spent seeing patients in community settings (43% and 0%, as compared with 72% at other sites), and the lowest scores on the ACT fidelity measure (3.1 and 2.4), both more than 1 SD below the mean of other IPCC sites.

OUTCOMES

Table 2 presents data on clinical and social adjustment outcomes at baseline and during the 2 years following study

Table 1. Baseline Data Comparing Veterans From Neuropsychiatric and General Medical and Surgical Sites

Characteristic*	General Medical and Surgical (n = 528)	Neuropsychiatric (n = 345)	χ^2 Test	Student t Test	P
Mean age, y (SD)	44 (13)	54 (12)		10.2	<.001
>45, %	35	68	90.1		<.001
Nonwhite, %	25	13	17.7		<.001
Never married, %	44	70	55.8		<.001
Working full- or part-time, %	25	4	64.2		<.001
VA compensation, %	57	53	1.3		
Preenrollment hospital days, No. (SD)†	123 (101)	452 (225)		25.0	<.001
Median	88	412			
Primary clinical diagnosis, %					
Schizophrenia	39	68	65.5		<.001
Other psychoses	13	10	1.5		
Bipolar disorder	13	6	10.2		<.01
Depression	10	4	9.7		<.01
Alcohol/drug abuse	7	8	0.3		
Personality disorder	8	0	27.1		<.001
Other	10	4	9.7		<.01
Dually diagnosed, %	30	23	4.9		<.05
Total pathology, BPRS (SD)	15 (8)	18 (9)		4.42	<.001
General severity, BSI (SD)	1.32 (0.82)	0.80 (0.65)		9.01	<.001
Alcohol index, ASI (SD)	0.12 (0.22)	0.10 (0.17)		1.52	
Global Assessment Scale (SD)	45 (11)	40 (12)		6.28	<.001
Community Living Skills (SD)	15 (3)	12 (4)		8.28	<.001

*VA indicates Veterans Affairs; BPRS, Brief Psychiatric Rating Scale; BSI, Brief Symptom Inventory; and, ASI, Addiction Severity Scale.

†Preenrollment hospital days is for a 2-year period.

entry. Comparison of outcomes at general medical and surgical sites across all time points using random regression analysis showed significant differences on only 1 measure (community living skills), with better functioning for IPCC patients (χ^2 test=8.65, $df=2$, $P<.01$). There were no significant differences between IPCC clients and controls on any measures at neuropsychiatric sites.

Figures in the “Comparative Change” columns of Table 2 show the difference between IPCC and standard VA treatment in the amount of change from baseline. Differences are minimal at most time points, but substantial differences favoring IPCC appear at general medical and surgical sites at the 2-year assessment on several measures. Significant differences favor IPCC treatment on total symptoms (BPRS) ($F=8.27$; $df=1,402$; $P<.004$); community living skills ($F=11.62$; $df=1,366$; $P<.001$); global functioning (GAS) ($F=11.12$; $df=1,409$; $P<.001$); and satisfaction with services ($F=5.93$; $df=1,324$; $P<.02$).

Comparison of IPCC and standard VA care on total days living in the community showed no significant difference at general medical and surgical sites (591 days for IPCC vs 579 for standard VA care; $F=1.12$; $df=1,525$; $P=.29$), but substantially greater community days at neuropsychiatric sites (375 days for IPCC vs 224 for standard VA care; $F=23.7$; $df=1,342$; $P<.001$). All but 2 of these differences remain significant even after reducing the standard significance level to .004 to adjust for 14 comparisons.

SERVICE USE AND COST

Table 3 shows that there were no statistically significant differences at general medical and surgical sites in

inpatient or outpatient service utilization during the 2-year follow-up period. Analysis of cost data also showed no significant differences between groups on standard inpatient or outpatient health care costs. However, adding the cost of IPCC treatment (\$5793 at general medical and surgical sites) increased total health care costs for the IPCC group by about \$5000 over the standard care group ($P<.002$). Non-health care costs were relatively small in magnitude and not significantly different across groups with 1 exception—the administrative cost of transferring payments was larger in the IPCC group, probably reflecting the case managers’ facilitation of access to benefits. In the final intention-to-treat analysis of societal costs at general medical and surgical sites, IPCC was about \$5000 more expensive than standard VA care ($P<.01$).

At neuropsychiatric sites there were significantly fewer hospital days and significantly greater outpatient visits for IPCC (Table 3). The most dramatic difference is noted for non-mental health hospital days reflecting care on long-term “intermediate care” units, a level of VA hospital care between acute hospitalization and nursing home care. Differences in hospital utilization at neuropsychiatric sites were attributable both to a greater proportion of patients having been discharged (82% for IPCC vs 57% for standard VA care; χ^2 test=24.9, $df=1$, $P<.001$) and to reduced hospital utilization among those who were discharged (average days, 216 for IPCC vs 275 for standard VA care; $t=2.2$, $df=241$, $P<.02$).

These differences in inpatient use resulted in more than \$35 000 lower inpatient costs for IPCC patients, and modestly increased outpatient costs. Total societal costs averaged \$34 000 less for IPCC patients at neuropsychiatric sites ($P<.001$).

Table 2. Clinical Outcomes: IPCC vs Standard VA care at General Medical and Surgical and Neuropsychiatric Facilities*

	General Medical and Surgical Sites†					Neuropsychiatric Sites‡				
	IPCC	SD	Standard Care	SD	Comparative Change§	IPCC	SD	Standard Care	SD	Comparative Change§
Total pathology (BPRS)										
Baseline	14.17	7.41	14.94	8.32	0.00	15.98	9.24	19.57	10.80	0.00
6 mo	14.49	8.78	13.45	6.95	1.80	19.00	9.45	23.36	11.34	-0.77
12 mo	13.96	9.31	14.42	9.27	0.31	18.95	9.83	20.90	10.18	1.65
18 mo	12.98	8.71	13.85	8.86	-0.10	17.42	8.84	19.81	10.05	1.20
24 mo	11.36	8.30	14.32	9.55	-2.19	15.60	9.20	17.56	11.08	1.63
General severity (BSI)										
Baseline	1.27	0.84	1.38	0.82	0.00	0.76	0.66	0.86	0.80	0.00
6 mo	1.22	0.88	1.29	0.87	0.03	0.84	0.67	0.89	0.83	0.05
12 mo	1.19	0.89	1.20	0.85	0.10	0.72	0.69	0.76	0.77	0.06
18 mo	1.16	0.90	1.19	0.83	0.07	0.67	0.64	0.76	0.81	0.02
24 mo	1.07	0.83	1.30	0.93	-0.12	0.60	0.60	0.73	0.77	-0.02
Alcohol index (ASI)										
Baseline	0.11	0.19	0.13	0.24	0.00	0.08	0.18	0.11	0.19	0.00
6 mo	0.07	0.16	0.07	0.14	0.02	0.03	0.10	0.04	0.13	0.02
12 mo	0.07	0.15	0.07	0.14	0.03	0.03	0.08	0.04	0.13	0.02
18 mo	0.07	0.13	0.08	0.16	0.00	0.03	0.09	0.03	0.11	0.03
24 mo	0.07	0.14	0.09	0.17	0.01	0.02	0.08	0.03	0.10	0.02
Global Assessment Scale										
Baseline	45.83	11.10	45.15	11.32	0.00	42.10	13.37	37.86	12.24	0.00
6 mo	49.54	12.89	47.91	11.89	0.95	43.47	12.74	39.94	12.60	-0.72
12 mo	49.15	13.36	49.59	12.24	-1.12	43.49	13.72	40.85	11.61	-1.61
18 mo	51.60	11.85	50.25	11.76	0.68	43.34	12.81	40.82	11.13	-1.72
24 mo	52.76	11.27	48.78	11.69	3.31	45.33	14.72	41.89	12.35	-0.81
Community Living Skills										
Baseline	14.66	3.38	14.58	3.52	0.00	12.51	4.64	11.85	4.20	0.00
6 mo	14.11	3.33	14.40	3.54	-0.38	13.22	4.47	13.68	3.87	-1.13
12 mo	14.00	3.78	14.23	3.76	-0.31	12.75	4.58	13.74	4.48	-1.66
18 mo	14.16	3.89	13.81	3.87	0.27	12.96	4.31	13.03	4.87	-0.73
24 mo	14.87	3.76	13.47	4.13	1.32	13.18	4.27	12.62	4.34	-0.09
Satisfaction										
Baseline	2.90	1.27	2.89	1.26	0.00	2.78	1.26	2.87	1.23	0.00
6 mo	3.03	1.23	3.09	1.21	-0.07	2.95	1.10	2.83	1.03	0.21
12 mo	3.06	1.26	3.01	1.25	0.05	2.98	1.19	2.90	1.13	0.17
18 mo	3.11	1.28	2.77	1.40	0.33	3.11	1.09	3.10	1.06	0.10
24 mo	3.17	1.19	2.82	1.37	0.33	3.00	1.26	3.13	0.96	-0.04

*IPCC indicates intensive psychiatric community care, VA, Veterans Affairs; BPRS, Brief Psychiatric Rating Scale; BSI, Brief Symptom Inventory; and ASI, Addiction Severity Index.

†For IPCC and standard care, n = 271 and 257 at baseline, respectively; 233 and 177 at 6 months; 239 and 181 at 12 months; 231 and 181 at 18 months; and 221 and 189 at 24 months.

‡For IPCC and standard care, n = 183 and 162 at baseline, respectively; 137 and 121 at 6 months; 132 and 110 at 12 months; 127 and 114 at 18 months; and 129 and 108 at 24 months.

§Comparative change is the difference (IPCC minus standard care) in change from baseline. Thus, negative values reflect greater decrease or less increase among IPCC patients on measures of pathology such as the BPRS, the BSI, and the ASI alcohol index. Positive values reflect greater increase or less decrease among IPCC patients on measures of functioning such as the Global Assessment Scale, the Community Living Skills Scale, and the satisfaction with services measure.

||At general medical and surgical sites, $\chi^2 = 8.65$, $df = 2$, $P < .01$.

EXCLUSION OF SITES WITH INCOMPLETE IMPLEMENTATION

Analyses were repeated excluding the 2 general medical and surgical sites that did not adequately implement the IPCC program. Although clinical outcome results did not change, hospital days at the remaining sites were lower for IPCC patients (99.8 hospital days for IPCC vs 128.9 for standard VA care; $F=6.01$, $df=1,344$, $P<.02$), as were inpatient costs (\$27 776 for IPCC vs \$35 570 for standard VA care; $F=5.21$, $df=1,344$, $P<.05$). Community days were significantly higher for IPCC patients (605.0 for IPCC vs 577.2 for standard VA care; $F=3.7$, $df=1,344$, $P<.05$). Societal costs dropped to \$44 772 for IPCC and

\$44 810 for standard VA care, a nonsignificant difference of \$38.

COST-EFFECTIVENESS RATIOS

At general medical and surgical sites cost-effectiveness ratios reveal costs of \$1705 per unit of improvement on the BPRS (95% confidence interval [CI], -\$1200 to \$4609); and \$1268 per unit of improvement on the GAS (95% CI, -\$880 to \$3416). However, with the 2 poor implementation sites excluded, cost-effectiveness ratios at general medical and surgical sites become -\$13 per unit of improvement on the BPRS (95% CI, -\$2494 to \$2520) and -\$10 per unit of improvement on the GAS (95% CI, -\$880

Table 3. Two-Year Utilization and Cost Data: IPCC vs Standard Care*

Utilization/Cost Category	General Medical and Surgical Sites						Neuropsychiatric Sites					
	IPCC		Standard Care		F†	P	IPCC		Standard Care		F‡	P
	(n = 271)	SD	(n = 257)	SD			(n = 183)	SD	(n = 162)	SD		
Utilization, No. of days												
Inpatient/residential care days	131.0	151.2	141.4	168.9	0.46	.50	319.8	290.9	512.5	366.0	26.82	<.001
Neuropsychiatric	101.5	101.0	105.6	116.0	1.32	.25	216.0	252.4	284.3	298.3	2.78	.10
Medical/surgical	29.5	112.6	35.8	115.3	0.31	.58	103.8	242.6	228.3	394.9	18.34	<.001
Outpatient services	128.8	150.4	130.5	148.1	0.09	.77	130.3	187.0	67.9	141.7	12.17	<.001
Neuropsychiatric	80.7	108.7	84.8	113.9	0.33	.57	47.6	95.2	33.6	80.7	1.13	.29
Medical/surgical	48.1	69.3	45.7	57.2	0.15	.70	82.7	116.6	34.2	81.4	18.07	<.001
Costs												
All health care costs, \$\$	51 047	41 227	46 093	41 676	9.78	.002	81 952	70 867	116 267	87 099	11.37	<.001
Inpatient/residential care	36 024	37 561	37 875	40 060	0.17	.68	69 527	71 429	113 654	89 029	15.25	<.001
Neuropsychiatric	29 404	30 291	30 164	32 945	0.22	.64	40 637	47 625	56 501	60 410	3.93	.05
Medical/surgical	6619	24 134	7711	20 957	0.21	.65	28 937	68 982	57 163	99 540	14.14	<.001
Outpatient care	15 023	13 509	8218	12 085	0.57	.45	12 425	9521	2613	5469	116.37	<.001
IPCC cost	7307	5848	0				7831	6305	0			
Mental health	5935	10 790	6447	11 219	0.47	.49	2597	5997	1821	4685	1.12	.29
Other outpatient	1781	2120	1770	1879	0.00	1.00	1997	1980	792	1288	45.46	<.001
Non-health care costs, \$	490	1332	398	1499	0.83	.36	503	429	384	387	6.21	.01
Productivity	253	820	214	766	0.44	.51	20	92	24	63	3.08	.08
Transfer payments	582	427	438	384	25.73	.00	494	379	394	376	1.79	.18
Criminal justice system	161	871	173	1189	0.00	.98	28	172	14	83	0.77	.38
Total cost to society, \$\$	51 537	41 372	46 491	41 638	6.20	.01	82 454	70 834	116 651	87 065	11.35	<.001

*IPCC indicates intensive psychiatric community care.

†df = 1,525 for all analyses.

‡df = 1,342 for all analyses.

§Statistical analysis on natural log transformation of cost data.

||Only administrative costs of transfer payments are included (see reference 23).

to \$3416). Negative ratios are not unambiguously interpretable, because large negative ratios could reflect either greater cost savings (a desirable result) or less effectiveness (an undesirable result). A negative cost-effectiveness ratio thus means that the treatment has greater cost-effectiveness but does not reflect its relative magnitude. The large CIs about the cost-effectiveness ratios are due to the characteristically large SDs of the cost data. Cost-effectiveness ratios were not calculated for neuropsychiatric sites, at which IPCC yielded no significant gain in effectiveness.

COMMENT

Although IPCC is one of the most thoroughly studied psychosocial interventions for people with severe mental illness,^{1-14,16-19} the present study adds to our understanding of this approach in 5 ways.

COST-EFFECTIVENESS

First and most broadly, this study confirms findings of other experimental studies by demonstrating that IPCC is a cost-effective treatment associated with long-term improvement in psychiatric status, social functioning, and consumer satisfaction at acute care sites and no deterioration at neuropsychiatric sites. After excluding sites that did not fully implement the program, both health care and total societal costs were equivalent at general medical and surgical sites and significantly lower at neuro-

psychiatric sites. Thus, IPCC is a cost-effective treatment that merits implementation among high users of hospital services of the type sampled here.

GENERALIZABILITY: EFFECTIVENESS VS EFFICACY

Second, because the programs evaluated in this study were implemented under more typical administrative arrangements than those in previous studies, our results suggest that the benefits of IPCC can be obtained under ordinary conditions, even when programs are not implemented by their original innovators. The IPCC program was implemented without either a detailed manual or extensive and continuous training. It thus appears to be effective as well as efficacious and illustrates the specific contribution of effectiveness evaluation to psychiatric research.³⁴

IMPLEMENTATION ASSESSMENT

Third, this study illustrates the importance of tracking program implementation and, albeit unintentionally, demonstrates the relationship of faithful implementation of IPCC to program effectiveness. The 2 general medical and surgical sites that failed to implement an IPCC program also failed to demonstrate the desired effect on hospital utilization. One of them, in fact, had the opposite of the intended effect, generating considerably greater hospital use among IPCC patients than among controls. Increased hospitalization has recently been reported in another study of case management,

with the interpretation that when programs are poorly implemented, case managers tend to facilitate hospital admission for their patients because they are aware of the depth of patients' pain but have only hospital care to offer.³⁵ These data suggest that costly interventions for vulnerable populations should not be implemented without appropriate training and continuous monitoring of structure, process, and outcomes of service delivery.

TIME COURSE OF IMPROVEMENT

Fourth, we observe that even when the clinical benefits of IPCC are not apparent for as long as 18 months of treatment, they may yet appear after further treatment. This finding underlines the importance of long-term follow-up and demonstrates the potential for delay in improvement with patients with severe and persistent mental illness.

DIFFERENTIAL EFFECTIVENESS ACROSS TREATMENT SETTINGS

Fifth and finally, the study shows that the effects of IPCC programs may be quite different in settings that treat different types of patients. The IPCC-related improvement in clinical status and social adjustment was only observed at acute care facilities treating younger, higher-functioning patients with greater subjective distress—patients who were more likely to have the capacity and motivation for change.

Total cost savings, in contrast, were significant only among the patients treated at neuropsychiatric facilities—older patients who had longer hospital stays. A study of deinstitutionalization at Northampton State Hospital, Northampton, Mass, also found older longer-stay patients to be least vulnerable for rehospitalization after community placement.³⁶

Perhaps the most important lesson of this study, in this respect, is that intensive, high-cost interventions must be carefully targeted at appropriate patient subgroups. In a previous article,³⁷ we showed that savings potentials are substantial only among psychiatric patients with the highest health care costs. It can not be overemphasized that all patients treated in this study were high service utilizers, representing the upper 5% to 10% of VA resource consumers. While many patients might derive clinical benefit from IPCC treatment, only those with high levels of inpatient service use are likely to show cost savings.

METHODOLOGIC ISSUES

Before concluding, several methodological limitations must be addressed. First, although raters were not part of the clinical team, their ratings were not conducted under double-blind conditions and it is possible that rater bias influenced the results. We think this is unlikely because 2 of the measures that showed significant response to IPCC treatment (community functioning and satisfaction) were based on self-report data and were therefore less subject to such bias. Second, it is possible that patients who were lost to follow-up were doing more poorly than others, artificially inflating ratings at the final interview. Follow-up rates at 24 months, however,

were quite reasonable (78% at general medical and surgical facilities 69% at neuropsychiatric facilities) and were in fact somewhat higher than at some earlier time points.

A final limitation is that the study was conducted entirely at VA facilities in the northeastern United States, in a sample of older men with limited substance abuse comorbidity, and thus may not be generalizable to the other settings. Further studies of this type are clearly needed to test the generality of our findings in other health systems and for other subgroups.

Accepted for publication November 18, 1997.

We would like to thank the many Veterans Affairs staff members (program directors, clinical, clerical, and evaluation staff) and the hundreds of veterans at Intensive Psychiatric Community Care demonstration sites who made this project possible. In particular, we are grateful for the support of the following people: M. Hahn, ACSW; N. Paziienza, PhD; and M. Coppola, ACSW, Albany, NY; N. Denton, MSW; J. Winarski, LICSW; D. Bitman, PhD; D. Chew, MSW; and W. Penk, PhD; Boston, Mass; M. Tsuang, MD, PhD; J. Simpson, PhD; E. Brown, PhD; G. Lockwood, MSS; N. Noel, PhD; J. O'Connor, OTR; M. Hastings-Procaccino, MSW; and J. Spoltore, PhD, Brockton/West Roxbury, Mass; D. Heimowitz, MSW; D. Harris-Cobbinah, CNS; S. Rochester, CNS; S. Gentile, MA; and E. Mayer, BSW, Bronx, NY; S. Krishnaswamy, MD; J. King; J. Jordan, MSW; C. Shams, RN; L. Smith, MSW; C. Glair, RN; and R. Sullivan, MSW, Buffalo, NY; B. Lewis, RN; R. Linares, MD; D. Fogarty; R. Davis, MS; and T. Roach, BA, Canandaigua, NY; A. Russo, MSW; E. Garrett, RN; and J. Ammazalorso, Montrose, NY; L. Lantinga, PhD; G. Kline, MA; and M. Baker, Syracuse, NY; L. Harkness, PhD; J. Brady, LPN; D. Deegan, MSW; B. Johnson, MSN; C. Sweeney, RN; H. Spencer, MA; and M. Cinquanta, BA, West Haven, Conn. We also acknowledge R. Baldino; D. Thompson, BS; B. Zigler, MS; A. Ackles, BS; and D. Showalter, BS, of the Northeast Program Evaluation Center, West Haven, Conn, for critical assistance with data preparation and analysis. Philip Leaf, PhD, and Robert Milstein, MD, PhD, assisted with the design of the study. Alan Fontana, PhD; Donald Hedeker, PhD; Marilyn Stolar; and Weichun Xu, PhD, assisted with statistical analysis and Linda Frisman, PhD, and Henry Dove, PhD, with cost estimation methods. Finally, we are grateful to Paul Errera, MD, VA Central Office; Donald Burnette, former Director, VA Eastern Region; Al Washko, former VA Region 1 Director; and Ted Stefos, PhD, for their continuous encouragement and support. Boris Astrachan, MD, provided valuable advice at many junctures. Elizabeth Brown, PhD; David Gittleson, DSW; Terrence Keane, PhD; Stephen Price, MD; and Ming Tsuang, MD, PhD, served on the advisory Northeast Regional Mental Health Task Force during the implementation of this project.

Reprints: Robert Rosenheck, MD, Northeast Program Evaluation Center, 950 Campbell Ave, West Haven, CT 06516 (e-mail: Robert.Rosenheck@Yale.edu).

REFERENCES

1. Bond GR, Miller LD, Krumwied RD, Ward RS. Assertive case management in three CMHCs: a controlled study. *Hosp Community Psychiatry*. 1989;39:411-418.
2. Hoult J, Reynolds I. Schizophrenia: a comparative trial of community-oriented and hospital-oriented psychiatric care. *Acta Psychiatr Scand*. 1984;69:359-372.

3. Mulder R. *Evaluation of the Harbinger Program, 1982-1985*. Lansing, Mich: Michigan Department of Mental Health; 1985.
4. Stein LI, Test MA. Alternative to mental hospital treatment, I: conceptual model, treatment program, and clinical evaluation. *Arch Gen Psychiatry*. 1980;37:392-397.
5. Wasylenki DA, Goering PN, Lancee WJ, Ballantyne R, Farkas M. Impact of a case manager program on psychiatric aftercare. *J Nerv Ment Dis*. 1985;173:303-308.
6. Weisbrod B, Test MA, Stein LI. Alternative to mental hospital treatment, II: economic cost-benefit analysis. *Arch Gen Psychiatry*. 1980;37:400-405.
7. Bond GR, McDonell EC, Miller LD. Assertive community treatment and reference groups: an evaluation of their effectiveness for young adults with serious mental illness and substance abuse problems. *Psychosoc Rehabil J*. 1991;15:31-43.
8. McFarlane WR, Stastny P, Deakins S. Family-aided assertive community treatment: a comprehensive rehabilitation and intensive case management approach for persons with schizophrenic disorders. *New Dir Ment Health Serv*. 1992;53:43-54.
9. Curtis JL, Millman EJ, Streuning E, D'Ercole A. Effect of case management on rehospitalization and utilization of ambulatory care services. *Hosp Community Psychiatry*. 1992;43:895-899.
10. Franklin J, Solovitz B, Mason M, Clemons JR, Miller GE. An evaluation of case management. *Am J Public Health*. 1987;77:674-678.
11. Rossler W, Loffler W, Fatkenheuer B, Reicher-Rossler A. Does case management reduce the rehospitalization rate? *Acta Psychiatr Scand*. 1992;86:445-449.
12. Scott JE, Dixon LB. Assertive community treatment and case management for schizophrenia. *Schizophr Bull*. 1995;21:657-668.
13. Burns BJ, Santos AB. Assertive community treatment: an update of randomized trials. *Psychiatr Serv*. 1995;46:669-675.
14. Olfson M. Assertive community treatment: an evaluation of the experimental evidence. *Hosp Community Psychiatry*. 1990;41:634-641.
15. Bachrach LL. Overview: model programs for chronic mental patients. *Am J Psychiatry*. 1980;137:1023-1031.
16. Rosenheck R, Neale M, Leaf P, Milstein R, Frisman L. Multisite experimental cost study of intensive psychiatric community care. *Schizophr Bull*. 1995;21:129-140.
17. Test MA. Continuity of care in community treatment. *New Dir Ment Health Serv*. 1979;2:15-23.
18. McGrew JH, Bond GR, Dietzen LL. Measuring the fidelity of implementation of a mental health program model. *J Consult Clin Psychol*. 1994;62:670-678.
19. Teague GB, Drake RE, Ackerson TH. Evaluating use of continuous treatment teams for persons with mental illness and substance abuse. *Psychiatr Serv*. 1995;46:689-695.
20. Teague GB. Assertive community treatment (ACT) implementation measure and analyses. Paper presented at: Annual Meeting of the Research Institute of the National Association of State Mental Health Program Directors; February 13, 1996; Bethesda, Md.
21. Overall JE, Gorham DR. The brief psychiatric rating scale. *Psychol Rep*. 1962;10:799-812.
22. Derogatis LR, Spencer N. *The Brief Symptom Index: Administration, Scoring and Procedure Manual*. Baltimore, Md: Johns Hopkins University Press; 1982.
23. McClellan T, Luborsky L, Woody G, O'Brien C. An improved diagnostic evaluation instrument for substance abuse patients: the Addiction Severity Index. *J Nerv Ment Dis*. 1980;168:26-33.
24. Endicott J, Spitzer RL, Fleiss JL, Cohen J. The Global Assessment Scale. *Arch Gen Psychiatry*. 1976;33:766-771.
25. Rosenheck RA, Neale MS, Frisman LK. Issues in estimating the cost of innovative mental health programs. *Psychiatr Q*. 1995;66:9-31.
26. Rosenheck RA, Frisman LK, Neale MS. Estimating the capital component of mental health care costs in the public sector. *Adm Policy Ment Health*. 1994;21:493-509.
27. Frisman LK, Rosenheck RA. How transfer payments are treated in cost-effectiveness and cost-benefit analysis. *Adm Policy Ment Health*. 1996;23:533-546.
28. Schobel BD. Administrative expenses under OASDI. *Soc Secur Bull*. 1981;44:21-28.
29. US Department of Justice. Office of Justice Programs, Bureau of Justice Statistics. *Justice Expenditures and Employment in the United States, 1988*. Washington, DC: US Government Printing Office; 1991. NCJ-125619.
30. Maguire K, Flanagan T, eds. US Department of Justice, Office of Justice Programs, Bureau of Justice Statistics. *Sourcebook of Criminal Justice Statistics: 1990*. Washington, DC: US Government Printing Office; 1991. NCJ-130580.
31. Gibbons RD, Hedeker D, Elkin I. Some conceptual and statistical issues in analysis of longitudinal psychiatric data. *Arch Gen Psychiatry*. 1993;50:739-750.
32. Gold MR, Siegel JE, Russell LB, Weinstein MC. *Cost-Effectiveness in Health and Medicine*. New York, NY: Oxford University Press; 1996.
33. O'Brien BJ, Drummond MF, Labelle R, Willan A. In search of power and significance: issues in the design and analysis of stochastic cost-effectiveness studies in health care. *Med Care*. 1994;32:150-163.
34. Salive ME, Mayfield JA, Weissman NW. Patient outcomes research teams and the agency for health care policy research. *Health Serv Res*. 1990;25:697-708.
35. D'Ercole A, Streuning E, Curtis JL, Millman EJ, Morris A. Effects of diagnosis, demographic characteristics, and case management on rehospitalization. *Psychiatr Serv*. 1997;48:682-688.
36. Geller JL, Fisher WH, Simon LJ, Wirth Cauchon JL. Second-generation deinstitutionalization, II: impact of Brewster v Dukakis on correlates of community and hospital utilization. *Am J Psychiatry*. 1990;147:988-993.
37. Rosenheck RA, Massari L, Frisman LK. Who should receive high cost mental health treatment and for how long? issues in the rationing of mental health care. *Schizophr Bull*. 1993;19:843-852.

ARCHIVES Circulation

The ARCHIVES is available by request to nonfederal physicians in the United States (50 states and Washington, DC) whose official American Medical Association masterfile record shows a primary specialty of psychiatry or child psychiatry in an office- or hospital-based practice as a staff physician, resident in training beyond the first year, or clinical fellow.

If you meet the above qualification criteria and are not currently receiving the ARCHIVES and would like to receive it each month, you must complete a free subscription request card. To receive a request card, please write to Kathryn Osten, American Medical Association, Circulation Processing Department, 515 N State St, Chicago, IL 60610 (fax 312-464-5831). A subscription request card will be sent to you in response. If you are a resident or fellow, please include verification of your training program and a complete mailing address.