

## Supplementary Online Content

Ebbert JO, Hatsukami DK, Croghan I, et al. Combination varenicline and bupropion SR for tobacco dependence treatment in cigarette smokers: a randomized trial. *JAMA*. doi:10.1001/jama.2013.283185

### **eTable .** Study Schedule

This supplementary material has been provided by the authors to give readers additional information about their work.

**eTable. Study Schedule**

The study schedule is represented below.

	Screening Phase			Medication Phase							Follow-Up Phase				
	Phone Session	Info Session	Baseline	3	4	5	6	7	8	9	10	11	12	13	14
Visit No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Visit Week	-2	-1	0	1	2	4	6	8	10	12	13	14	16	26	52
Visit Type	P	C	C	P	C	C	C	C	C	C	P	C	P	C	C
<b>Visit Window (in days)<sup>1</sup></b>			± 3	± 3	± 3	± 7	± 7	± 7	± 7	± 7	± 3	± 3	± 7	± 14	± 14
Quit Date				X <sup>2</sup>											
Informed Consent		X													
Inclusion/Exclusion	X	X	X												
Physical exam			X												
Vitals		X	X		X	X	X	X	X	X		X		X	X
Medical History		X	X												
Demographics		X													
FTND <sup>3</sup>			X												
BDI-II <sup>4</sup>		X			X	X		X				X		X	X
C-SSRS <sup>5</sup>			X		X	X		X				X		X	X
Expectancies Questionnaire			X												
mCEQ <sup>6</sup>			X		X	X	X	X	X	X					
SEQ-12 <sup>7</sup>			X												
Daily Diary (w/ MNWS-R <sup>8</sup> )		X	X	X	X	X			X	X	X	X			
Urine Pregnancy Test		X					X			X					
Study Medication Dispensed			X		X	X	X	X	X						
Tobacco Outcome Measure		X	X	X	X	X	X	X	X	X	X	X	X	X	X
Concomitant Therapies	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Adverse Events			X	X	X	X	X	X	X	X	X	X	X	X	X
Behavioral Counseling			X	X	X	X	X	X	X	X	X	X	X	X	X
Expired CO <sup>9</sup>		X	X		X	X	X	X	X	X		X		X	X
End of Study Assessment Questionnaire															X

1 - If a subject cannot reschedule their visit within the study visit window, that visit is categorized as “missed”. A subject is allowed 3 consecutive missed visits before “drop” procedure is commenced. Please see NRP SOP for details of “drop” procedure.

2 - TQD is Day # 8 of varenicline + bupropion SR (active) or varenicline + bupropion SR (identical placebo)

3 - FTND = Fagerström Test for Nicotine Dependence

4 - BDI-II = Beck Depression Inventory-II

5 - C-SSRS = Columbia Suicide Severity Rating Scale

6 - mCEQ = modified Cigarette Evaluation Questionnaire (CEQ). Completed by non-abstainers.

7 - SEQ-12 = Self-Efficacy Questionnaire

8 - MNWS-R = Minnesota Nicotine Withdrawal Scale - Revised

9 - CO = An expired air CO of less than or equal to 8 ppm will be used to verify self-reported smoking abstinence.