

Scan for Author
Video Interview

Prevalence of and Risk Factors for Autopsy-Determined Atherosclerosis Among US Service Members, 2001-2011

Bryant J. Webber, MD

Peter G. Seguin, MD, MPH

Daniel G. Burnett, MD, MPH

Leslie L. Clark, PhD, MS

Jean L. Otto, DrPH, MPH

AGE-ADJUSTED ISCHEMIC HEART disease mortality rates in the United States have declined by 72% since their peak in 1968^{1,2}—a phenomenon attributed about equally to reduction in risk factors and expansion of therapies.^{3,4} These preventive and therapeutic advancements derive from an improved elucidation of the disease process. An early breakthrough in the understanding of the natural history of atherosclerotic heart disease was achieved in 1953, when Enos and colleagues⁵ at the Armed Forces Institute of Pathology reported a 77% prevalence of coronary atherosclerosis among US soldiers killed in the Korean War. By demonstrating anatomically that atherosclerosis affected a large proportion of young individuals without clinical evidence of heart disease, their study revolutionized the understanding of the onset and progression of cardiovascular disease.⁶ A follow-up report in the Vietnam War era,⁷ along with a number of autopsy studies in the civilian population,⁸⁻¹⁴ provided additional evidence that the onset of atherosclerosis may occur at an early age.

For editorial comment see p 2624.

Author Video Interview available at www.jama.com.

Context Autopsies of US service members killed in the Korean and Vietnam wars demonstrated that atherosclerotic changes in the coronary arteries can appear early in the second and third decades of life, long before ischemic heart disease becomes clinically apparent.

Objective To estimate the current prevalence of coronary and aortic atherosclerosis in the US armed forces.

Design, Setting, and Participants Cross-sectional study of all US service members who died of combat or unintentional injuries in support of Operations Enduring Freedom and Iraqi Freedom/New Dawn between October 2001 and August 2011 and whose cardiovascular autopsy reports were available at the time of data collection in January 2012. Prevalence of atherosclerosis was analyzed by various demographic characteristics and medical history. Classifications of coronary atherosclerosis severity were determined prior to data analysis and designed to provide consistency with previous military studies: minimal (fatty streaking only), moderate (10%-49% luminal narrowing of ≥ 1 vessel), and severe ($\geq 50\%$ narrowing of ≥ 1 vessel).

Main Outcome Measures Prevalence of coronary and aortic atherosclerosis in the US armed forces and by age, sex, self-reported race/ethnicity, education, occupation, service branch and component, military rank, body mass index at military entrance, and *International Classification of Diseases, Ninth Revision, Clinical Modification*, diagnoses of cardiovascular risk factors.

Results Of the 3832 service members included in the analysis, the mean age was 25.9 years (range, 18-59 years) and 98.3% were male. The prevalence of any coronary atherosclerosis was 8.5% (95% CI, 7.6%-9.4%); severe coronary atherosclerosis was present in 2.3% (95% CI, 1.8%-2.7%), moderate in 4.7% (95% CI, 4.0%-5.3%), and minimal in 1.5% (95% CI, 1.1%-1.9%). Service members with atherosclerosis were significantly older (mean [SD] age, 30.5 [8.1] years) than those without (mean [SD] age, 25.3 [5.6] years; $P < .001$). Comparing atherosclerosis prevalence among with those with no cardiovascular risk factor diagnoses (11.1% [95% CI, 10.1%-12.1%]), there was a greater prevalence among those with a diagnosis of dyslipidemia (50.0% [95% CI, 30.3%-69.7%]; age-adjusted prevalence ratio [PR], 2.09 [95% CI, 1.43-3.06]), hypertension (43.6% [95% CI, 27.3%-59.9%]; age-adjusted PR, 1.88 [95% CI, 1.34-2.65]), or obesity (22.3% [95% CI, 15.9%-28.7%]; age-adjusted PR, 1.47 [95% CI, 1.10-1.96]), but smoking (14.1% [95% CI, 8.0%-20.2%]) was not significantly associated with a higher prevalence of atherosclerosis (age-adjusted PR, 1.12 [95% CI, 0.73-1.74]).

Conclusion Among deployed US service members who died of combat or unintentional injuries and received autopsies, the prevalence of atherosclerosis varied by age and cardiovascular risk factors.

JAMA. 2012;308(24):2577-2583

www.jama.com

Author Affiliations: Uniformed Services University of the Health Sciences, Bethesda, Maryland (Drs Webber and Burnett); Armed Forces Medical Examiner System, Dover Air Force Base, Delaware (Dr Seguin); Armed Forces Health Surveillance Center, Silver Spring, Maryland (Drs Clark and Otto); and Henry M.

Jackson Foundation for the Advancement of Military Medicine Inc, Bethesda (Dr Otto).

Corresponding Author: Bryant J. Webber, MD, Uniformed Services University of the Health Sciences, PMB, 4301 Jones Bridge Rd, Room A1040A, Bethesda, MD 20814 (bryant.webber@usuhs.edu).

Since the publication of these studies, health policies for children and young adults in the general population^{15,16} as well as the military population^{17,18} of the United States have been implemented to reduce the risk of cardiovascular disease associated with risk factors such as hypertension, diabetes, cholesterol, and smoking. We conducted a cross-sectional study to assess the current prevalence of autopsy-determined atherosclerosis among US service members who died in support of combat operations.

METHODS

Every US service member who died in support of Operations Enduring Freedom (OEF) or Iraqi Freedom/New Dawn (OIF/OND) between October 2001 and August 2011 (OIF began in March 2003 and was renamed OND in September 2010) was included in this cross-sectional study. To maintain continuity with previous military studies, those with a manner of death labeled as “suicide,” “natural,” or “undetermined” were excluded. Active duty service members and activated reservists of all service branches (ie, Air Force, Army, Coast Guard, Marine Corps, and Navy) were eligible for entry. This study was approved by the Office of Research at the Uniformed Services University of the Health Sciences; as deceased persons research, it did not require institutional review board approval.

Autopsy, Demographic, and Diagnostic Data Sources

We obtained autopsy reports from the Armed Forces Medical Examiner Tracking System (AFMETS) and demographic and medical encounter data from the Defense Medical Surveillance System (DMSS). AFMETS, managed by the Mortality Surveillance Division of the Armed Forces Medical Examiner System, Dover, Delaware, is the central registry for US military fatalities.¹⁹ DMSS, managed by the Armed Forces Health Surveillance Center, Silver Spring, Maryland, is the central repository for medical surveillance data

for the armed forces, which contains longitudinal data on service members, including almost complete ascertainment of all medical encounters since 1990, as well as demographic and deployment data.²⁰

Demographic data included age, sex, race/ethnicity, highest education level achieved before death, occupation, service branch, active or reserve component, and military rank. On entering the military, service members self-select their race/ethnicity from a list of fixed categories; race/ethnicity was reported because of a known association with atherosclerosis.^{8,9,13} Medical encounter data included body mass index (BMI, calculated as weight in kilograms divided by height in meters squared) at military entrance and history of atherosclerosis, ischemic heart disease, and major cardiovascular risk factors (ie, hypertension, dyslipidemia, obesity, diabetes mellitus, impaired fasting glucose, and smoking) based on *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*, codes. Service members were considered to have a history of any of these diseases or risk factors if they had at least 1 inpatient encounter or 2 outpatient encounters within 180 days with a defining diagnosis in any diagnostic position. All risk factor data were recorded prior to death.

Evaluation of Atherosclerotic Lesions

All US military deaths associated with combat or unintentional injuries in support of OEF or OIF/OND—whether the service member died in theater or after evacuation—required a comprehensive autopsy performed by a board-certified pathologist. For those who died in theater, autopsies were conducted at Dover Port Mortuary, Dover Air Force Base, Delaware, and were typically completed within 72 hours of death. For those who were wounded, medically evacuated, and subsequently died of wounds, autopsies were often conducted at the local military treatment facility, including National Naval Medical Center, Bethesda, Mary-

land (n=32); Brooke Army Medical Center, Fort Sam Houston, Texas (n=28); and Walter Reed Army Medical Center, Washington, DC (n=20). Early in OEF, some autopsies were conducted at Landstuhl Regional Medical Center, Germany (n=68).

All autopsies included an examination of the cardiovascular system. After removal of the heart from the pericardial sac and anterior chest chamber, the major coronary arteries (ie, left main, left anterior descending, circumflex, and right coronary) were inspected in the fresh state. The inferior portions of the ventricles were then sectioned and the valves opened according to the physiologic sequence of blood flow. The aorta was opened longitudinally, beginning typically at the aortic valve and carried distally through the bifurcation and into both common iliac arteries. The pathologist documented the location and degree of all atherosclerotic lesions based on gross examination.

To provide consistency with previous military studies, we determined the following classifications of coronary atherosclerosis severity prior to data analysis: minimal (fatty streaking only), moderate (10%-49% luminal narrowing of ≥ 1 vessel), and severe ($\geq 50\%$ narrowing of ≥ 1 vessel). We also documented the presence of aortic atherosclerosis.

Statistical Analyses

We used summary statistics to describe the population. We evaluated the prevalence of atherosclerosis by various demographic characteristics and diagnostic histories and reported unadjusted and Mantel-Haenszel age-adjusted prevalence ratios (PRs) with 95% confidence intervals. We used logistic regression to evaluate risk factor combinations and to assess for effect modification, reporting odds ratios (ORs). To compare included and excluded participants, we used χ^2 and Fisher exact tests of homogeneity (for categorical variables) and unpaired *t* tests (for continuous variables).

All analyses were performed with Stata/IC version 11.2 (StataCorp). *P* values less than .05 were considered statistically significant; all *P* values were based on 2-sided tests.

RESULTS

Of the 6191 fatalities in the case identification period, 4143 cases were identified by the Armed Forces Medical Examiner System (532 were initially excluded because the manner of death was other than combat or unintentional injury and 1516 because of injury to the cardiovascular system precluding complete autopsy). On review of the reports, we excluded an additional 311 cases (235 with no autopsy on file, 69 with damage preventing coronary examination, and 7 indicating organ donation). Included in the analysis were 3832 service members aged 18 to 59 years, of whom 98.3% were male.

Included and excluded participants were mostly homogenous demographically, although the excluded were younger (mean [SD] ages, 25.9 [6.2] vs 27.0 [6.6], respectively; *P* = .002) and had a smaller proportion of college graduates (10.0% vs 14.8%; *P* = .008); the groups had similar mean (SD) entrance BMIs (included, 24.3 [3.6] vs excluded, 24.4 [3.2]; *P* = .85) and similar proportions with 1 or more cardiovascular risk factor diagnoses (included, 8.5% vs excluded, 10.9%; *P* = .14).

Atherosclerosis Prevalence

The overall prevalence of coronary or aortic atherosclerosis was 12.1% (95% CI, 11.1%-13.1%). The prevalence of coronary and aortic atherosclerosis was 2.1% (95% CI, 1.7%-2.6%). The prevalence of any coronary atherosclerosis was 8.5% (95% CI, 7.6%-9.4%); severe coronary atherosclerosis was present in 2.3% (95% CI, 1.8%-2.7%), moderate in 4.7% (95% CI, 4.0%-5.3%), and minimal in 1.5% (95% CI, 1.1%-1.9%). The prevalence of any aortic atherosclerosis was 5.7% (95% CI, 5.0%-6.5%).

Factors Associated With Prevalent Atherosclerosis

Age consistently produced the strongest association with prevalent atherosclerosis. Service members with atherosclerosis (mean [SD] age, 30.5 [8.1] years) were approximately 5 years older than those without (mean [SD] age, 25.3 [5.6] years; *P* < .001); those aged 40 years and older had about 7 times the prevalence of disease as compared with those aged 24 years and younger (45.9% vs 6.6%; unadjusted PR, 6.95; 95% CI, 5.49-8.80) (TABLE 1).

Lower education level and higher military entrance BMI were significantly associated with prevalent atherosclerosis, after adjusting for age. As compared with those who completed high school or less, those who completed at least some college had lower prevalence of disease (11.2% vs 13.8%; age-adjusted PR, 0.65; 95% CI, 0.45-0.92); this association did not extend to college or higher-level graduates. As compared with those with a normal BMI (18.5-24.9) on military entrance, those with a BMI in the overweight (25.0-29.9) or obese (≥ 30.0) range had a significantly higher prevalence of atherosclerosis (overweight: 12.7% vs 7.6%; age-adjusted PR, 1.39; 95% CI, 1.10-1.75; obese: 15.8% vs 7.6%; age-adjusted PR, 1.87; 95% CI, 1.35-2.60). Age-adjusted PRs by sex, race/ethnicity, occupation, service branch, active/reserve component, and military rank were not significant (Table 1).

Age-adjusted atherosclerosis prevalence was associated with several diagnoses (TABLE 2). As compared with those with no major cardiovascular risk factor diagnoses, those with a diagnosis of dyslipidemia (50.0% vs 11.1%; age-adjusted PR, 2.09; 95% CI, 1.43-3.06), hypertension (43.6% vs 11.1%, age-adjusted PR, 1.88; 95% CI, 1.34-2.65), or obesity (22.3% vs 11.1%; age-adjusted PR, 1.47; 95% CI, 1.10-1.96) had a significantly higher prevalence of atherosclerosis. Of all risk factor combinations, only obesity and hypertension (*n* = 8) resulted in a statistically significant association after adjusting for

age (50.0% vs 11.1%; age-adjusted OR, 3.14; 95% CI, 1.54-6.44).

Of the 464 service members with atherosclerosis at autopsy, only 2 (0.5%) had been clinically diagnosed with atherosclerosis and none had been diagnosed with ischemic heart disease.

COMMENT

The prevalence of autopsy-determined coronary atherosclerosis among US service members was 8.5% in our study population. Older age; lower educational level; higher BMI at military entrance; and prior diagnoses of dyslipidemia, hypertension, and obesity were associated with a higher prevalence of atherosclerosis.

This prevalence demonstrates a steep decline from the rates of 77% noted in the Korean War⁵ and 45% in the Vietnam War⁷ (TABLE 3)—a decline that can be explained by either real or artifactual phenomena. Within the latter category are several methodological discrepancies among the 3 studies. First, while the Vietnam study employed angiography on perfusion-fixed coronary arteries, the Korean and current studies relied on macroscopic visualization, which is likely less sensitive to detect the earliest atherosclerotic changes. Second, the Korean study grouped fibrous thickening with fatty streaking as atherosclerosis, but it is now understood that intimal thickening may occur physiologically as a non-atherosclerotic adaptation.²³ Third, while the Vietnam study used an absolute scale for grading plaque size, the present study enlisted the percentage-obstructed method used in the Korean study. Fourth, the autopsies in the earlier studies were conducted for scientific purposes, such as the study of wound ballistics, but the autopsies reviewed for this study were ordered to be conducted routinely rather than selected by specific patterns of injury; our method is therefore likely to have minimized expectation bias.²⁴

Selection biases may also attenuate direct comparisons of these studies. In light of the possibility that recent military volunteers have healthier cardio-

vascular profiles than their conscripted counterparts in the Korean and Vietnam wars, the potential emerges for a self-selection or “healthy warrior” bias when juxtaposing studies conducted during and after the draft era. In addition, advancements in battlefield medicine and technology may have saved the lives of wounded soldiers with atherosclerosis, who, by virtue of compromised perfusion, would have died in earlier conflicts; the prevalence of

Table 1. Prevalence of Coronary and/or Aortic Atherosclerosis by Demographic Characteristic

Characteristics	No. of Patients With Atherosclerosis/Total No.	Atherosclerosis Prevalence, % (95% CI)	Prevalence Ratio (95% CI)	
			Unadjusted	Age-Adjusted
Total	464/3832	12.1 (11.1-13.1)		
Age, y				
<25	135/2047	6.6 (5.5-7.7)	1 [Reference]	
25-29	103/931	11.1 (9.0-13.1)	1.68 (1.31-2.14)	
30-39	154/697	22.1 (19.0-25.2)	3.35 (2.70-4.15)	
≥40	72/157	45.9 (38.0-53.7)	6.95 (5.49-8.80)	
Sex				
Female	5/67	7.5 (1.0-13.9)	1 [Reference]	1 [Reference]
Male	459/3765	12.2 (11.1-13.2)	1.63 (0.70-3.81)	1.71 (0.75-3.88)
Race/ethnicity				
White, non-Hispanic	328/2787	11.8 (10.6-13.0)	1 [Reference]	1 [Reference]
Black, non-Hispanic	40/311	12.9 (9.1-16.6)	1.09 (0.80-1.49)	0.92 (0.68-1.24)
Hispanic	52/424	12.3 (9.1-15.4)	1.04 (0.79-1.37)	1.10 (0.85-1.43)
Asian/Pacific Islander	22/130	16.9 (10.4-23.5)	1.44 (0.97-2.13)	1.43 (0.98-2.09)
Other ^a	22/180	12.2 (7.4-17.1)	1.04 (0.69-1.56)	0.84 (0.57-1.25)
Education				
High school or less	354/3159	11.2 (10.1-12.3)	1 [Reference]	1 [Reference]
Some college	31/225	13.8 (9.2-18.3)	1.23 (0.87-1.73)	0.65 (0.45-0.92)
College	54/343	15.7 (11.9-19.6)	1.40 (1.08-1.83)	0.84 (0.64-1.10)
Postcollege	13/45	28.9 (15.1-42.7)	2.58 (1.61-4.12)	0.73 (0.44-1.21)
Unknown	12/60	20.0 (9.6-30.4)	1.78 (1.07-2.99)	1.28 (0.78-2.10)
Occupation				
Combat	283/2503	11.3 (10.1-12.5)	1 [Reference]	1 [Reference]
Health care	14/140	10.0 (5.0-15.0)	0.88 (0.53-1.47)	0.75 (0.46-1.25)
Administrative/supply	68/423	16.1 (12.6-19.6)	1.42 (1.11-1.81)	1.10 (0.87-1.39)
Other ^b	99/766	12.9 (10.5-15.3)	1.14 (0.92-1.42)	1.05 (0.86-1.30)
Service branch				
Marine Corps	74/930	8.0 (6.2-9.7)	1 [Reference]	1 [Reference]
Army	359/2745	13.1 (11.8-14.3)	1.64 (1.29-2.09)	1.15 (0.90-1.48)
Navy	16/90	17.8 (9.7-25.8)	2.23 (1.36-3.67)	1.37 (0.75-2.54)
Air Force	15/66	22.7 (12.3-33.1)	2.86 (1.74-4.69)	1.38 (0.77-2.50)
Coast Guard	0/1			
Component				
Active	356/3250	11.0 (9.9-12.0)	1 [Reference]	1 [Reference]
Reserve	108/582	18.6 (15.4-21.7)	1.69 (1.39-2.06)	1.04 (0.85-1.28)
Military rank ^c				
Junior enlisted	171/2179	7.8 (6.7-9.0)	1 [Reference]	1 [Reference]
Junior officer	38/284	13.4 (9.4-17.4)	1.71 (1.23-2.37)	0.91 (0.61-1.35)
Senior enlisted	233/1309	17.8 (15.7-19.9)	2.27 (1.88-2.72)	1.10 (0.87-1.40)
Senior officer	22/60	36.7 (24.1-49.2)	4.67 (3.25-6.71)	0.94 (0.49-1.81)
Military entrance BMI ^d				
<18.5	3/62	4.8 (0.0-10.3)	0.64 (0.21-1.95)	0.53 (0.17-1.68)
18.5-24.9	125/1653	7.6 (6.3-8.8)	1 [Reference]	1 [Reference]
25.0-29.9	156/1229	12.7 (10.8-14.6)	1.68 (1.34-2.10)	1.39 (1.10-1.75)
≥30.0	41/259	15.8 (11.4-20.3)	2.09 (1.51-2.90)	1.87 (1.35-2.60)
Not on file	139/629	22.1 (18.8-25.3)	2.92 (2.34-3.65)	1.41 (1.07-1.87)

Abbreviation: BMI, body mass index.

^aAmerican Indian, Alaskan Native, multiracial, unknown, and declined to report.

^bCommunications, intelligence, electrical, mechanical, engineering, maintenance, scientist, and unknown.

^cJunior enlisted: E-1 through E-4; junior officer: O-1 through O-3 and W-1 through W-3; senior enlisted: E-5 through E-9; senior officer: O-4 through O-6 and W-4 through W-5.

^dCalculated as weight in kilograms divided by height in meters squared.

atherosclerosis at autopsy, in other words, may appear higher in the Korean and Vietnam wars because of unequal combat casualty care. Demographic and socioeconomic changes in the military population over this period may also account for the decline in atherosclerosis; because these data are unavailable for the Korean and Vietnam war studies, comparisons should be made cautiously. Finally, the smaller proportion of combat fatalities autopsied in the Korean and Vietnam war studies—less than 1% in both cases, as compared with more than 60% in the current study—may have resulted in less representative samples.

In light of historic epidemiology, these artifactual explanations are matched by compelling factual alter-

natives. Autopsies conducted for general aviation mishaps over the past 5 decades have generally demonstrated declines in any and severe coronary atherosclerosis, although shifting demographics complicate interpretation of these findings.²⁵⁻²⁷ Less subjectively, prevalence of cardiovascular risk factors in the general US population,²⁸⁻³² with the exceptions of obesity³³ and diabetes,³⁴ are trending down. Although similar trend data are not available in the military, a recent study of more than 280 000 Air Force members found lower prevalences of hypertension, dyslipidemia, and diabetes as compared with the general US population.³⁵ It is not unreasonable to assume that the prevalence of these risk factors among deployed US troops has declined since

Korea and Vietnam. Smoking rates have also decreased in the military, often surpassing absolute rate reductions in the civilian population.^{31,36} The majority (51%) of service members smoked cigarettes in 1980, but by 1998 this figure had decreased to 30%.³⁶ This 40% reduction in smoking prevalence within the military since the end of the Vietnam War likely contributed to our results.

Although the military culture tolerated or even promoted tobacco use in earlier eras, more recent Department of Defense (DoD) policy initiatives have aimed to reduce smoking. These have included discontinuation of cigarettes in K- and C-rations, efforts to deglamorize tobacco, and making Navy ships smoke-free.³⁷ Current DoD Instruc-

Table 2. Prevalence of Coronary and/or Aortic Atherosclerosis by ICD-9-CM Diagnoses of Major Cardiovascular Risk Factors

ICD-9-CM Diagnosis	No. of Patients With Atherosclerosis/Total No.	Atherosclerosis Prevalence, % (95% CI)	Prevalence Ratio (95% CI)	
			Unadjusted	Age-Adjusted
None	389/3506	11.1 (10.1-12.1)	1 [Reference]	1 [Reference]
Obesity	37/166	22.3 (15.9-28.7)	2.01 (1.49-2.71)	1.47 (1.10-1.96)
Smoking	18/128	14.1 (8.0-20.2)	1.27 (0.82-1.96)	1.12 (0.73-1.74)
Hypertension	17/39	43.6 (27.3-59.9)	3.93 (2.72-5.68)	1.88 (1.34-2.65)
Dyslipidemia	14/28	50.0 (30.3-69.7)	4.51 (3.08-6.60)	2.09 (1.43-3.06)
Diabetes/IFG	2/10	20.0 (0.0-50.2)	1.80 (0.52-6.25)	0.58 (0.17-1.97)
Obesity and hypertension	4/8	50.0 (5.3-94.7)	4.51 (2.24-9.07) ^a	3.14 (1.54-6.44) ^a

Abbreviations: ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IFG, impaired fasting glucose.

^aReflects odds ratio.

Table 3. Coronary Autopsy Studies in the US Armed Forces: Korean War, Vietnam War, and OEF and OIF/OND

	Korean War ^{5,21}	Vietnam War ⁷	OEF and OIF/OND
US deaths, No.	36 574 ^a	58 220 ^a	6191 ^b
Autopsies included in the study, No. (% of total deaths)	300 (0.8)	105 (0.2)	3832 (61.9)
Mean (SD) age, y	22.1 (NR)	22.1 (4.4)	25.9 (6.6)
Race, % white	NR	86.7	72.7
Coronary lesion grading			
Minimal	Fibrous thickening or streaking causing insignificant narrowing	Fibrous thickening or single plaques <5 mm in greatest diameter	Streaking causing insignificant luminal narrowing
Moderate	10%-49% narrowing	Single plaques <1 cm	10%-49% narrowing
Severe	≥50% narrowing	Single plaques >1 cm or confluent smaller plaques	≥50% narrowing
Prevalence of coronary atherosclerosis, %			
Minimal	35	NR	1.5
Moderate	27	NR	4.7
Severe	15	5	2.3
Any	77	45	8.5

Abbreviations: OEF, Operation Enduring Freedom; OIF, Operation Iraqi Freedom; OND, Operation New Dawn; NR, not reported.

^aPer Congressional Research Service.²²

^bDuring case identification period of October 2001 through August 2011, per Armed Forces Medical Examiner System.

tions seek to bring cigarette prices on military installations to within 5% of the local economy³⁸ and to make DoD facilities smoke-free.³⁹ Service-specific implementation of these policies have led to such initiatives as banning cigarettes from Air Force medical campuses⁴⁰ and prohibiting smoking on submarines.⁴¹ Despite these policy initiatives and improvements, current military smoking rates (30.5%)³⁶ continue to exceed civilian rates among similar age groups (21.4% in 18- to 24-year-olds and 23.7% in 25- to 44-year-olds).³²

The findings of this study should be interpreted cautiously in light of its other limitations. Because autopsies were performed by multiple pathologists over the span of a decade, interobserver and intersession variation may exist. Studies by the World Health Organization, however, have demonstrated high reproducibility of coronary and aortic atherosclerotic grading among medical examiners and over time.^{42,43} Based on the 2008 DoD health-related behavior survey, we also suspect that ICD-9-CM diagnoses failed to capture many cardiovascular risk factor exposures. In this survey of more than 28 000 active duty personnel, 30.5% had smoked cigarettes in the previous 30 days and 12.6% were classified as obese based on BMI.³⁶ The degree of undercapture of these and other cardiovascular risk factors would likely not vary by the presence or absence of atherosclerosis, however, resulting in a nondifferential misclassification that would bias our results toward the null.

Our findings suggest that the prevalence of atherosclerosis has declined among deployed US service members since the Korean and Vietnam wars, and the work of Ford and associates^{3,4} indicates that the prevalence of ischemic heart disease in the general population of US adults has declined over a similar time period. However, targets for further improvement remain. The 2008 DoD health-related behavior survey identifies smoking, obesity, diet, and physical activity as prime targets for public health intervention within the

military community,³⁶ and data from the National Health and Nutrition Examination Survey indicate that only 1.2% of US adults⁴⁴ achieve the 7 lifestyle metrics¹⁵ correlated with a stepwise reduction in all-cause and cardiovascular mortality.⁴⁴ To help “identify those personnel with a potential high risk for cardiovascular disease,”¹⁷ the DoD should consider automatic coding within their electronic health record for diagnoses such as obesity and smoking that are objectively measured during health care visits. Military and civilian health care systems should continue to help patients reduce their cardiovascular risk factors, beginning in childhood and continuing throughout adult life. Despite remarkable progress in prevention and treatment, cardiovascular disease remains the leading cause of death in the United States and other developed nations, and even small improvements in the prevalence of smoking and other risk factors may reduce death rates further and prolong healthy lives.⁴⁵

Author Contributions: Dr Webber had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Webber, Seguin, Burnett, Clark, Otto.

Acquisition of data: Webber, Seguin, Clark.

Analysis and interpretation of data: Webber, Seguin, Burnett, Clark, Otto.

Drafting of the manuscript: Webber, Seguin, Burnett, Clark.

Critical revision of the manuscript for important intellectual content: Webber, Seguin, Burnett, Clark, Otto.

Statistical analysis: Webber, Burnett.

Administrative, technical, or material support: Seguin, Otto.

Study supervision: Seguin, Burnett, Clark, Otto.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Funding/Support: This study was supported by the Uniformed Services University of the Health Sciences (USUHS), Armed Forces Medical Examiner System (AFMES), and Armed Forces Health Surveillance Center (AFHSC).

Role of the Sponsor: The sponsors had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation or review of the manuscript. USUHS, AFMES, and AFHSC approved the manuscript prior to submission.

Disclaimer: This work is the sole responsibility of the authors and does not represent the official views of USUHS, AFMES, AFHSC, or Department of Defense.

Online-Only Material: The Author Video Interview is available at <http://www.jama.com>.

Additional Contributions: We thank USUHS faculty members Roger Gibson, DVM, MPH, PhD; Ann Scher, PhD; and Elizabeth Rouse, MD, for their help with study

design, statistical analyses, and forensic pathology consultation, respectively; Steve Taubman, PhD, and Joel Gaydos, MD, MPH, of AFHSC, for data acquisition and for scientific review of the manuscript, respectively; and the study's participants, for their service. No contributors received compensation in addition to their federal government salary.

REFERENCES

- Age-adjusted death rates for 69 selected causes by race and sex using year 2000 standard population: United States, 1968-78. National Center for Health Statistics. <http://www.cdc.gov/nchs/data/dvs/mx196878.pdf>. Accessed October 12, 2012.
- Miniño AM, Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2008. *Natl Vital Stat Rep*. 2011;59(10):1-126.
- Ford ES, Ajani UA, Croft JB, et al. Explaining the decrease in US deaths from coronary disease, 1980-2000. *N Engl J Med*. 2007;356(23):2388-2398.
- Ford ES, Capewell S. Proportion of the decline in cardiovascular mortality disease due to prevention versus treatment: public health versus clinical care. *Annu Rev Public Health*. 2011;32:5-22.
- Enos WF, Holmes RH, Beyer J. Coronary disease among United States soldiers killed in action in Korea: preliminary report. *JAMA*. 1953;152(12):1090-1093.
- Strong JP. Landmark perspective: coronary atherosclerosis in soldiers: a clue to the natural history of atherosclerosis in the young. *JAMA*. 1986;256(20):2863-2866.
- McNamara JJ, Molot MA, Stremple JF, Cutting RT. Coronary artery disease in combat casualties in Vietnam. *JAMA*. 1971;216(7):1185-1187.
- Strong JP, McGill HC Jr. The natural history of coronary atherosclerosis. *Am J Pathol*. 1962;40:37-49.
- Tejada C, Strong JP, Montenegro MR, Restrepo C, Solberg LA. Distribution of coronary and aortic atherosclerosis by geographic location, race, and sex. *Lab Invest*. 1968;18(5):509-526.
- Holme I, Solberg LA, Weissfeld L, et al. Coronary risk factors and their pathway of action through coronary raised lesions, coronary stenoses and coronary death: multivariate statistical analysis of an autopsy series: the Oslo Study. *Am J Cardiol*. 1985;55(1):40-47.
- Berenson GS, Srinivasan SR, Bao W, Newman WP III, Tracy RE, Wattigney WA. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults: the Bogalusa Heart Study. *N Engl J Med*. 1998;338(23):1650-1656.
- Reed DM, MacLean CJ, Hayashi T. Predictors of atherosclerosis in the Honolulu Heart Program: I, Biologic, dietary, and lifestyle characteristics. *Am J Epidemiol*. 1987;126(2):214-225.
- Relationship of atherosclerosis in young men to serum lipoprotein cholesterol concentrations and smoking: a preliminary report from the Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. *JAMA*. 1990;264(23):3018-3024.
- Joseph A, Ackerman D, Talley JD, Johnstone J, Kupersmith J. Manifestations of coronary atherosclerosis in young trauma victims: an autopsy study. *J Am Coll Cardiol*. 1993;22(2):459-467.
- Lloyd-Jones DM, Hong Y, Labarthe D, et al; American Heart Association Strategic Planning Task Force and Statistics Committee. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2010;121(4):586-613.
- Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and

- adolescents. National Heart Lung and Blood Institute. http://www.nhlbi.nih.gov/guidelines/cvd_ped/summary.htm. Accessed November 27, 2012.
17. DoD physical fitness and body fat program: directive 1308.1: June 30, 2004. Department of Defense. <http://www.dtic.mil/whs/directives/corres/pdf/130801p.pdf>. Accessed November 27, 2012.
 18. Health promotion and disease/injury prevention: directive 1010.10: August 23, 2003. Department of Defense. <http://www.dtic.mil/whs/directives/corres/pdf/101010p.pdf>. Accessed November 27, 2012.
 19. Gardner JW, Cozzini CB, Kelley PW, et al. The Department of Defense Medical Mortality Registry. *Mil Med*. 2000;165(7)(suppl 2):57-61.
 20. Rubertone MV, Brundage JF. The Defense Medical Surveillance System and the Department of Defense serum repository: glimpses of the future of public health surveillance. *Am J Public Health*. 2002;92(12):1900-1904.
 21. Enos WF Jr, Beyer JC, Holmes RH. Pathogenesis of coronary disease in American soldiers killed in Korea. *J Am Med Assoc*. 1955;158(11):912-914.
 22. Leland A, Oboroceanu MJ. American war and military operations casualties: lists and statistics [RL32492]: February 26, 2010. Congressional Research Service. <http://www.scribd.com/doc/34858912/Congressional-Research-Service-American-War-and-Military-Operations-Casualties-Lists-and-Statistics>. Accessed November 29, 2012.
 23. Nakashima Y, Wight TN, Sueishi K. Early atherosclerosis in humans: role of diffuse intimal thickening and extracellular matrix proteoglycans. *Cardiovasc Res*. 2008;79(1):14-23.
 24. Sackett DL. Bias in analytic research. *J Chronic Dis*. 1979;32(1-2):51-63.
 25. Taneja N, Wiegmann DA. Prevalence of cardiovascular abnormalities in pilots involved in fatal general aviation airplane accidents. *Aviat Space Environ Med*. 2002;73(10):1025-1030.
 26. Booze CF Jr, Staggs CM. A comparison of post-mortem coronary atherosclerosis findings in general aviation pilot fatalities. *Aviat Space Environ Med*. 1987;58(4):297-300.
 27. Pettyjohn FS, McMeekin RR. Coronary artery disease and preventive cardiology in aviation medicine. *Aviat Space Environ Med*. 1975;46(10):1299-1304.
 28. QuickStats: average total cholesterol level among men and women aged 20-74 years: National Health and Nutrition Examination Survey, United States, 1959-1962 to 2007-2008. *MMWR Morb Mortal Wkly Rep*. 2009;58(37):1045.
 29. Burt VL, Cutler JA, Higgins M, et al. Trends in the prevalence, awareness, treatment, and control of hypertension in the adult US population: data from the health examination surveys, 1960 to 1991. *Hypertension*. 1995;26(1):60-69.
 30. Yoon SS, Ostchega Y, Louis T. Recent trends in the prevalence of high blood pressure and its treatment and control, 1999-2008. *NCHS Data Brief*. 2010;(48):1-8.
 31. Centers for Disease Control and Prevention (CDC). Tobacco use: United States, 1900-1999. *MMWR Morb Mortal Wkly Rep*. 1999;48(43):986-993.
 32. Centers for Disease Control and Prevention (CDC). Cigarette smoking among adults and trends in smoking cessation: United States, 2008. *MMWR Morb Mortal Wkly Rep*. 2009;58(44):1227-1232.
 33. Ogden CL, Fryar CD, Carroll MD, Flegal KM. Mean body weight, height, and body mass index, United States 1960-2002. *Adv Data*. 2004;(347):1-17.
 34. Long-term trends in diagnosed diabetes. CDC's Division of Diabetes Translation. http://www.cdc.gov/diabetes/statistics/slides/long_term_trends.pdf. Accessed July 24, 2012.
 35. Hatzfeld JJ, LaVeist TA, Gaston-Johansson FG. Racial/ethnic disparities in the prevalence of selected chronic diseases among US Air Force members, 2008. *Prev Chronic Dis*. 2012;9:E112.
 36. Bray RM, Pemberton MR, Hourani LL, et al. 2008 Department of Defense Survey of Health Related Behaviors Among Active Duty Military Personnel. Research Triangle Park, NC: RTI International; 2009.
 37. Joseph AM, Muggli M, Pearson KC, Lando H. The cigarette manufacturers' efforts to promote tobacco to the US military. *Mil Med*. 2005;170(10):874-880.
 38. Armed services exchange policy: instruction 1330.09: December 7, 2005. Department of Defense. <http://www.dtic.mil/whs/directives/corres/pdf/133009p.pdf>. Accessed November 28, 2012.
 39. Smoke-free DoD facilities: instruction 1010.15: January 2, 2001. Department of Defense. http://biotech.law.lsu.edu/blaw/dodd/corres/pdf/i101015_010201/i101015p.pdf. Accessed November 29, 2012.
 40. Tobacco use in the Air Force: instruction 40-102: March 26, 2012. United States Air Force. <http://www.e-publishing.af.mil/shared/media/epubs/AFI40-102.pdf>. Accessed November 28, 2012.
 41. Commander Submarine Force Atlantic, Commander Submarine Force US Pacific Fleet. *Submarine Force Tobacco Cessation Program, COMSUBLANT/COMSUBPAC NOTICE 5100*. Norfolk, VA: Department of the Navy; 2010.
 42. Zhdanov VS, Sternby NH, Vikhert AM, Galakhov IE. Development of atherosclerosis over a 25 year period: an epidemiological autopsy study in males of 11 towns. *Int J Cardiol*. 1999;68(1):95-106.
 43. Uemura K, Sternby N, Vanecek R, Vihert A, Kagan A. Grading atherosclerosis in aorta and coronary arteries obtained at autopsy: application of a tested method. *Bull World Health Organ*. 1964;31:297-320.
 44. Yang Q, Cogswell ME, Flanders WD, et al. Trends in cardiovascular health metrics and associations with all-cause and CVD mortality among US adults. *JAMA*. 2012;307(12):1273-1283.
 45. Wilkins JT, Ning H, Berry J, Zhao L, Dyer AR, Lloyd-Jones DM. Lifetime risk and years lived free of total cardiovascular disease. *JAMA*. 2012;308(17):1795-1801.