

Supplementary Online Content

Ebbert JO, Hatsukami DK, Croghan I, et al. Combination varenicline and bupropion SR for tobacco dependence treatment in cigarette smokers: a randomized trial. *JAMA*. doi:10.1001/jama.2013.283185

eTable . Study Schedule

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Study Schedule

The study schedule is represented below.

	Screening Phase			Medication Phase							Follow-Up Phase				
	Phone Session	Info Session	Baseline	3	4	5	6	7	8	9	10	11	12	13	14
Visit No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Visit Week	-2	-1	0	1	2	4	6	8	10	12	13	14	16	26	52
Visit Type	P	C	C	P	C	C	C	C	C	C	P	C	P	C	C
Visit Window (in days)¹			± 3	± 3	± 3	± 7	± 7	± 7	± 7	± 7	± 3	± 3	± 7	± 14	± 14
Quit Date				X ²											
Informed Consent		X													
Inclusion/Exclusion	X	X	X												
Physical exam			X												
Vitals		X	X		X	X	X	X	X	X		X		X	X
Medical History		X	X												
Demographics		X													
FTND ³			X												
BDI-II ⁴		X			X	X		X				X		X	X
C-SSRS ⁵			X		X	X		X				X		X	X
Expectancies Questionnaire			X												
mCEQ ⁶			X		X	X	X	X	X	X					
SEQ-12 ⁷			X												
Daily Diary (w/ MNWS-R ⁸)		X	X	X	X	X			X	X	X	X			
Urine Pregnancy Test		X					X			X					
Study Medication Dispensed			X		X	X	X	X	X						
Tobacco Outcome Measure		X	X	X	X	X	X	X	X	X	X	X	X	X	X
Concomitant Therapies	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Adverse Events			X	X	X	X	X	X	X	X	X	X	X	X	X
Behavioral Counseling			X	X	X	X	X	X	X	X	X	X	X	X	X
Expired CO ⁹		X	X		X	X	X	X	X	X		X		X	X
End of Study Assessment Questionnaire															X

1 - If a subject cannot reschedule their visit within the study visit window, that visit is categorized as “missed”. A subject is allowed 3 consecutive missed visits before “drop” procedure is commenced. Please see NRP SOP for details of “drop” procedure.

2 - TQD is Day # 8 of varenicline + bupropion SR (active) or varenicline + bupropion SR (identical placebo)

3 - FTND = Fagerström Test for Nicotine Dependence

4 - BDI-II = Beck Depression Inventory-II

5 - C-SSRS = Columbia Suicide Severity Rating Scale

6 - mCEQ = modified Cigarette Evaluation Questionnaire (CEQ). Completed by non-abstainers.

7 - SEQ-12 = Self-Efficacy Questionnaire

8 - MNWS-R = Minnesota Nicotine Withdrawal Scale - Revised

9 - CO = An expired air CO of less than or equal to 8 ppm will be used to verify self-reported smoking abstinence.