

Supplementary Online Content

Paige NM, Miake-Lye IM, Booth MS, et al. Association of spinal manipulative therapy with clinical benefit and harm for acute low back pain: systematic review and meta-analysis. *JAMA*. doi:10.1001/jama.2017.3086

eAppendix. Systematic Review Search Strategies

eTable 1. Evidence Table of 26 Randomized Clinical Trials of Spinal Manipulative Therapy for Acute Low Back Pain

eTable 2. Quality Scores of 26 Randomized Clinical Trials of Spinal Manipulation Therapy for Acute Low Back Pain

eReferences

This supplementary material has been provided by the authors to give readers additional information about their work.

eAppendix. Systematic review search strategies

SEARCH STRATEGY FOR “CHIROPRACTIC” SYSTEMATIC REVIEWS

DATABASE SEARCHED:

Cochrane Database of Systematic Reviews and Other Reviews

NO DATE OR LANGUAGE LIMITATIONS

SEARCH STRATEGY:

'chiroprac* in Title, Abstract, Keywords
Cochrane Reviews (17)
Other Reviews (44)

SEARCH STRATEGY:

"Manipulation, Spinal"

Cochrane Databased Search Strategy #2:

spine or spinal or neck or back or cervi*
and
(smt or manipulat* or chiropract*):ti,ab,kw

Dates:

2011-present,

Limit to the Cochrane Systematic Reviews, Other Reviews (DARE), Technology Assessments, and Economic Evaluations databases.

Forward search on:

Hurwitz EL, Aker PD, Adams AH, Meeker WC, Shekelle PG. Manipulation and mobilization of the cervical spine. A systematic review of the literature. Spine (Phila Pa 1976). Aug 1 1996;21(15):1746-1759; discussion 1759-1760.

2. Update Search Strategies

SPINAL MANIPULATION THERAPY – 2015 UPDATE SEARCH METHODOLOGY

DATABASE SEARCHED & TIME PERIOD COVERED:

COCHRANE CENTRAL – 1/1/2011-2/06/2017

SEARCH STRATEGY:

#1 MeSH descriptor: [Back] explode all trees

#2 MeSH descriptor: [Buttocks] this term only

#3 MeSH descriptor: [Leg] this term only

#4 MeSH descriptor: [Back Pain] explode all trees

#5 MeSH descriptor: [Back Pain] 1 tree(s) exploded

#6 MeSH descriptor: [Back Injuries] explode all trees

#7 MeSH descriptor: [Low Back Pain] this term only

#8 MeSH descriptor: [Sciatica] this term only

© 2017 American Medical Association. All rights reserved.

#9 low next back next pain

#10 lbp

#11 #1 or #2 or #3 or #5 or #6 or #7 or #8 or #9 or #10

#12 MeSH descriptor: [Musculoskeletal Manipulations] explode all trees

#13 MeSH descriptor: [Chiropractic] explode all trees

#14 manip*

#15 MeSH descriptor: [Osteopathic Medicine] explode all trees

#16 osteopath*

#17 chiropract*

#18 #12 or #13 or #14 or #15 or #16 or #17

#19 #11 and #18

=====
DATABASE SEARCHED & TIME PERIOD COVERED:
MEDLINE ON OVID – 1/1/2011-2/06/2017

SEARCH STRATEGY:

1 Clinical Trial.pt.

2 randomized.ab,ti.

3 placebo.ab,ti.

4 dt.fs.

5 randomly.ab,ti.

6 trial.ab,ti.

7 groups.ab,ti.

8
1 or 2 or 3 or 4 or 5 or 6 or 7

9 Animals/

10 Humans/

11
9 not (9 and 10) Including Related Terms

12

© 2017 American Medical Association. All rights reserved.

8 not 11

13 dorsalgia.ti,ab.

14 exp Back Pain/

15 backache.ti,ab.

16 (lumbar adj pain).ti,ab.

17 coccyx.ti,ab.

18 coccydynia.ti,ab.

19 sciatica.ti,ab.

20 sciatica/

21 spondylosis.ti,ab.

22 lumbago.ti,ab.

23 exp low back pain/

24

13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23

25 exp Manipulation, Chiropractic/

26 exp Manipulation, Orthopedic/

27 exp Manipulation, Osteopathic/

28 exp Manipulation, Spinal/

29 exp Musculoskeletal Manipulations/

30 exp Chiropractic/

31 manipulation.mp.

32 manipulate.mp.

33 exp Orthopedics/

34 exp Osteopathic Medicine/

35

25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34

36

12 and 24 and 35

37

36 and 2011:2015.(sa_year).

=====

DATABASE SEARCHED & TIME PERIOD COVERED:
EMBASE – 1/1/2011-2/06/2017

SEARCH STRATEGY:

#2 'clinical article'/exp OR 'clinical study'/exp OR 'clinical trial'/de OR 'controlled study'/de OR 'randomized controlled trial'/de OR 'major clinical study'/de OR 'double blind procedure'/de OR 'multicenter study'/de OR 'single blind procedure'/de OR 'phase 3 clinical trial'/de OR 'phase 4 clinical trial'/de OR 'crossover procedure'/de OR 'placebo'/de

#6 allocat*

#7 assign*

#8 blind*

#12 clinical NEAR/25 (study OR trial*)

#13 compar*

#14 control*

#17 'cross over'

#18 'cross-over'

#19 'crossover'

#20 factorial

#21 'follow up'

#22 follow* NEAR/3 up

#23 'follow up'

#24 placebo*

#25 prospectiv*

#26 random*

#27 (singl* OR doubl* OR trebl* OR tripl*) NEAR/25 (blind* OR mask*)

#28 trial

#29 versus OR vs

#30

#6 OR #7 OR #8 OR #12 OR #13 OR #14 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29

#31

#2 OR #30

© 2017 American Medical Association. All rights reserved.

#34 dorsalgia

#35 'back pain'

#36 lumbar NEAR/2 pain

#37 coccyx

#38 coccydynia

#39 sciatica

#40 spondylosis

#41 lumbago

#42 'backache'/exp OR 'ischialgia'/exp OR 'low back pain'/exp

#43

#34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40 OR #41 OR #42

#44 'chiropractic'/exp OR 'orthopedic manipulation'/exp OR 'manipulative medicine'/exp OR 'osteopathic medicine'/exp OR 'orthopedics'/exp

#45 manipulation

#46 manipulate

#47 osteopathy

#48

#44 OR #45 OR #46 OR #47

#49

#31 AND #43 AND #48

#50

#31 AND #43 AND #48 AND [humans]/lim

#51

#31 AND #43 AND #48 AND [humans]/lim AND [2011-2015]/py

=====
DATABASE SEARCHED & TIME PERIOD COVERED:
CINAHL – 1/1/2011-2/06/2017

SEARCH STRATEGY:

Search modes - Find all search terms (For all search statements)

S1 randomized controlled trials

S2 randomized controlled trials

S3 PT clinical trial

© 2017 American Medical Association. All rights reserved.

S4 (MH "Clinical Trials+")

S5 clin* n25 trial*

S6 (singl* or doubl* or trebl* or tripl*) n25 (blind* or mask*)

S7 (MH "Placebos")

S8 (MH "Study Design+")

S9 (MH "Comparative Studies")

S10 (MH "Evaluation Research+")

S11 (MH "Prospective Studies+")

S12 "follow up studies" OR "follow-up studies" OR "followup studies" OR "follow-up study" OR "follow up study" OR "followup study"

S13 control* or prospectiv* or volunteer*

S14 placebo* OR random* OR (latin n2 square*)

S15

S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14

S18 TI dorsalgia OR AB dorsalgia

S19 (MH "Back Pain+")

S20 TI backache OR AB backache

S21 TI lumbar n2 pain OR AB lumbar n2 pain

S22 TI coccyx pain OR AB lumbar n2 pain

S23 TI coccyx OR AB coccyx

S24 TI coccydynia OR AB coccydynia

S25 TI sciatica OR AB sciatica

S26 (MH "Sciatica")

S27 TI spondylosis OR AB spondylosis

S28 TI lumbago cronico OR AB spondylosis

S29 TI lumbago OR AB lumbago

S30 (MH "Low Back Pain")

S31

S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27 OR S28 OR
S29
OR S30

S32 (MH "Chiropractic+")

S33 (MH "Manipulation, Chiropractic")

S34 (MH "Manipulation, Orthopedic")

S35 (MH "Manipulation, Osteopathic")

S36 (MH "Manual Therapy+")

S37 (MH "Orthopedics")

S38 (MH "Osteopathy+")

S39 manipulation

S40 manipulate

S41

S32 OR S33 OR S34 OR S35 OR S36 OR S37 OR S38 OR S39 OR S40

S42

s15 AND S31 AND S41

S43

S15 AND S31 AND S41

DATABASE SEARCHED & TIME PERIOD COVERED:

PubMed – 1/1/2015-2/06/2017

SEARCH STRATEGY:

Manipulation, Chiropractic[mh] OR Manipulation, Orthopedic[mh] OR Manipulation,
Osteopathic[mh] OR Manipulation, Spinal[mh] OR Musculoskeletal Manipulations[mh] OR
Chiropractic[mh] OR Orthopedics OR Osteopathic Medicine
AND

"Low Back Pain"[Mesh] OR low back pain*[tiab] OR "Back"[Mesh] OR dorsalgia[tiab] OR Back
Pain[mh] OR backache[tiab] OR "lumbar pain"[tiab] OR coccyx[tiab] OR coccydynia[tiab] OR
sciatica[tiab] OR sciatica[mh] OR spondylosis[tiab] OR lumbago[tiab]

AND

Randomized Controlled Trial" [Publication Type] OR "Randomized Controlled Trials as
Topic"[Mesh] OR random*[tiab] OR rct* OR systematic[tiab] OR systematic[sb] OR Clinical
Trial[pt] OR randomized[tiab] OR placebo[tiab] OR randomly[tiab] OR trial[tiab] OR groups[tiab]

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Bergquist-Ullman, et al., 1977 ¹	Industry	87% male	34 years	14% of patients had a straight leg raise test positive at less than 60 degrees	Pain index	43	back school (instruction and exercise)	N=44	10 day median: 20 3 week median: 19 6 week median: 22
						42	non-thrust manipulation	N=50	10 day median: 22 3 week median: 18 6 week median: 21
						42	diathermy according to Cyriax, Kaltenborn, Lewit, and Janda	N=56	10 day median: 28 3 week median: 25 6 week median: 17
Blomberg, et al., 1994 ^{2,6}	Primary care	52% male	37 years	10% with "true radicular pain"	Disability Rating Score (function)	No baseline data	usual medical care	N=48	3 days mean: 4.6 1 week mean: 3.9 2 week mean: 3.2 3 week mean: 3
							mix of thrust and non-thrust manipulation, some patients also got steroid injections of the parasacroccygeal structures as described by Cyriax	N=53	3 days mean: 3.5 1 week mean: 2.6 2 week mean: 1.8 3 week mean: 1.4
					Pain score		usual medical care	N=48	3 days mean: 4.8 1 week mean: 4.2 2 week mean: 3.4 3 week mean: 3.4
							mix of thrust and non-thrust manipulation, some patients also got steroid injections of the parasacroccygeal structures as described by Cyriax	N=53	3 days mean: 3.8 1 week mean: 3.1 2 week mean: 2 3 week mean: 1.7

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Cherkin, et al., 1998 ⁷	Primary care patients from health maintenance organization	52% male	41 years	Sciatica excluded	Roland Morris Disability questionnaire (function)	12.1 (CI: 11.2-13.1)	thrust manipulation	N=122	4 week mean: 3.7 (2.9 SD)
						11.7 (CI: 10.4-13.0)	physical therapy according to McKenzie	N=136	4 week mean: 4.1 (3.3 SD)
						11.7 (CI: 10.4-13.0)	educational booklet	N=66	4 week mean: 4.9 (3.8 SD)
					Bothersomeness of symptoms (pain)	5.5 (CI: 5.1-5.8)	thrust manipulation	N=122	4 week mean: 1.9 (1.5 SD)
						6 (CI: 5.6-6.5)	physical therapy according to McKenzie	N=136	4 week mean: 2.3 (1.9 SD)
						5.3 (CI: 4.9-5.7)	educational booklet	N=66	4 week mean: 3.1 (2.4 SD)
Childs, et al., 2004 ⁸	8 physical therapy clinics in the United States	58% male	34 years	24% had "symptoms distal to knee"	Oswestry disability questionnaire (function)	41.4 (10.1 SD)	thrust manipulation	N=70	1 week mean: 14.6 4 week mean: 8.4
						40.9 (10.8 SD)	low stress aerobic exercise and lumbar spine strengthening program according to Agency for Health Care Policy and Research guidelines	N=61	1 week mean: 35 4 week mean: 23
Cramer, et al., 1993 ⁹	Clinical chiropractic college	57% male	Not reported	Patients with "compressive neuropathy" we excluded	Visual Analogue Scale (pain)	71.8 (14.8 SD)	non-thrust manipulation and electrical stimulation and cold pack	N=17	10 day mean: 38.6 (25.2 SD)
						72 (19.2 SD)	detuned ultrasound and cold pack	N=18	10 day mean: 42 (28.8 SD)
					Oswestry disability questionnaire (function)	17.6 (11.9 SD)	non-thrust manipulation and electrical stimulation and cold pack	N=17	10 day mean: 7.3 (6.8 SD)
						14.9 (5.0 SD)	detuned ultrasound and cold pack	N=18	10 day mean: 8.0 (7.6 SD)

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Cruser, et al., 2012 ¹⁰	United States military facility	55% male	27 years	Not reported	Visual Analogue Scale (pain)	5.2 (2.1 SD)	mix of thrust and non-thrust manipulation, soft tissue stretching, myofascial release, counterstrain muscle energy, sacroiliac articulation	N=30	4 week mean: 2.0 (1.5 SD)
						5.5 (2.2 SD)	usual medical care	N=30	4 week mean: 3.7 (2.4 SD)
					Roland Morris Disability questionnaire (function)	12.4 (5.3 SD)	mix of thrust and non-thrust manipulation, soft tissue stretching, myofascial release, counterstrain muscle energy, sacroiliac articulation	N=30	4 week mean: 4.4 (5.9 SD)
						12.5 (6.0 SD)	usual medical care	N=30	4 week mean: 7.31 (6.3 SD)
Delitto, et al., 1993 ¹¹	Physiotherapy department	58% male	33 years	21% had "leg symptoms"	Oswestry disability questionnaire (function)	33 (5 SD)	thrust manipulation and extension exercises according to McKenzie and hand-heel rock exercise	N=14	3 day mean: 20 (5 SD) 5 day mean: 10 (5 SD)
						41 (5 SD)	flexion exercises according to Williams	N=10	3 day mean: 36 (5 SD) 5 day mean: 32 (4 SD)
Erhard, et al., 1994 ¹²	Physiotherapy department	62% male	44 years	8% had "leg symptoms"	Oswestry disability questionnaire (function)	45 (12 SD)	thrust manipulation and extension exercises according to McKenzie	N=12	3 day mean: 20 (8 SD) 5 day mean: 8 (8 SD)
						40 (12 SD)	extension exercises according to McKenzie	N=12	3 day mean: 35 (8 SD) 5 day mean: 25 (14 SD)
Farrell, et al., 1982 ¹³	Setting unclear	62% male	42 years	Not reported	Subjective pain rating	4.95	non-thrust manipulation according to Stoddart and Maitland	N=24	3 week mean: 0.3
						5.3	physical therapy and diathermy, isometric abdominal exercises and ergonomic instructions	N=24	3 week mean: 0.3

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Fritz, et al., 2015 ¹⁴	Primary care	48% male	37 years	Patients with presence of pain or numbness distal to the knee were excluded	Numeric pain rating of low back pain severity	no baseline data	thrust manipulation and exercises	N=108	4 week mean: 1.7 (1.9 SD)
						no baseline data	standard medical care and self-help booklet	N=112	4 week mean: 2.1 (1.9 SD)
					Oswestry disability questionnaire (function)	no baseline data	thrust manipulation and exercises	N=108	4 week mean: 11.1 (12.5 SD)
						no baseline data	standard medical care and self-help booklet	N=112	4 week mean: 14.5 (13.2 SD)
Glover, et al., 1974 ¹⁵	Work medical center	89% male	39 years	Not reported	Percent pain relief	no baseline data	diathermy	N=41	3 day mean: 56 1 week mean: 80
						no baseline data	non-thrust manipulation	N=43	3 day mean: 50 1 week mean: 75
Godfrey, et al., 1984 ¹⁶	Patients referred from primary care	Not reported	42 years	Not reported	General symptomatology (number of patients with marked improvement) (pain)	no baseline data	thrust manipulation according to Maigne		2-3 week: 14/39 (35.9%)
						no baseline data	light effleurage and minimal electrostimulation		2-3 week: 7/33 (21.2%)
					Activities of Daily Living (number of patients with moderate improvement) (function)	no baseline data	thrust manipulation according to Maigne		2-3 week: 7/24 (29.2%)
						no baseline data	light effleurage and minimal electrostimulation		2-3 week: 5/17 (29.4%)

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Goertz, et al., 2013 ¹⁷	United States army medical center	86% male	26 years	43% had "radicular signs"	Numerical pain rating scale	5.8 (2.1 SD)	standard medical care and brief massage, ice or heat, McKenzie exercises, stretching exercises	N=46	2 week mean: 6.1 4 week mean: 5.2
						5.8 (1.5 SD)	thrust manipulation	N=45	2 week mean: 3.9 4 week mean: 3.9
					Roland Morris Disability questionnaire (function)	12.7 (5.1 SD)	standard medical care and brief massage, ice or heat, McKenzie exercises, strengthening exercises	N=46	2 week mean: 12.9 4 week mean: 12
						11 (4.2 SD)	thrust manipulation	N=45	2 week mean: 8.9 4 week mean: 8
Grunnesjo, et al., 2004 ¹⁸⁻²⁰	Nine primary health care and one outpatient orthopedic hospital department	56% male	41 years	8% had "verified herniations"	Pain last 24 hours	52.2 (CI: 46.7-57.8)	stay active	N=71	5 week mean: 29.7 (25.8 SD)
						54.7 (CI: 49.8-59.6)	mix of thrust and non-thrust manipulation and stay active and in some patients a steroid injection in the parasacroccygeal region	N=89	5 week mean: 20.8 (23.3 SD)
					All disability rating variables	52 (CI: 47.4-56.6)	stay active	N=71	5 week mean: 31.9 (21.9 SD)
						57.8 (CI: 53.7-61.8)	mix of thrust and non-thrust manipulation and stay active and in some patients a steroid injection in the parasacroccygeal region	N=89	5 week mean: 25.8 (22.1 SD)
Hadler, et al., 1987 ²¹	Primary care	57% male	Not reported	Not reported	Roland Morris Disability questionnaire (function)	no baseline data	mobilization	N=28	9 day mean: 4.5 12 day mean: 3.7
						no baseline data	thrust manipulation	N=26	9 day mean: 3.7 12 day mean: 3.4

© 2017 American Medical Association. All rights reserved.

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up					
Hallegraeff, et al., 2009 ²²	Three physical therapy and manual therapy centers	55% male	39 years	Patients with symptoms distal to the knee were excluded	Oswestry disability questionnaire (function)	0.24 (0.18 SD)	thrust manipulation	N=31	2.5 week mean: 0.14 (0.17 SD)					
						0.26 (0.12 SD)	physical therapy	N=33	2.5 week mean: 0.14 (0.12 SD)					
					Visual Analogue Scale (pain)	42.7 (18.4 SD)	thrust manipulation	N=31	2.5 week mean: 19 (16.9 SD)					
						54 (17.5 SD)	physical therapy	N=33	2.5 week mean: 24.8 (20.1 SD)					
Hancock, et al., 2007 ²³	Patients referred from primary care	56% male	41 years	Patients with "nerve root compromise" were excluded	Numerical pain rating scale negative effect size favors manipulation	no baseline data	non-thrust manipulation	N=59	1 week effect size: 0.2 (CI: -0.3-0.7) 2 week effect size: -0.4 (CI: -1.0, 0.1) 4 week effect size: -0.2 (CI: -0.7, 0.3)					
						no baseline data	detuned pulsed ultrasound (sham)	N=60						
					Roland Morris Disability questionnaire (function) negative effect size favors manipulation	no baseline data	non-thrust manipulation	N=59	1 week effect size: -0.7 (CI: -2.1, 0.6) 2 week effect size: -1.4 (CI: -2.7, -0.1) 4 week effect size: -1 (CI: -2.1, 0.1)					
						no baseline data	detuned pulsed ultrasound (sham)	N=60						
					Heymann, et al., 2013 ²⁴	5 orthopedic or general practices	60% male	37 years	Not reported	Roland Morris Disability questionnaire (function)	13.5 (5.6 SD)	thrust manipulation	N=38	1 week mean: 5.8
											14.4 (4.8 SD)	analgesic (diclofenac)	N=37	1 week mean: 9.7
15 (3.8 SD)	sham	N=25	no data provided											
Visual Analogue Scale (pain)	no baseline data	thrust manipulation	N=38	1 week mean: 10										
	no baseline data	analgesic	N=37	1 week mean: 30										
	no baseline data	sham	N=25	1 week mean: no data provided										

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Hoiriis, et al., 2004 ²⁵	Patients recruited via advertisement	57% male	42 years	Patients with "known or suspected disk herniation" were excluded	Visual Analogue Scale (pain)	4.52 (1.82 SD)	thrust manipulation	N=34	2 week mean: 2.4 (2.2 SD) 4 week mean: 1.7 (1.9 SD)
						3.9 (2.0 SD)	muscle relaxants (cyclobenzaprine or carisoprodol or methocarbamol)	N=36	2 week mean: 2.7 (2.2 SD) 4 week mean: 2.2 (2.2 SD)
						3.8 (1.6 SD)	sham	N=40	2 week mean: 3.2 (2.4 SD) 4 week mean: 2.2 (2.0 SD)
					Oswestry disability questionnaire (function)	24.8 (11.5 SD)	thrust manipulation	N=46	2 week mean: 17.0 (13.8 SD) 4 week mean: 11.9 (11.9 SD)
						22.8 (12.9 SD)	muscle relaxants (cyclobenzaprine or carisoprodol or methocarbamol)	N=47	2 week mean: 17.0 (12.2 SD) 4 week mean: 16.0 (16.1 SD)
						24.8 (11.7 SD)	sham	N=48	2 week mean: 19.3 (13.7 SD) 4 week mean: 16.3 (12.6 SD)
Juni, et al., 2009 ²⁶	Patients referred from emergency department or a general practice	64% male	35 years	Patients with "signs of nerve root irritation or compression" were excluded	Roland Morris Disability questionnaire (function)	12.8 (5.1 SD)	Mix of thrust and non-thrust manipulation	N=52 12.8 (5.1 SD)	2 week mean: 5.8 (5.7 SD)
						14.3 (4.9 SD)	analgesic (paracetamol, diclofenac, or dihydrocodeine)	N=52	2 week mean: 5.2 (7.0 SD)
					Pain intensity, BS-11 score positive favors manipulation	6.3 (2.2 SD)	mix of thrust and non-thrust manipulation	N=52	Difference of 0.5 (2.6 SD)
						6.8 (2.2 SD)	Analgesic (paracetamol, diclofenac, or dihydrocodeine)	N=52	

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
MacDonald, et al., 1990 ²⁷	General practice	41% male	Not reported	Patients with "neurologic deficits" were excluded	Improvement in the disability index	6.4 (3 SD)	thrust manipulation and advice on posture, exercises and avoidance of occupational stress	N=36	2 week mean: 4.1 (3.5 SD)
						6.1 (2.5 SD)	advice on posture, exercise, and avoidance of occupational stress	N=30	2 week mean: 4.4 (3.5 SD)
Morton, 1999 ²⁸	Patients referred from primary care	34% male	44 years	Patients with "abnormalities on neurologic exam" were excluded	Roland Morris Disability questionnaire (function)	10.6 (5.2 SD)	thrust manipulation	N=15	1 week mean: 6.9 (4.1 SD) 2 week mean: 6.0 (2.3 SD) 3 week mean: 3.7 (3.7 SD) 4 week mean: 1.9 (2.5 SD)
						10.1 (6.4 SD)	spinal stabilizing exercises	N=14	1 week mean: 9.1 (5.9 SD) 2 week mean: 7.9 (6.3 SD) 3 week mean: 7 (6.1 SD) 4 week mean: 6 (5.2 SD)
					Visual Analogue Scale (pain)	49.7 (23.6 SD)	thrust manipulation	N=15	1 week mean: 27.6 (15.2 SD) 2 week mean: 17.4 (13.9 SD) 3 week mean: 7.5 (6.4 SD) 4 week mean: 2.4 (3 SD)
						46.6 (25.1 SD)	spinal stabilizing exercises	N=14	1 week mean: 46.4 (23.3 SD) 2 week mean: 36.6 (24.6 SD) 3 week mean: 34.5 (23 SD) 4 week mean: 25.4 (17.3 SD)

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up	
Postacchini, et al., 1988 ²⁹	Hospital outpatient department	51% male	38 years	Not reported	Improvement in low back pain from pre-treatment	no baseline data	thrust manipulation	N=53	3 week mean: 8.5	
						no baseline data	back school	N=17	3 week mean: 10.4	
						no baseline data	analgesics (diclofenac)	N=49	3 week mean: 9.4	
						no baseline data	physiotherapy of light massage, analgesic currents, and diathermy	N=47	3 week mean: 8.1	
						no baseline data	bed rest	N=29	3 week mean: 6.6	
						no baseline data	topical gel	N=46	3 week mean: 5.8	
Rasmussen, 1979 ³⁰	Hospital department of physical medicine and rheumatology	Not reported	35 years	Patients with "signs of root pressure" were excluded	Number of patients with total restoration of all symptoms	no baseline data	non-thrust manipulation	N=12	11/12 (91.7%)	
						no baseline data	diathermy	N=12	3/12 (25%)	
Skargren, et al., 1998 ³¹	Primary care centers	38% male	41 years	Not reported	Visual Analogue Scale (pain) negative favors manipulation	56 (22 SD)	thrust manipulation	N=172	4-5 week difference: -0.16 (CI: -6.47, 6.15)	
						61 (21 SD)	physiotherapy	N=144		
						Oswestry disability questionnaire (function) negative favors manipulation	35 (17 SD)	thrust manipulation	N=172	4-5 week difference: -1.49 (CI: -5.51, 2.54)
							37 (16 SD)	physiotherapy	N=144	

eTable 1. Evidence table of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain (continued)

Author, Year	Setting	% Male	Mean Age	Presence of Leg Pain or Sciatica	Outcome	Baseline value	Treatment arms	Sample Size	Follow-up
Waterworth, et al., 1985 ³²	General practice	62% male	36 years	Not reported	Score of lower back pain	2.1	non-thrust manipulation	N=38	12 day mean: 0.42
						2.1	analgesic (diflunisal)	N=36	12 day mean: 0.44
						2	physiotherapy including local heat, ultrasound, and flexion and extension exercises	N=34	12 day mean: 0.38
					Patient has overall improvement score of excellent	no baseline data	non-thrust manipulation	N=38	23/38 (60.5%)
						no baseline data	analgesic (diflunisal)	N=36	15/36 (41.7%)
						no baseline data	physiotherapy including local heat, ultrasound, and flexion and extension exercises	N=34	13/34 (38.2%)

eTable 2. Quality scores of 26 randomized clinical trials of spinal manipulative therapy for acute low back pain

Article	Randomization	Concealment	Blinding, provider	Blinding, patient	Blinding, outcome	Dropouts	Timing	Intention to Treat	Baseline Differences	Co-interventions	Compliance	Total
Bergquist-Ullman, M., 1977 ¹	+	?	-	-	-	-	+	-	?	?	-	2
Blomberg, S., 1994 ²	+	+	-	-	-	+	-	+	-	+	+	6
Cherkin, D. C., 1998 ⁷	+	+	-	-	-	+	+	+	+	-	?	6
Childs, J. D., 2004 ⁸	+	+	-	-	-	+	+	+	+	?	+	7
Cramer, G. D., 1993 ⁹	?	?	-	-	-	+	+	+	?	?	?	3
Cruser, A., 2012 ¹⁰	?	+	-	-	-	+	+	+	+	+	+	7
Delitto, A., 1993 ¹¹	+	-	-	-	?	?	+	+	+	?	?	4
Erhard, R. E., 1994 ¹²	+	?	-	-	?	-	+	-	+	?	?	3
Farrell, J. P., 1982 ¹³	?	?	-	-	-	+	+	?	+	?	?	3
Fritz, J. M., 2015 ¹⁴	+	+	-	-	-	+	+	+	+	-	+	7
Glover, J. R., 1974 ¹⁵	+	?	-	-	-	?	+	?	?	?	?	2
Godfrey, C. M., 1984 ¹⁶	+	?	-	+	+	-	+	-	?	+	+	6
Goertz, C. M., 2013 ¹⁷	+	+	-	-	-	-	+	+	+	+	+	7
Grunnesjö, M. I., 2004 ¹⁸⁻²⁰	+	+	-	-	-	+	+	+	+	?	+	7
Hadler, N. M., 1987 ²¹	?	?	-	+	-	+	+	-	-	?	?	3
Hallegraeff, J. M., 2009 ²²	+	+	-	-	-	+	+	+	-	?	+	6
Hancock, M. J., 2007 ²³	+	+	-	+	-	+	+	+	+	+	+	9
Heymann, W. J., 2013 ²⁴	?	+	-	+	+	-	+	+	+	?	?	6
Hoiriis, K. T., 2004 ²⁵	+	?	-	-	-	-	+	+	?	?	?	3
Juni, P., 2009 ²⁶	+	+	-	-	-	+	+	+	?	+	+	7
MacDonald, R. S., 1990 ²⁷	?	?	-	-	-	+	+	-	+	?	+	4
Morton, J. E., 1999 ²⁸	+	-	-	-	-	?	+	+	?	?	?	3
Postacchini, F., 1988 ²⁹	?	?	-	-	-	+	+	-	?	-	?	2
Rasmussen, G., 1979 ³⁰	?	?	-	-	-	+	+	-	?	?	?	2
Skargren, E. I., 1998 ³¹	?	?	-	-	-	-	+	-	+	-	?	2
Waterworth, R. F., 1985 ³²	+	?	-	-	-	+	+	-	?	?	?	3

* +=yes, -=no, ?=unsure/don't know; full criteria specified in Cochrane Back Group Risk of Bias Tool.³³

eReferences

1. Bergquist-Ullman M, Larsson U. Acute low back pain in industry. A controlled prospective study with special reference to therapy and confounding factors. *Acta Orthop Scand*. 1977(170):1-117.
2. Blomberg S, Svardsudd K, Tibblin G. A randomized study of manual therapy with steroid injections in low-back pain. Telephone interview follow-up of pain, disability, recovery and drug consumption. *Eur Spine J*. 1994;3(5):246-254.
3. Blomberg S, Hallin G, Grann K, Berg E, Sennerby U. Manual therapy with steroid injections--a new approach to treatment of low back pain. A controlled multicenter trial with an evaluation by orthopedic surgeons. *Spine (Phila Pa 1976)*. 1994;19(5):569-577.
4. Blomberg S, Svardsudd K, Mildenerger F. A controlled, multicentre trial of manual therapy in low-back pain. Initial status, sick-leave and pain score during follow-up. *Scand J Prim Health Care*. 1992;10(3):170-178.
5. Blomberg S, Tibblin G. A controlled, multicentre trial of manual therapy with steroid injections in low-back pain: functional variables, side effects and complications during four months follow-up. *Clinical Rehabilitation*. 1993;7(1):49-62.
6. Blomberg S, Svardsudd K, Tibblin G. Manual therapy with steroid injections in low-back pain. Improvement of quality of life in a controlled trial with four months' follow-up. *Scand J Prim Health Care*. 1993;11(2):83-90.
7. Cherkin DC, Deyo RA, Battie M, Street J, Barlow W. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med*. 1998;339(15):1021-1029.
8. Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med*. 2004;141(12):920-928.
9. Cramer GD, Humphreys CR, Hondras MA, McGregor M, Triano JJ. The Hmax/Mmax ratio as an outcome measure for acute low back pain. *J Manipulative Physiol Ther*. 1993;16(1):7-13.
10. Crusier A, Maurer D, Hensel K, Brown SK, White K, Stoll ST. A randomized, controlled trial of osteopathic manipulative treatment for acute low back pain in active duty military personnel. *Journal of manual & manipulative therapy*. 2012;20(1):5-15.
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/708/CN-00897708/frame.html>.
11. Delitto A, Cibulka MT, Erhard RE, Bowling RW, Tenhula JA. Evidence for use of an extension-mobilization category in acute low back syndrome: a prescriptive validation pilot study. *Phys Ther*. 1993;73(4):216-222; discussion 223-218.
12. Erhard RE, Delitto A, Cibulka MT. Relative effectiveness of an extension program and a combined program of manipulation and flexion and extension exercises in patients with acute low back syndrome. *Phys Ther*. 1994;74(12):1093-1100.
13. Farrell JP, Twomey LT. Acute low back pain. Comparison of two conservative treatment approaches. *Med J Aust*. 1982;1(4):160-164.
14. Fritz JM, Magel JS, McFadden M, et al. Early physical therapy vs usual care in patients with recent-onset low back pain: A randomized clinical trial. *JAMA - Journal of the American Medical Association*. 2015;314(14):1459-1467.
15. Glover JR, Morris JG, Khosla T. Back pain: a randomized clinical trial of rotational manipulation of the trunk. *Br J Ind Med*. 1974;31(1):59-64.
16. Godfrey CM, Morgan PP, Schatzker J. A randomized trial of manipulation for low-back pain in a medical setting. *Spine (Phila Pa 1976)*. 1984;9(3):301-304.
17. Goertz CM, Long CR, Hondras MA, et al. Adding chiropractic manipulative therapy to standard medical care for patients with acute low back pain: results of a pragmatic randomized comparative effectiveness study. *Spine*. 2013;38(8):627-634.
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/298/CN-00967298/frame.html>.
18. Grunnesjo MI, Bogefeldt JP, Svardsudd KF, Blomberg SI. A randomized controlled clinical trial of stay-active care versus manual therapy in addition to stay-active care: functional variables and pain. *J Manipulative Physiol Ther*. 2004;27(7):431-441.
19. Bogefeldt J, Grunnesjo MI, Svardsudd K, Blomberg S. Sick leave reductions from a comprehensive manual therapy programme for low back pain: the Gotland Low Back Pain Study. *Clin Rehabil*. 2008;22(6):529-541.

20. Grunnesjö MI, Bogefeldt JP, Blomberg SI, Strender LE, Svärdsudd KF. A randomized controlled trial of the effects of muscle stretching, manual therapy and steroid injections in addition to 'stay active' care on health-related quality of life in acute or subacute low back pain. *Clinical rehabilitation*. 2011;25(11):999-1010.
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/578/CN-00865578/frame.html>.
21. Hadler NM, Curtis P, Gillings DB, Stinnett S. A benefit of spinal manipulation as adjunctive therapy for acute low-back pain: a stratified controlled trial. *Spine (Phila Pa 1976)*. 1987;12(7):702-706.
22. Hallegraeff JM, de Greef M, Winters JC, Lucas C. Manipulative therapy and clinical prediction criteria in treatment of acute nonspecific low back pain. *Percept Mot Skills*. 2009;108(1):196-208.
23. Hancock MJ, Maher CG, Latimer J, et al. Assessment of diclofenac or spinal manipulative therapy, or both, in addition to recommended first-line treatment for acute low back pain: a randomised controlled trial. *Lancet*. 2007;370(9599):1638-1643.
24. Heymann WJ, Schloemer P, Timm J, Muehlbauer B. Spinal high-velocity low amplitude manipulation in acute nonspecific low back pain: a double-blinded randomized controlled trial in comparison with diclofenac and placebo. *Spine*. 2013;38(7):540-548.
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/716/CN-00878716/frame.html>.
25. Hoiriis KT, Pflieger B, McDuffie FC, et al. A randomized clinical trial comparing chiropractic adjustments to muscle relaxants for subacute low back pain. *J Manipulative Physiol Ther*. 2004;27(6):388-398.
26. Juni P, Battaglia M, Nuesch E, et al. A randomised controlled trial of spinal manipulative therapy in acute low back pain. *Ann Rheum Dis*. 2009;68(9):1420-1427.
27. MacDonald RS, Bell CM. An open controlled assessment of osteopathic manipulation in nonspecific low-back pain. *Spine (Phila Pa 1976)*. 1990;15(5):364-370.
28. Morton JE. Manipulation in the treatment of acute low back pain. *Journal of Manual & Manipulative Therapy*. 1999;7(4):182-189.
29. Postacchini F, Facchini M, Palieri P. Efficacy of various forms of conservative treatment in low back pain: a comparative study. *Neuro-orthopedics*. 1988;6(1):28-35.
30. Rasmussen G. Manipulation in treatment of low back pain: a randomized clinical trial. *Man Med*. 1979;1:8-10.
31. Skargren EI, Carlsson PG, Oberg BE. One-year follow-up comparison of the cost and effectiveness of chiropractic and physiotherapy as primary management for back pain. Subgroup analysis, recurrence, and additional health care utilization. *Spine (Phila Pa 1976)*. 1998;23(17):1875-1883; discussion 1884.
32. Waterworth RF, Hunter IA. An open study of diflunisal, conservative and manipulative therapy in the management of acute mechanical low back pain. *N Z Med J*. 1985;98(779):372-375.
33. Furlan AD, Pennick V, Bombardier C, van Tulder M. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine*. 2009;34(18):1929-1941.