

Supplementary Online Content

Soneji S, Barrington-Trimis JL, Wills TA, et al. Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: a systematic review and meta-analysis. *JAMA Pediatr*. Published online June 26, 2017. doi:10.1001/jamapediatrics.2017.1488

eTable 1. Summary of Search Results

eTable 2. Pubmed Run (Conducted on July 28, 2016)

eTable 3. Cochrane Library (Wiley) Run (Conducted on July 28,2016)

eTable 4. Keyword Search Terms for Society for Nicotine & Tobacco and Society for Behavioral Medicine Annual Meetings and NIH Tobacco Regulatory Science Conference

eTable 5. Psychosocial and Behavioral Characteristics Included in Studies

eTable 6. Quality of Studies Assessment (Newcastle – Ottawa Scale)

eTable 7. Risk of Bias (ROBINS-I)

eTable 8. Pooled Adj. Odds Ratio Varying Prob. of Publishing Study with Largest Standard Error

eTable 9. Pooled Adj. Odds Ratio: Copas Selection Model and Random Effects Model

eTable 10. Source of Data from Each Study in Systematic Review and Meta-Analysis

eFigure 1. Embase Run (Conducted on July 28, 2016)

eFigure 2. Web of Science Run (Conducted on July 28, 2016)

eFigure 3. Copas Selection Modelling

eAppendix 1. Selection Bias: Copas Selection Modelling

eTable 1. Summary of Search Results

Database	Platform	Years covered	Dates Conducted	# Results
Medline	Pubmed	1946-current	July 28, 2016 December 15, 2016 February 13, 2017	2471
Embase	Embase.com	1974-current	July 28, 2016 December 13, 2016 February 7, 2017	3156
Web of Science	WOS	1900-current	July 28, 2016 December 15, 2016 February 13, 2017	2055
Cochrane Library	Wiley	Issue #, date DSR - Iss7, July 2016 Dare-Iss2, April 2015 Trials-Iss6, June 2016 methods-Iss3, July 2012 tech-Iss2, April 2016	July 28, 2016 December 15, 2016 February 17, 2017	240
Society for Research on Nicotine & Tobacco	—	2016	September 1, 2016	803
Society for Behavioral Medicine	—	2016	September 1, 2016	34
NIH Tobacco Regulatory Science Conference	—	2016	September 1, 2016	167
Total				8926
Total with Duplicates Removed				6959

eTable 2. Pubmed Run (Conducted on July 28, 2016)

Search	Query	Items Found
#3	Search (#1 AND #2)	2,044
#2	Search Tobacco Use[mesh] OR Tobacco[mesh] OR Tobacco use disorder[mesh] OR Tobacco Products[mesh] OR Cigarette*[tiab] OR Tobacco[tiab] OR Smoking[tiab] OR smoker*[tiab] OR cigar*[tiab]	292,817
#1	Search Electronic Cigarettes[mesh] OR (Nebulizers and Vaporizers[mesh] AND (tobacco[mesh] OR tobacco[tiab] OR nicotine[mesh] OR nicotine[tiab])) OR (Drug Delivery Systems[mesh] AND (tobacco[mesh] OR tobacco[tiab] OR nicotine[mesh] OR nicotine[tiab])) OR Electronic Cigarette*[tiab] OR E-Cig*[tiab] OR electronic nicotine delivery system*[tiab] OR vape*[tiab] OR vaping[tiab]	2,231

Table 3. Cochrane Library (Wiley) Run (Conducted on July 28,2016)

ID	Search	Hits
#1	Electronic Cigarette* or E-Cig* or electronic nicotine delivery system* or vape* or vaping or nebulize*:ti,ab,kw (Word variations have been searched)	3,344
#2	Cigarette* or Tobacco or Smoking or smoker* or cigar*:ti,ab,kw (Word variations have been searched)	20,039
#3	#1 and #2	196

eTable 4. Keyword Search Terms for Society for Nicotine & Tobacco and Society for Behavioral Medicine Annual Meetings and NIH Tobacco Regulatory Science Conference

Keywords
Electronic cigarette*
E-cig*
Electronic nicotine delivery system
Vape*
Vaping
Tobacco
Smoking
Cigarette

eTable 5. Psychosocial and Behavioral Characteristics Included in Studies

Study	Covariate	Details
Leventhal et al. (2015)	Depressive Symptoms	20-item Center for Epidemiologic Studies Depression Scale
	Impulsivity	5-item Temperament and Character Inventory impulsivity subscale
	Delinquent Behavior	Sum of frequency ratings for engaging in 11 different behaviors (e.g., stealing, lying to parents)
	Substance Use	Ever use of alcohol and 13 separate illicit and prescription substances of abuse
	Family History of Smoking	“Does anyone in your immediate family (brothers, sisters, parents, or grandparents) have a history of smoking cigarettes?”
	Peer Smoking	“In the last 30 days, how many of your 5 closest friends have smoked cigarettes?”
	Smoking Susceptibility	“Would you try smoking a cigarette if one of your best friends offered it to you?” “Do you think you would smoke in the next 6 months?” “Are you curious about smoking?” (definitely not, probably not, probably yes, definitely yes)
	Smoking Expectancies	Average of the 2 responses for “I think I might enjoy ... smoking” and (reversed) “I think I might feel bad ... from smoking.” (1 strongly disagree, 2 disagree, 3 agree, 4 strongly agree)
Primack et al. (2015)	Sensation Seeking	Composite measure of sensation-seeking tendency previously found to be related to cigarette smoking and other high-risk health behaviors based on 6 items, such as “I like to do dangerous things”
	Parental Smoking	Never (0), former (1), occasional (2), and daily (3), and scores for mothers and fathers were averaged
	Peer Smoking	# of respondents’ close friends smoked cigarettes, with response choices of none (0), few (1), more than a few (2), or most (3)
Barrington-Trimis et al. (2016)	Cigarette Use in Home	“Does anyone who lives with you now use cigarettes?”
	Peer Smoking	“How many of your □4 closest friends use [cigarettes]?” (0–4 friends)
	Peer Acceptability of Smoking	“How would your best friends act toward you if you used cigarettes?” (very unfriendly, unfriendly, friendly, or very friendly)
Primack et al. (2016)	Self-Esteem	Single-Item Self-Esteem Scale
	Sensation Seeking	4-item validated Likert-type scale that included items such as “I like to do dangerous things”
	Rebelliousness	Validated Likert-type subscale of Smith and Fogg that included items such as “I tend to go against the rules”
Unger et al. (2016)	Past Month Use of Alcohol	—
	Past Month Use of Other Tobacco Products	Hookah, cigars, little cigars, smokeless tobacco
Hornik	Sensation Seeking	4-item validated Likert-type scale that included items such as “I like to do frightening things”

et al. (2016)	Ever Cigarette Use	“Have you ever tried smoking cigarettes, even one or two puffs?”
	Cigarette Use in Home	“Does anyone who lives with you now smoke cigarettes?”
	Peer Smoking	“How many of your 4 closest friends smoke cigarettes?” (0-4 friends)
Spindle et al. (2016)	Depression	4 items from the Symptom Checklist (SCL)-90. Items are measured on 5-point Likert scales and included “feeling blue,” “worrying too much about things,” “feeling hopeless about the future,” “feeling no interest in things.”
	Anxiety	4 items from the SCL-90. Items are measured on 5-point Likert scales and included “feeling fearful,” “suddenly scared for no reason,” “nervousness or shakiness inside,” “spells of terror or panic.”
	Impulsivity	5 subscales from the UPPS-P Impulsive Behavior Scale. Subscales each consisted of 3 items measured on 4-point Likert scales. Subscales included: lack of perseverance, lack of premeditation, negative urgency, positive urgency, and sensation seeking.
	Stressful Life Events	12 items inquiring about potentially stressful life events in the past 12 months (e.g., “separation from loved one or close friend”). Stressful life events were summed to create an overall score.
	Peer Deviance	6 items addressing how many of the student’s friends (from “none” to “all”) had smoked cigarettes, drank alcohol, gotten drunk, had problems with alcohol, been in trouble with the law, and smoked marijuana. Items were summed to create an overall peer deviance score.
	Other Tobacco Use	Ever use of “smokeless tobacco,” “little cigars/cigarillos,” and “hookah.”
Miech et al. (2017)	Binge Drinking, Past 2 Weeks	“Think back over the last 2 weeks. How many times (if any) have you had 5 or more drinks in a row?”
	Marijuana Use, Past 30 Days	“On how many occasions (if any) have you used marijuana (weed, pot) or hashish (hash, hash oil) during the last 30 days?”

eTable 6. Quality of Studies Assessment (Newcastle – Ottawa Scale)

Category	Criteria	Leventhal et al. (2015)	Primack et al. (2015)	Wills et al. (2015)	Barrington-Trimis et al. (2015)	Unger et al. (2016)	Hornik & Gibson (2016)	Primack et al. (2016)	Spindle et al. (2016)	Miech et al. (2017)
Selection	Representativeness of the exposed cohort	b*	b*	b*	b*	b*	b*	b*	b*	b*
	Selection of the non-exposed cohort	a*	a*	a*	a*	a*	a*	a*	a*	a*
	Ascertainment of exposure	c	c	c	c	c	c	c	c	c
	Demonstration that outcome of interest was not present at start of study	a*	a*	a*	a*	a*	a*	a*	a*	a*
Comparability	Comparability of cohorts on the basis of the design or analysis	*	*	*	*	*	*	*	*	*
Outcome	Assessment of outcome	c	c	c	c	c	c	c	c	c
	Was follow-up long enough for outcomes to occur ¹	a*	a*	a*	a*	a*	a*	a*	a*	a*
	Adequacy of follow-up of cohorts ²	b*	c	c	c	b*	d	c	b*	c
Total # of stars (*)		6	5	5	5	6	5	5	6	5

¹6 months considered adequate follow up time

²Studies with <20% loss to follow-up received one star for adequacy of follow-up of cohorts

Note: Lev=Leventhal et al. (2015); Pri=Primack et al. (2015); Wil=Wills et al. (2016); Bar=Barrington-Trimis et al. (2016); Ung=Unger et al. (2016); Hor=Hornik et al. (2016); Spi=Spindle et al. (2016); Mie=Miech et al. (2017). Letters refer to the grade as denoted by the Newcastle – Ottawa Scale rubric (http://www.ohri.ca/programs/clinical_epidemiology/nosgen.pdf).

eTable 7. Risk of Bias (ROBINS-I)

Study	Confounding	Selection	Measurement of Intervention	Missing Data	Measurement of Outcomes	Reported Result	Overall
Lev (2015)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Pri (2015)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Wil (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Bar (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Pri (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Ung (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Hor (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Spi (2016)	Moderate	Low	Low	Low	Low	Moderate	Moderate
Mie (2017)	Moderate	Low	Low	Low	Low	Moderate	Moderate

Note: Moderate=the study is sound for a non-randomized study with regard to this domain but cannot be considered comparable to a well-performed randomized trial; Low=the study is comparable to a well-performed randomized trial with regard to this domain. Lev=Leventhal et al. (2015); Pri=Primack et al. (2015); Wil=Wills et al. (2016); Bar=Barrington-Trimis et al. (2016); Ung=Unger et al. (2016); Hor=Hornik et al. (2016); Spi=Spindle et al. (2016); Mie=Miech et al. (2017).

eTable 8. Pooled Adj. Odds Ratio Varying Prob. of Publishing Study with Largest Standard Error

Probability of publishing study with largest standard error	OR [95% CI]	P-value for hypothesis of overall treatment effect	P-value for hypothesis that no selection remains unexplained
1	3.46 (2.34-5.14)	<0.001	0.065
0.85	3.32 (2.24-4.93)	<0.001	0.090
0.73	3.16 (2.14-4.66)	<0.001	0.114
0.63	3.01 (2.05-4.40)	<0.001	0.142
0.55	2.86 (1.97-4.14)	<0.001	0.184
0.48	2.72 (1.89-3.90)	<0.001	0.246
0.42	2.59 (1.82-3.69)	<0.001	0.340

Note: Adj.=Adjusted; Prob.=Probability; OR=odds ratio; CI=confidence interval

eTable 9. Pooled Adj. Odds Ratio: Copas Selection Model and Random Effects Model

Model	OR [95% CI]	P-value for hypothesis of overall treatment effect	P-value for hypothesis that no selection remains unexplained
Copas Selection	3.16 (2.14-4.66)	< 0.0001	0.114
Random Effects	3.50 (2.38-5.16)	< 0.0001	—

Note: Adj.=Adjusted; OR=odds ratio; CI=confidence interval

eTable 10. Source of Data from Each Study in Systematic Review and Meta-Analysis

Study¹	Summary Information	Transition Probabilities	Unadjusted OR	Adjusted OR
Leventhal et al. (2015)	Age of Sample: Table 1 Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Figure	Derived from 1-6 month and 7-12 month transition probabilities in Table 3	Table 4	Table 4
Primack et al. (2015)	Age of Sample: Table 1 Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Methods	Table 2	Derived from counts in Table 2	Table 3
Wills et al. (2015)	Age of Sample: Methods Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Methods	Derived from counts in Table 2	Derived from counts in Table 2	Table 4
Barrington-Trimis et al. (2016)	Age of Sample: Results Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Figure 1	Table 2	Derived from counts in Table 2	Table 2
Unger et al. (2016)	Age of Sample: Results Study Period: Material and Methods Follow-Up Period: Material and Methods Loss to Follow-Up: Material and Methods	Figure 1	Derived from counts and transition probabilities in Results	Table 1
Hornik et al. (2016)	Age of Sample: Method Study Period: Method Follow-Up Period: Method Loss to Follow-Up: Method	Additional analysis by authors	Additional analysis by authors	Additional analysis by authors
Primack et al. (2016)	Age of Sample: Methods Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Methods	Table 2	Derived from counts in Results and transition probabilities in Table 2	Table 3
Spindle et al. (2017)	Age of Sample: Methods Study Period: Methods	Derived from counts in Table 1	Derived from counts in Table 1	Table 3

	Follow-Up Period: Methods Loss to Follow-Up: Methods			
Miech et al. (2017)	Age of Sample: Methods Study Period: Methods Follow-Up Period: Methods Loss to Follow-Up: Methods	Table 2	Derived from transition probabilities in Table 2 and counts in Appendix Table A1	Table 2

¹Each study is hyperlinked to its unique journal or conference proceeding website

Note: OR=odds ratio.

eFigure 1. Embase Run (Conducted on July 28, 2016)

The screenshot displays the Embase search interface. At the top, the search bar contains the query "#5 OR #6". Below the search bar, there are navigation options: Search, Mapping, Date, Sources, Fields, Quick limits, EBM, Pub. types, Languages, Gender, Age, and Animal. A "Search" button is visible on the left.

On the left side, there are "Results Filters" with expand/collapse options and an "Apply" button. The filters include: Sources, Drugs, Diseases, Devices, Floating Subheadings, Age, Gender, Study types, Publication types, Journal titles, Publication years, Authors, Conference Abstracts, Drug Trade Names, Drug Manufacturers, Device Trade Names, and Device Manufacturers.

The main area shows a "History" table with the following entries:


Search ID	Query	Count
#7	#5 OR #6	2,741
#6	#3 AND #4	2,037
#5	#1 AND #2	1,797
#4	'tobacco use'/exp OR 'tobacco'/exp OR 'tobacco dependence'/exp	328,298
#3	'electronic cigarette'/exp OR ('vaporizer'/exp AND ('tobacco'/exp OR tobacco.ab,ti OR 'nicotine'/exp OR nicotine.ab,ti)) OR ('drug delivery system'/exp AND ('tobacco'/exp OR tobacco.ab,ti OR 'nicotine'/exp OR nicotine.ab,ti))	3,213
#2	cigarette*.ab,ti OR 'tobacco product'*.ab,ti OR 'tobacco use'.ab,ti OR 'tobacco dependence'.ab,ti OR 'tobacco abuse'.ab,ti OR 'tobacco consumption'.ab,ti OR smoking.ab,ti OR smoker*.ab,ti OR cigar*.ab,ti OR tobacco.ab,ti	337,891
#1	'electronic cigarette'*.ab,ti OR 'e cig'*.ab,ti OR 'electronic nicotine delivery system'*.ab,ti OR vape*.ab,ti OR vaping.ab,ti	2,082

Below the history table, it indicates "2,741 results for search #7 | Show all abstracts". The "Results" section shows a list of search results, sorted by "Publication Year". The first three results are:

- Electronic-cigarette use by individuals in treatment for substance abuse: A survey of 24 treatment centers in the United States**
 Gubner N.R., Andrews K.B., Mohammad-Zadeh A., Lisha N.E., Guydish J.
 [In Process] *Addictive Behaviors* 2016 63 (45-50)
 Embase [Abstract](#) [Index Terms](#) [View Full Text](#) [PENN TEXT](#)
- Smokers' and e-cigarette users' perceptions of modified risk warnings for e-cigarettes**
 Wackowski O.A., O'Connor R.J., Strasser A.A., Hammond D., Villanti A.C., Delnevo C.D.
 [In Process] *Preventive Medicine Reports* 2016 4 (309-312)
 Embase [Abstract](#) [Index Terms](#) [View Full Text](#) [PENN TEXT](#)
- Associations between tobacco and nicotine product use and depressive symptoms among college students in Texas**
 Bandiera F.C., Loukas A., Wilkinson A.V., Perry C.L.
Addictive Behaviors 2016 63 (19-22)
 Embase [Abstract](#) [Index Terms](#) [View Full Text](#) [PENN TEXT](#)


At the bottom left, there is a "records.ris" download button. At the bottom right, there is a "Show all downloads..." button.

eFigure 2. Web of Science Run (Conducted on July 28, 2016)

Search History: **Web of Science™ Core Collection** 

Set	Results	
		Save History / Create Alert Open Saved History
# 5	2,199	#4 AND #3 <i>Indexes=SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, ESCI Timespan=All years</i>
# 4	11,862	TOPIC: (Electronic Cigarette* OR E-Cig* OR electronic nicotine delivery system* OR vape* OR vaping OR nebulize*) <i>Indexes=SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, ESCI Timespan=All years</i>
# 3	330,947	TOPIC: (Cigarette* OR Tobacco OR Smoking OR smoker* OR cigar*) <i>Indexes=SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, ESCI Timespan=All years</i>

eFigure 3. Copas Selection Modelling



eAppendix. Selection Bias: Copas Selection Modeling

Two of the seven studies on cigarette smoking initiation fell outside the 95% confidence intervals denoted by the diagonal dashed lines shown in the funnel plot (Figure A1, Panel A), which suggests possible heterogeneity and publication bias. We then assessed the sensitivity of the meta-analysis to selection mechanisms of varying strength.^{1,2} Specifically, γ_0 is approximately equal to the probit of the probability that a study with a large standard error is published and γ_1 is approximately equal to the probit of the probability that a study with precision equal to the inverse of its standard error is published. The contour plot (Figure A1, Panel B) suggests that the logarithm of the estimated adjusted pooled odds ratio from the meta-analysis may be sensitive (i.e., varies between 0.7 and 1.25) to the range of (γ_0, γ_1) values. We further explore this sensitivity in Figure A1, Panels C and D. As the probability of publishing the study with the largest standard error decreases from 100% to 40%, the estimated adjusted pooled odds ratio decreases from 3.57 ($e^{1.27}$) to 2.59 ($e^{0.95}$; Figure A1, Panel C). Notably, the confidence interval of the adjusted pooled odds ratio remains above 1 (i.e., confidence interval of log odds ratio remains above 0) across the range of probabilities of publishing the study with the largest standard error. For each of the selection probabilities shown in Figure A1, Panel C, the Copas selection model calculates a p-value for the test of any remaining selection bias. Selection mechanisms for which this p-value is not statistically significant (i.e., p-value $\geq 10\%$) correspond to more plausible estimates of the adjusted pooled odds ratio under the Copas selection model.¹ The model indicates statistically significant residual publication bias (i.e., p-value $< 10\%$) until the probability of publishing the study with the largest standard error falls just below 65%. In other words, estimated adjusted pooled odds ratios corresponding to probabilities of publishing the study with the largest standard error below 60% are the most plausible under the model. Notably, all of these estimated adjusted pooled odds ratios are statistically significant (Table E1). Overall, adjusting for selection bias, the estimated adjusted pooled odds ratio equaled 3.16 (95% CI: 2.14-4.66) compared to 3.50 (95% CI: 2.38-5.16) under the baseline random effects model (Table E2).

References

1. Carpenter JR, Schwarzer G, Rücker G, Küstler R. Empirical evaluation showed that the Copas selection model provided a useful summary in 80% of meta-analyses. *J Clin Epidemiol.* 2009;62(6):624-631.e4. doi:10.1016/j.jclinepi.2008.12.002.
2. Schwarzer G, Carpenter J, Rücker G. Empirical evaluation suggests Copas selection model preferable to trim-and-fill method for selection bias in meta-analysis. *J Clin Epidemiol.* 2010;63(3):282-288. doi:10.1016/j.jclinepi.2009.05.008.