



Original Investigation | Psychiatry

Association of Economic Status and Educational Attainment With Posttraumatic Stress Disorder

A Mendelian Randomization Study

Renato Polimanti, PhD; Andrew Ratanatharathorn, PhD; Adam X. Maihofer, MS; Karmel W. Choi, PhD; Murray B. Stein, MD, MPH; Rajendra A. Morey, MD; Mark W. Logue, PhD; Caroline M. Nievergelt, PhD; Dan J. Stein, FRCP, PhD, DPhil; Karestan C. Koenen, PhD; Joel Gelernter, MD; for the Psychiatric Genomics Consortium Posttraumatic Stress Disorder Working Group

Abstract

IMPORTANCE There is a well-established negative association of educational attainment (EA) and other traits related to cognitive ability with posttraumatic stress disorder (PTSD), but the underlying mechanisms are poorly understood.

OBJECTIVES To investigate the association of PTSD with traits related to EA.

DESIGN, SETTING, AND PARTICIPANTS Genetic correlation, polygenic risk scoring, and mendelian randomization (MR) were conducted including 23 185 individuals with PTSD and 151 309 control participants from the Psychiatric Genomics Consortium for PTSD and up to 1 131 881 individuals assessed for EA and related traits from UK Biobank, 23andMe, and the Social Science Genetic Association Consortium. Data were analyzed from July 3 through November 19, 2018.

MAIN OUTCOMES AND MEASURES Genetic correlation obtained from linkage disequilibrium score regression, phenotypic variance explained by polygenic risk scores, and association estimates from MR.

RESULTS Summary association data from multiple genome-wide association studies were available for a total of 1 180 352 participants (634 391 [53.7%] women). Posttraumatic stress disorder showed negative genetic correlations with EA ($r_g = -0.26$; $SE = 0.05$; $P = 4.60 \times 10^{-8}$). Mendelian randomization analysis, conducting considering a random-effects inverse-variance weighted method, indicated that EA has a negative association with PTSD ($\beta = -0.23$; 95% CI, -0.07 to -0.39 ; $P = .004$). Investigating potential mediators of the EA-PTSD association, propensity for trauma exposure and risk-taking behaviors were observed as risk factors for PTSD independent of EA (trauma exposure: $\beta = 0.37$; 95% CI, 0.19 to 0.52 ; $P = 2.57 \times 10^{-5}$; risk-taking: $\beta = 0.76$; 95% CI, 0.38 to 1.13 ; $P = 1.13 \times 10^{-4}$), while income may mediate the association of EA with PTSD (MR income: $\beta = -0.18$; 95% CI, -0.29 to -0.07 ; $P = .001$; MR EA: $\beta = -0.23$; 95% CI, -0.39 to -0.07 ; $P = .004$; multivariable MR income: $\beta = -0.32$; 95% CI, -0.57 to 0.07 ; $P = .02$; multivariable MR EA: $\beta = -0.04$; 95% CI, -0.29 to 0.21 ; $SE = 0.13$; $P = .79$).

CONCLUSIONS AND RELEVANCE Large-scale genomic data sets add further evidence to the negative association of EA with PTSD, also supporting the role of economic status as a mediator in the association observed.

JAMA Network Open. 2019;2(5):e193447. doi:10.1001/jamanetworkopen.2019.3447

Key Points

Question Which mechanisms underlie the negative association of posttraumatic stress disorder (PTSD) with traits related to educational attainment (EA)?

Findings In this mendelian randomization study based on large-scale genomic data sets, including data from more than 1 million individuals, EA was negatively associated with PTSD, also supporting the role of economic status as a mediator in this association.

Meaning This study suggests that economic status may mediate the association of EA with PTSD independent of the brain mechanisms associated with EA.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY License.

Introduction

Posttraumatic stress disorder (PTSD) is a psychological condition that occurs in some individuals after exposure to a major traumatic event. Prospective studies have suggested that many variables previously considered outcomes of trauma are likely to be pretrauma risk factors.¹ Among these complex associations, that of PTSD with cognitive ability and educational attainment (EA; ie, the number of years of schooling that individuals complete) is among the most puzzling.

While there is a robust epidemiologic literature on the negative association of PTSD with cognitive ability and EA,²⁻⁵ the underlying mechanisms remain unclear. Reverse causation (ie, when an outcome precedes and causes the exposure)⁶⁻⁸ is a key obstacle to disentangling the direction of the mechanisms that associate these 2 phenotypes.

Adequately powered genome-wide association studies (GWASs) are able to dissect the predisposition to complex traits. This requires extremely large sample sizes to detect the polygenic architecture of complex traits, overcoming their heterogeneity to have sufficient power to find risk loci of very small effect.⁹ By combining such small-effect loci, it is possible to build genetic instruments that can be used to investigate complex epidemiological associations, such as the underlying mechanisms connecting PTSD, traits related to EA, and the potential mediation of other pretrauma risk factors. In particular, genetic information can remove the bias of reverse causation from analysis of the association of PTSD with cognition and education. Genetic variants are allocated at conception and do not change throughout life, and they can be used to define reliable genetic instruments that can be applied in a mendelian randomization (MR) analysis.¹⁰⁻¹⁴ The basic principle in MR is that an instrumental variable based on genetic variants associated with a phenotype can be used to represent, or mirror, the disease risk associated with that phenotype without the presence of possible environmental confounders.¹⁵

Previous large-scale GWASs conducted by the Social Science Genetic Association Consortium (SSGAC)¹⁶⁻¹⁹ investigated traits related to EA, identifying a number of loci and biological pathways regulating brain mechanisms at the basis of human cognitive ability. A 2018 genome-wide analysis of multiple brain disorders and phenotypes²⁰ showed pervasive shared heritability of these traits with EA and related traits. Based on the GWAS regarding PTSD newly generated by the Psychiatric Genomics Consortium (PGC),²¹ we applied multiple statistical methods to large-scale genomic data sets to investigate the mechanisms involved in the association of PTSD with traits related to EA and potential pretrauma risk factors.

Methods

This study was conducted using summary association data generated by previous studies. Owing to the use of previously collected, deidentified, aggregated data, this study did not require institutional review board approval. Ethical approval had been obtained in all original studies.^{16,21} Summary data were available for a total of 1 180 352 participants. Multiple statistical methods were applied to these data sets to investigate the association of PTSD with EA and related traits. A schematic workflow summarizing the analyses conducted is reported in eFigure 1 in the [Supplement](#). These analyses were conducted from July 3 through November 19, 2018. The study was reported in accordance to the Strengthening the Reporting of Genetic Association Studies (STREGA) reporting guideline.²²

Cohorts Investigated

Genome-wide information regarding PTSD was derived from the freeze-2 analysis conducted by the Psychiatric Genomics Consortium Posttraumatic Stress Disorder (PGC-PTSD) Working Group.²¹ In this analysis, lifetime and/or current PTSD status was assessed using various instruments and different versions of the *Diagnostic and Statistical Manual of Mental Disorders* (Third Edition Revised, Fourth Edition, and Fifth Edition). We focused on the data generated from the analysis of individuals

of European descent (23 185 individuals with PTSD; 151 309 control participants) because the genome-wide analyses of traits associated with EA were conducted only on this ancestry group.

Genome-wide information regarding traits related to EA were derived from the GWAS meta-analysis by the SSGAC,¹⁶ which investigated EA as the primary phenotype in a total of 1 131 881 individuals (EA1M). Additional information about the definition of the EA phenotype is available in the eAppendix in the [Supplement](#). In the same SSGAC study,¹⁶ 3 additional phenotypes were investigated. Of these, 2 were analyzed exclusively among research participants of the personal genomics company 23andMe. Participants were asked to rate their mathematical ability (MA; n = 564 698; very poor, 0; poor, 1; about average, 2; good, 3; excellent, 4) and to list the most advanced math course they had successfully completed (MC; n = 430 445; prealgebra, 1; algebra, 2; geometry, 3; trigonometry, 4; precalculus, 5; calculus, 6; vector calculus, 7; >vector calculus, 8). The third phenotype investigated, cognitive performance, was assessed in 257 828 participants from the Cognitive Genomics Consortium study and the UK Biobank.¹⁶ For the data sets including participants from 23andMe, we had access only to summary association data of the top 10 000 variants. Accordingly, the data sets derived from 23andMe were not used for the reverse analysis (ie, estimating the association of PTSD with traits related to EA).

A summary of the data sets tested is reported in the **Table**. Since UK Biobank participants were included in both PGC-PTSD and SSGAC studies, some analyses were conducted on a PGC-PTSD subsample that excluded the UK Biobank cohort (PGC-PTSD freeze-1.5 data set: 12 823 individuals with PTSD; 35 648 control participants). Excluding UK Biobank, negligible overlap is present between PGC-PTSD and SSGAC cohorts (eAppendix in the [Supplement](#)).

We also used data from the UK Biobank to investigate the possible mediation of PTSD risk factors in their associations with traits related to EA. Self-reported risk-taking behaviors were assessed with the question, "Would you describe yourself as someone who takes risks?" (UK Biobank data field 2040). Income was assessed with, "Average total household income before tax" (UK Biobank data field 738). We also considered traumatic events assessed in the UK Biobank (eTable 1 in the [Supplement](#)). Genome-wide information regarding these traits was derived from GWAS summary association data generated from the UK Biobank.²³

Genetic Correlation and Definition of the Genetic Instruments

Linkage disequilibrium (LD) score regression was used to estimate the genetic correlation among the traits investigated.²⁴ Since sample overlap between the GWAS tested does not affect the results obtained from this method,²⁴ we were able to calculate pairwise genetic correlations, including those between data sets that had overlapping information with the UK Biobank participants.

Table. Traits Tested With Corresponding Information Regarding Sample Size, Data Available, and Cohorts Included

Trait	Abbreviation	Sample Size	Cohort	Summary Statistics
Posttraumatic stress disorder	PTSD freeze-1.5	12 823 with PTSD; 35 648 controls	PGC	Full
Posttraumatic stress disorder	PTSD freeze-2	23 185 with PTSD; 151 309 controls	PGC, UKB	Full
Cognitive performance	CP	257 828	SSGAC, UKB	Full
Educational attainment	EA	766 345	SSGAC, UKB	Full
Educational attainment	EA1M	1 131 881	SSGAC, UKB, 23andMe	Top 10 000
Self-reported math ability	MA	564 692	23andMe	Top 10 000
Most advanced math course completed	MC	430 439	23andMe	Top 10 000
Risk-taking behaviors	RT	348 549	UKB	Full
Income	Income	311 028	UKB	Full
Physically abused by family as a child	PAC	117 838	UKB	Full

Abbreviations: PGC, Psychiatric Genomics Consortium; PTSD, posttraumatic stress disorder; SSGAC, Social Science Genetic Association Consortium; UKB, UK Biobank.

The polygenic risk scores (PRSs) were calculated after using P value–informed clumping with an LD cutoff of $R^2 = 0.001$ within a 10 000-kilobase window, excluding the major histocompatibility complex region of the genome because of its complex LD structure and including only variants with a minor allele frequency less than 1%. The European samples from the 1000 Genomes Project were used as the LD reference panel.²⁵ The PRS analysis was conducted on the basis of the GWAS summary association data using the gtx R package incorporated in PRSice software.²⁶ For each PRS analysis, we calculated an approximate estimate of the explained variance from a multivariate regression model.²⁷ For the traits related to EA, we considered a genome-wide significance threshold ($P < 5.00 \times 10^{-8}$). For other traits (ie, PTSD, income, risk-taking behaviors, and trauma exposure), owing to the limited power to detect a large number of genome-wide significant loci, we evaluated multiple P value thresholds (PT; PT = 5.00×10^{-8} , 10^{-7} , 10^{-6} , 10^{-5} , 10^{-4} , .001, .05, 1, .3, .5, and PT < 1) to increase the variance explained by the genetic instruments. To account for the multiple PTs tested, we considered $P < .005$ as the significance threshold in the PRS analysis. Tests of statistical significance were 2-tailed. The results of the PRS analyses were used to define the genetic instruments for each pairwise comparison to be investigated further via the MR approach.

Mendelian Randomization

To assess the association among the traits tested, we used GWAS summary association data to conduct 2-sample MR analyses.²⁸ As mentioned earlier, genetic instruments were based on the results obtained in the PRS analysis. Since different MR methods have different sensitivities to different potential issues, accommodate different scenarios, and vary in their statistical efficiency,¹¹ we considered a range of MR methods. The primary analysis was conducted considering a random-effects inverse-variance weighted (IVW) method.²⁹ The secondary MR methods included MR Egger,³⁰ simple mode,³¹ weighted median,³² and weighted mode.³¹ These MR analyses were conducted using the TwoSampleMR R package.²⁹ Additionally, owing to the fact that some traits showed a limited number of associated genome-wide significant variants, genetic instruments associated with PTSD, income, risk-taking behaviors, and trauma exposure were based on suggestive PTs similar to previous MR studies.³³⁻³⁵ We verified these IVW estimates using the MR-Robust Adjusted Profile Score (MR-RAPS) approach, which is a method designed to identify and estimate confounded associations using weak genetic instrument variables.^{36,37} We conducted multiple sensitivity analyses with respect to the MR tests conducted to exclude possible biases (horizontal pleiotropy, ie, the variants included in the genetic instrument having an effect on disease outside their effects on the exposure in MR^{38,39}) under different scenarios in the MR estimates. These included the IVW heterogeneity test,²⁹ the MR-Egger intercept,³⁰ the MR-RAPS overdispersion test,³⁶ and the MR-Pleiotropy Residual Sum and Outlier (MR-PRESSO) global test.⁴⁰ Finally, a leave-1-out analysis was conducted to identify potential outliers among the variants included in the genetic instruments tested. The MR results without evidence of horizontal pleiotropy and heterogeneity were entered in the multivariable MR (MVMR) analysis⁴¹ conducted using the IVW approach. This method permits evaluation of the independent association of each risk factor with the outcome, similar to the simultaneous assessment of several treatments in a factorial randomized trial.⁴¹

Comparing the utility of MR with MVMR methods, MR estimates the total association of the exposure with the outcome, whereas MVMR estimates the direct association of each exposure with the outcome.⁴² In this scenario, MVMR is not a form of mediation analysis but instead estimates the direct association of the exposure with the outcome that does not act via the mediator.⁴² The MVMR analysis was conducted using the MendelianRandomization R package.⁴³

Enrichment Analysis

We tested for functional differences between the traits of interest by means of enrichment analyses based on tissue-specific and cell type–specific gene expression reference panels.⁴⁴⁻⁴⁸ These analyses were conducted using the MAGMA tool⁴⁹ implemented in FUMA.⁵⁰

Results

Data were available for a total of 1 180 352 participants (634 391 [53.7%] women). Information regarding PTSD was available for 23 185 individuals with PTSD and 151 309 control participants (174 494 individuals; 15%) from the PGC-PTSD Working Group and regarding EA for 1 131 881 individuals (96%) from the total sample.

Genetic Correlation and Polygenic Risk Scoring

Our study investigated multiple data sets generated from different cohorts and with different data availability (Table). Using LD score regression and data sets with full GWAS summary association data, we observed a negative genetic correlation of PTSD (freeze-2) with EA ($r_g = -0.26$; SE = 0.05; $P = 4.60 \times 10^{-8}$) and cognitive performance ($r_g = -0.16$; SE = 0.05; $P = 9.00 \times 10^{-4}$). The PRS analyses were conducted using PTSD as the target and traits related to EA as the training data set, considering only genome-wide significant variants ($P < 5.00 \times 10^{-8}$; eFigure 2 in the [Supplement](#)). This analysis was conducted excluding those pairwise comparisons that would have included data sets with the UK Biobank as an overlapping cohort. The most significant PRS association was observed between MC genome-wide significant PRS with respect to PGC-PTSD freeze-2 data ($R^2 = 0.04\%$; SE = 0.0001; $P = 1.13 \times 10^{-8}$). The same PTSD data set showed a weaker association with MA PRS ($R^2 = 0.01\%$; SE = 0.00003; $P = .004$). Significant associations were also observed with respect to PGC-PTSD freeze-1.5 outcome for EA ($R^2 = 0.03\%$; SE = 0.0001; $P = 6.16 \times 10^{-4}$), MC PRS ($R^2 = 0.03\%$; SE = 0.0001; $P = 7.86 \times 10^{-4}$), and EA1M PRS ($R^2 = 0.03\%$; SE = 0.0001; $P = .002$). The phenotypic variance explained by the significant PRS is in line with the cross-phenotype association expected between 2 complex traits with a moderate genetic correlation. No association of the PRS of cognitive performance and MA was observed with respect to PGC-PTSD freeze-1.5 data set (cognitive performance: $R^2 < 0.01\%$; $P = .15$; MA: $R^2 < 0.01\%$; $P = .11$), and accordingly, these phenotypes were not investigated further. We tested the reverse direction (ie, PTSD as base and traits related to EA as target). Owing to the limited number of genome-wide significant loci in the PGC-PTSD analysis, the PRS analysis was conducted considering multiple-association PTs to include at least 10 LD-independent variants in each PRS tested (eFigure 3 in the [Supplement](#)). We observed a nominally significant association between the PGC-PTSD freeze-1.5 PRS (PT = 10^{-5}) and EA ($R^2 = 0.0004\%$; SE = 0.000002; $P = .04$) that would not survive a Bonferroni correction for the number of PTs tested.

Mendelian Randomization


Based on the PRS results, we conducted MR tests using 3 genetic instruments based on genome-wide significant variants. These included MC, EA, and EA1M, tested with respect to PGC-PTSD data sets (freeze-2 and/or freeze-1.5, depending on UK Biobank overlap). The reverse MR test was conducted on the basis of PGC-PTSD freeze-1.5 data (PT = 5.00×10^{-5}) with respect to the EA data set. It was not possible to conduct reverse analyses for the MC and EA1M data sets because we did not have access to the full GWAS summary association data for these studies. A significant association was found between MC and PGC-PTSD freeze-2 data (IVW: $\beta = -0.41$; 95% CI, -0.59 to -0.23 ; $P = 3.46 \times 10^{-6}$). Concordant results were observed when considering other MR methods ([Figure 1](#)). The significance was replicated with the MR-RAPS approach ($\beta = -0.42$; 95% CI, -0.60 to -0.24 ; SE = 0.09; $P = 3.70 \times 10^{-6}$), and no outliers were identified by the leave-1-out analysis (eFigure 4 in the [Supplement](#)). However, we observed the presence of possible bias in this result owing to heterogeneity and/or pleiotropy (IVW heterogeneity test: $Q = 267.4$; $df = 192$; $P = 2.61 \times 10^{-4}$; MR-RAPS overdispersion test: estimated pleiotropy variance, 0.0001; $P = .007$; MR-PRESSO global test: observed residual sum of squares, 281.4; $P = 5.00 \times 10^{-4}$). Thus, we identified the outliers on the basis of the MR-RAPS standardized residuals ($-1.96 > z > 1.96$) and verified their contributions on the results of the IVW heterogeneity test (eFigure 5 in the [Supplement](#)). Removing the outliers from the MC genetic instrument, we confirmed the association

of MC with PTSD in the freeze-2 data set (IVW: $\beta = -0.39$; 95% CI, -0.57 to 0.21 ; $P = 4.25 \times 10^{-7}$; MR-RAPS: $\beta = -0.39$; 95% CI, -0.55 to -0.23 ; $P = 1.06 \times 10^{-6}$) and the lack of evidence of possible confounders from the sensitivity analyses (eTable 2 and eFigure 6 in the Supplement). We verified the reliability of this MR finding considering PGC-PTSD freeze-1.5 data as the outcome and MC, EA, and EA1M as exposures. We observed comparable MR results across the genetic instruments generated from different cohorts and the 2 versions of the PGC-PTSD data sets (Figure 2; eFigure 7 in the Supplement). However, consistent with the lower power of PGC-PTSD freeze-1.5 data, we observed a reduction of the significance (MC for PGC-PTSD freeze-2 data set: IVW: $\beta = -0.41$; 95% CI, -0.59 to -0.23 ; $P = 3.46 \times 10^{-6}$; MC for PGC-PTSD freeze-1.5 data set: IVW: $\beta = -0.22$; 95% CI, -0.38 to -0.06 ; $P = .004$; EA for PGC-PTSD freeze-1.5 data set: $\beta = -0.23$; 95% CI, -0.39 to -0.07 ; $P = .004$). We verified that no bias in the MR results was present owing to palindromic variants with an ambiguous allele frequency⁵¹ and to the presence of assortative mating in EA⁵² (eTable 3 and eAppendix in the Supplement).

To support further that MC and EA are associated with the same mechanism, we conducted a MVMR analysis, which showed that these 2 associations are not independent from each other (eFigure 8 in the Supplement), and their relationship with PTSD should be shared. Accordingly, we used EA as a proxy of MC in the subsequent analyses because we had full access only to the former data set.


We tested the reverse association, considering PTSD as the risk factor (ie, exposure) and EA as the outcome of the MR analysis. We included PTSD genetic instrument variants with a PTSD GWAS

Figure 1. Single-Nucleotide Polymorphism (SNP) Repeated Effects on Posttraumatic Stress Disorder (PTSD) and Most Advanced Math Course Completed (MC)



SNP exposure (MC associations, β) and SNP outcome (PTSD freeze-2 associations, log odds ratio [OR]) coefficients used in the mendelian randomization analysis. Crosses represent 95% CIs for each association.

Figure 2. Estimated Associations Considering Different Traits Associated With Educational Attainment (EA) and 2 Versions of the Posttraumatic Stress Disorder (PTSD) Data Set



EA1M indicates sample including participants from Social Science Genetic Association Consortium, UK Biobank, and 23andMe; IVW, inverse-variance weighted; MC, most advanced math course completed; MR-RAPS, mendelian randomization-robust adjusted profile score; PTSD2, PTSD freeze 2 data set; and PTSD1.5, PTSD freeze 1.5 data set.

$P = 10^{-5}$ considering the PGC-PTSD freeze-1.5 data. No significant association was observed (IVW: $\beta = 0.006$; 95% CI, -0.002 to 0.014 ; $P = .16$; MR-RAPS: $\beta = 0.0006$; 95% CI, -0.007 to 0.008 ; $P = .10$). To further confirm the absence of reverse association, we conducted an MR analysis including all LD-independent variants in the genetic instrument and applied the MR-RAPS method only. No directional association of PTSD with EA was observed ($\beta = -0.0006$; 95% CI, -0.002 to 0.002 ; $P = .55$), but we confirmed a directional association of EA with PTSD ($\beta = -0.27$; 95% CI, -0.38 to -0.15 ; $P = 8.06 \times 10^{-6}$). These outcomes were stable across different adjustments of the MR-RAPS method (eTable 4 in the [Supplement](#)).

Multivariable MR Analysis

To further investigate the association of EA with PTSD, we tested 3 potential mediators (risk-taking behaviors, income, and trauma exposure) in an MVMR analysis. Before entering these potential mediators in the MVMR analysis, we verified the reliability of each genetic instrument by conducting a standard 2-sample MR and verifying the evidence of bias owing to heterogeneity and horizontal pleiotropy. Because of the limited number of genome-wide significant variants with respect to these traits, we conducted a PRS considering multiple PTs as described earlier, to determine the best genetic instrument for each trait with respect to the PGC-PTSD freeze-1.5 data set (eFigure 9 in the [Supplement](#)). The best results were observed for PT = 5.00×10^{-4} with risk-taking behaviors ($R^2 = 0.06\%$; SE = 0.0001; $P = 1.53 \times 10^{-5}$) and PT = .001 for income ($R^2 = 0.09\%$; SE = 0.0002; $P = 5.67 \times 10^{-8}$). The MR analysis based on these genetic instruments confirmed that PTSD is associated with the genetic instruments related to risk-taking behaviors (IVW: $\beta = 0.76$; 95% CI, 0.38 to 1.13; $P = 1.13 \times 10^{-4}$; MR-RAPS: $\beta = 0.76$; 95% CI, 0.32 to 1.21; $P = 6.75 \times 10^{-4}$), and income (IVW: $\beta = -0.18$; 95% CI, -0.29 to -0.07 ; $P = .001$; MR-RAPS: $\beta = -0.19$; 95% CI, -0.31 to -0.07 ; $P = .003$). No evidence of heterogeneity or pleiotropy was observed in either analysis (eTable 5 in the [Supplement](#)).

Multiple traumatic events were assessed in the UK Biobank (eTable 1 in the [Supplement](#)), and they showed genetic correlation with each other (eTable 6 in the [Supplement](#)). We selected 4 traumatic experiences that showed a similar pattern of genetic correlation with respect to PTSD, EA, and the other 2 potential mediators (eFigure 10 in the [Supplement](#)). The most informative PRS across the 4 traumatic experiences tested was observed at PT = .001 (eFigure 11 in the [Supplement](#)). Then, we conducted an MR analysis testing different trauma-related genetic instruments with respect to PGC-PTSD freeze-1.5 data. We observed significant associations not affected by confounders (eTable 7 in the [Supplement](#)) for 3 traumatic experiences and, conducting an MVMR analysis, identified "physically abused by family as a child" (UK Biobank data field 20488) as the most informative genetic instrument for trauma exposure (MR analysis: $\beta = 0.36$; 95% CI, 0.19 to 0.52; $P = 2.57 \times 10^{-5}$; MVMR analysis: $\beta = 0.26$; 95% CI, -0.51 to 0.01 ; SE = 0.129; $P = .04$) (eFigure 12 in the [Supplement](#)).


In the MVMR analysis, we observed that trauma exposure and risk-taking behaviors were independent risk factors for PTSD (ie, the results obtained from the MR IVW and MVMR IVW analyses were both significant; **Figures 3A, B, and C**). Conversely, the genetic instrument related to income potentially mediates the association of the EA genetic instrument with PTSD (Figure 3A and D). Educational attainment has a significant association with respect to PTSD ($\beta = -0.23$; 95% CI, -0.39 to -0.07 ; $P = .004$), but when adjusted by income, this result is null ($\beta = -0.04$; 95% CI -0.30 to 0.21 ; $P = .79$). Conversely, the association of income with PTSD was still significant when adjusted by EA (unadjusted: $\beta = -0.18$; 95% CI, -0.30 to -0.06 ; $P = .001$; adjusted: $\beta = -0.32$; 95% CI, -0.57 to -0.07 ; SE = 0.13; $P = .02$).

Enrichment Analysis

Although there is a large genetic correlation between EA and income ($r = 0.81$; $P < 6.10 \times 10^{-308}$), income is significantly more correlated with PTSD than EA (EA: $r = -0.26$; SE = 0.05; $P = 4.60 \times 10^{-8}$; income: $r = -0.45$; SE = 0.06; $P = 9.98 \times 10^{-16}$; z for EA \times income = 2.65; P for


EA \times income = 0.008). Both traits are enriched for the transcriptomic profile of multiple brain tissues (eg, cerebellar hemisphere, EA: β = 0.01; 95% CI, 0.07-0.12; P = 1.49×10^{-16} ; income: β = 0.04; 95% CI, 0.02-0.06; P = 3.83×10^{-7}) and neuronal cell types (eg, γ -aminobutyric acid [GABA]-ergic neurons, EA: β = 0.14; 95% CI, 0.07-0.21; P = 8.96×10^{-5} ; income: β = 0.05; 95% CI, 0.01-0.09; P = .02), but the enrichment signals of the EA GWAS are more significant than the ones observed in the income GWAS (**Figure 4**; eTable 8 in the [Supplement](#)), showing that EA data are more informative for the brain processes expected to be associated with cognition than income data.

Figure 3. Multivariable Mendelian Randomization Analysis Considering Associations of Educational Attainment (EA), Posttraumatic Stress Disorder (PTSD), and Other Traits



IVW indicates inverse-variance weighted.

Figure 4. Enrichment Analyses for Educational Attainment and Income



A, Tissue-specific enrichments (brain tissues are indicated with solid circles; nonbrain tissues are indicated with open circles). B, Cell-type enrichments that were at least nominally significant ($P < .05$) in both traits. BA9 indicates Brodmann area 9; BA24,

Brodmann area 24; exCA1, excitatory neurons in hippocampal subfield CA1; exDG, excitatory neurons in dentate gyrus; exPFC1, excitatory neurons in prefrontal cortex 1; GABA, γ -aminobutyric acid; and PFC, prefrontal cortex.

Discussion

Our analysis was based mainly on data from investigations of traits associated with EA. Previous studies have demonstrated that genetic results deriving from them are mainly informative regarding the brain mechanisms at the basis of human cognitive ability.¹⁶⁻¹⁹ Our analysis made use of these data to show that traits associated with cognitive ability have an association with PTSD. This result was consistent across traits in independent cohorts, even when adjusting the analysis for assortative mating present in these traits.⁵² In line with an association direction from cognition to PTSD, no evidence of reverse association was observed.

Although our findings are consistent with a specific direction, we had to evaluate whether other factors could be responsible for this association. Accordingly, we tested 3 phenotypes of known relevance for PTSD: propensity to trauma exposure,⁵³ risk-taking behaviors,⁵⁴ and economic status.⁵⁵ Our MVMR analysis clearly showed that, while propensity to trauma exposure and risk-taking behaviors are independent PTSD risk factors, the directional association of EA with PTSD is associated with economic status, which is the driving force of the association. Educational attainment and income showed a large genetic overlap, and while PTSD showed a higher correlation with income than with EA, our investigation showed that EA is more informative for the brain mechanisms that are considered responsible for a predisposition to high cognitive ability. Since income appears to be responsible for the EA-PTSD association, we hypothesize that this mechanism is not related to cognitive ability but rather to other risk factors. This is also supported by our finding that, unlike EA and MC (traits that should be more associated with socioeconomic status), cognitive performance and MA showed a lower genetic association with PTSD. A 2018 genome-wide investigation of social stratification⁵⁶ showed that cognition and socioeconomic status are correlated with a wide range of factors, including personality, psychological traits, mental health, substance use, physical health, reproductive behaviors, and anthropometric traits. Additionally, income may reflect indirect effects, such as those induced by genetic nurture on EA.^{57,58} Socioeconomic factors might also be associated with the outcome of PTSD, given that individuals with poorer outcomes are more likely to be included in studies of prevalent cases.

To our knowledge, this study represents the first MR analysis to investigate the underlying mechanisms linking cognitive ability to PTSD. It is based on the largest genome-wide data sets for these traits available at this time. The EA-PTSD association observed is in line with several prospective studies,¹ and the association of socioeconomic status with PTSD is also a well-established risk factor reported in several observational studies.^{55,59} Compared with these previous investigations, the current analyses are based on a much larger population (>1 million individuals) than would ever be feasible for a traditional experimental design (which would include randomized interventions and measurements of outcomes) and without the ethical quandaries that would accompany such randomizations. Also, these results are not expected to be affected by reverse association because of the genetic information used. The findings show that income may explain the EA-PTSD association, suggesting that brain mechanisms related to cognitive ability are not directly responsible for the association observed.

Limitations

Our study has limitations. The present study is based on genetic information generated from the investigation of heterogeneous, clinically defined phenotypes such as PTSD. We tested this complex trait with respect to a series of complex, socially contextualized phenotypes, including EA, income, risk-taking behaviors, and predisposition to traumatic events. Although we used appropriate statistical methods and conducted the analyses across multiple independent cohorts, findings related to genetic data associated with these phenotypes need to be interpreted cautiously.^{58,60} The results of our current analysis are also limited by the statistical power of the PTSD GWAS data sets, which may have limited our ability to observe the reverse association of PTSD with EA. However, we used multiple methods that showed significant associations when applied to similarly powered

GWAS data sets. Another potential limitation is owing to the pervasive presence of horizontal pleiotropy among complex traits.³⁹ We applied multiple sensitivity analyses that accounted for different scenarios related to the potential confounding effect of horizontal pleiotropy and heterogeneity in the genetic instruments applied in our MR analyses. Although no evidence of bias was observed by the methods used, our current findings could be affected by an unaccounted confounder.

Conclusions

This study provides new evidence to elucidate the association of EA with PTSD, pointing toward risk factors associated with economic status rather than brain pathways. These findings have relevant implications with respect to our understanding of the pretrauma risk factors associated with increased vulnerability to PTSD. Additionally, MR analysis should consider testing the independence of multiple correlated risk factors with respect to the outcome of interest. This is particularly relevant when investigating the potential role of EA in human phenotypes and disorders.

ARTICLE INFORMATION

Accepted for Publication: March 19, 2019.

Published: May 3, 2019. doi:[10.1001/jamanetworkopen.2019.3447](https://doi.org/10.1001/jamanetworkopen.2019.3447)

Open Access: This is an open access article distributed under the terms of the [CC-BY License](https://creativecommons.org/licenses/by/4.0/). © 2019 Polimanti R et al. *JAMA Network Open*.

Corresponding Author: Renato Polimanti, PhD, Yale University School of Medicine, Veterans Affairs Connecticut Healthcare Center, 116A2, 950 Campbell Ave, West Haven, CT 06516 (renato.polimanti@yale.edu).

Author Affiliations: Department of Psychiatry, Yale University School of Medicine, West Haven, Connecticut (Polimanti, Gelernter); Veterans Affairs Connecticut Healthcare Center, West Haven, Connecticut (Polimanti, Gelernter); Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York (Ratanatharathorn); Department of Psychiatry, School of Medicine, University of California, San Diego (Maihofer, M. B. Stein, Nievergelt); Veterans Affairs Center of Excellence for Stress and Mental Health and Research Service, Veterans Affairs San Diego Healthcare System, San Diego, California (Maihofer, Nievergelt); Stanley Center for Psychiatric Research, Broad Institute, Cambridge, Massachusetts (Choi); Department of Psychiatry, Massachusetts General Hospital, Boston (Choi); Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Choi, Koenen); Psychiatry Service, Veterans Affairs San Diego Healthcare System, San Diego, California (M. B. Stein); Department of Psychiatry, Duke University, Durham, North Carolina (Morey); Durham Veterans Affairs Medical Center, Durham, North Carolina (Morey); National Center for PTSD Behavioral Science Division, Veterans Affairs Boston Healthcare System, Boston, Massachusetts (Logue); Department of Psychiatry, School of Medicine, Boston University, Boston, Massachusetts (Logue); South African Medical Research Council Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry, University of Cape Town, Cape Town, South Africa (D. J. Stein); Department of Genetics, School of Medicine, Yale University, New Haven, Connecticut (Gelernter); Department of Neuroscience, School of Medicine, Yale University, New Haven, Connecticut (Gelernter).

Author Contributions: Dr Polimanti had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Polimanti, Choi, M. B. Stein, Morey, Koenen.

Acquisition, analysis, or interpretation of data: Polimanti, Ratanatharathorn, Maihofer, Choi, Logue, Nievergelt, D. J. Stein, Koenen, Gelernter.

Drafting of the manuscript: Polimanti.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Polimanti, Ratanatharathorn, Koenen.

Obtained funding: Nievergelt, Koenen.

Administrative, technical, or material support: Logue, Nievergelt, Koenen, Gelernter.

Supervision: Gelernter.

Conflict of Interest Disclosures: Dr Choi reported grants from National Institute of Mental Health during the conduct of the study. Dr M. B. Stein reported grants from the US Department of Defense and the National Institute of Mental Health during the conduct of the study and receiving payments for his editorial work on the journals *Biological Psychiatry* and *Depression and Anxiety* as well as the health professional reference UpToDate. Dr D. J. Stein reported grants from the South African Medical Research Council during the conduct of the study and receiving personal fees from Lundbeck and Sun Pharmaceutical Industries. Dr Gelernter reported grants from the National Institutes of Health during the conduct of the study. No other disclosures were reported.

Funding/Support: This study was supported by grants R01 MH106595 and U01 MH109532 from the National Institutes of Health and by the Veterans Affairs National Center for Posttraumatic Stress Disorder Research.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Group Information: The Psychiatric Genomics Consortium Posttraumatic Stress Disorder (PGC-PTSD) Working Group members are Caroline M. Nievergelt (Department of Psychiatry, University of California, San Diego); Adam X. Maihofer (Department of Psychiatry, University of California, San Diego); Torsten Klengel (Department of Psychiatry, Harvard Medical School, Boston, Massachusetts); Elizabeth G. Atkinson (Broad Institute, Stanley Center for Psychiatric Research, Cambridge, Massachusetts); Chia-Yen Chen (Broad Institute, Stanley Center for Psychiatric Research); Karmel W. Choi (Broad Institute, Stanley Center for Psychiatric Research); Jonathan R. I. Coleman (Institute of Psychiatry, Psychology and Neuroscience, Social, Genetic and Developmental Psychiatry Centre, King's College London, London, United Kingdom); Shareefa Dalvie (South African Medical Research Council Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry, University of Cape Town, Cape Town, Western Cape, South Africa); Laramie E. Duncan (Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, California); Mark W. Logue (Veterans Affairs Boston Healthcare System, National Center for PTSD, Boston, Massachusetts); Allison C. Provost (Cohen Veterans Bioscience, Cambridge, Massachusetts); Andrew Ratanatharathorn (Department of Epidemiology, Harvard T.H. Chan School of Public Health, Harvard University, Boston, Massachusetts); Murray B. Stein (Department of Psychiatry, University of California, San Diego); Katy Torres (Department of Psychiatry, University of California, San Diego); Allison E. Aiello (Department of Epidemiology, University of North Carolina at Chapel Hill); Lynn M. Almli (Carter Consulting, Atlanta, Georgia); Ananda B. Amstadter (Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics, Richmond); Søren B. Andersen (Research and Knowledge Centre, The Danish Veteran Centre, Ringsted, Denmark); Ole A. Andreassen (Institute of Clinical Medicine, University of Oslo, Oslo, Norway); Paul A. Arbisi (Mental Health Service Line, Minneapolis Veterans Affairs Health Care System, Minneapolis, Minnesota); Allison E. Ashley-Koch (Department of Psychiatry, Duke University, Durham, North Carolina); S. Bryn Austin (Division of Adolescent and Young Adult Medicine, Boston Children's Hospital, Boston, Massachusetts); Esmina Avdibegovic (Department of Psychiatry, University Clinical Center of Tuzla, Tuzla, Bosnia and Herzegovina); Dragan Babić (Department of Psychiatry, University Clinical Center of Mostar, Mostar, Bosnia and Herzegovina); Marie Bækvad-Hansen (Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark); Dewleen G. Baker (Department of Psychiatry, University of California, San Diego); Jean C. Beckham (Department of Psychiatry and Behavioral Sciences, Duke University, Durham, North Carolina); Laura J. Bierut (Department of Psychiatry, Washington University in Saint Louis School of Medicine, St Louis, Missouri); Jonathan I. Bisson (Medical Research Council Centre for Psychiatric Genetics and Genomics, National Centre for Mental Health, Cardiff University, Cardiff, United Kingdom); Marco P. Boks (Utrecht Brain Center Rudolf Magnus, Department of Translational Neuroscience, University Medical Center, Utrecht, the Netherlands); Elizabeth A. Bolger (Department of Psychiatry, Harvard Medical School); Anders D. Børglum (Centre for Integrative Sequencing, iSEQ, Aarhus University, Aarhus, Denmark); Bekh Bradley (Mental Health Service Line, Atlanta Veterans Affairs Health Care System, Decatur, Georgia); Megan Brashear (School of Public Health and Department of Epidemiology, Louisiana State University Health Sciences Center, New Orleans); Jerome Breen (National Institute for Health Research Biomedical Research Centre at the Maudsley, King's College London, London, United Kingdom); Richard A. Bryant (Department of Psychology, University of New South Wales, Sydney, Australia); Angela C. Bustamante (Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor); Jonas Bybjerg-Grauholm (Department for Congenital Disorders, Statens Serum Institut); Joseph R. Calabrese (Department of Psychiatry, University Hospitals, Cleveland, Ohio); José M. Caldas-de-Almeida (Chronic Diseases Research Centre [CEDOC], Lisbon Institute of Global Mental Health, Lisbon, Portugal); Anders M. Dale (Department of Radiology, Department of Neurosciences, University of California, San Diego); Mark J. Daly (Psychiatric and Neurodevelopmental Genetics Unit [PNGU], Massachusetts General Hospital, Boston); Nikolaos P. Daskalakis (Cohen Veterans Bioscience); Jürgen Deckert (Center of Mental Health, Psychiatry, Psychosomatics and Psychotherapy, University Hospital of Würzburg, Würzburg, Germany); Douglas L. Delahanty (Department of Psychological Sciences, Kent State University, Kent, Ohio); Michelle F. Dennis (Duke Molecular Physiology Institute, Duke University, Durham, North Carolina); Seth G. Disner (Research Service Line, Minneapolis Veterans Affairs

Health Care System, Minneapolis, Minnesota); Katharina Domschke (Department of Psychiatry and Psychotherapy, Faculty of Medicine, Medical Center, University of Freiburg, Freiburg, Germany); Alma Dzubur-Kulenovic (University Clinical Center of Sarajevo, Department of Psychiatry, Sarajevo, Bosnia and Herzegovina); Christopher R. Erbes (Center for Care Delivery and Outcomes Research [CCDOR], Minneapolis Veterans Affairs Health Care System, Minneapolis, Minnesota); Alexandra Evans (Medical Research Council Centre for Psychiatric Genetics and Genomics, National Centre for Mental Health, Cardiff University); Lindsay A. Farrer (Department of Medicine, Boston University School of Medicine, Boston, Massachusetts); Norah C. Feeny (Department of Psychological Sciences, Case Western Reserve University, Cleveland, Ohio); Janine D. Flory (Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York); David Forbes (Department of Psychiatry, University of Melbourne, Melbourne, Australia); Carol E. Franz (Department of Psychiatry, University of California, San Diego); Sandro Galea (Department of Psychological and Brain Sciences, Boston University); Melanie E. Garrett (Department of Psychiatry and Behavioral Sciences, Duke University); Bizu Gelaye (Department of Epidemiology, Harvard T.H. Chan School of Public Health, Harvard University); Joel Gelernter (Department of Psychiatry, US Department of Veterans Affairs, West Haven, Connecticut); Elbert Geuze (Research Center Military Mental Healthcare, Netherlands Ministry of Defence, Utrecht, the Netherlands); Charles Gillespie (Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, Georgia); Aferdita Goci Uka (Department of Psychiatry, University Clinical Centre of Kosovo, Prishtina, Republic of Kosovo); Scott D. Gordon (Department of Genetics and Computational Biology, Queensland Institute of Medical Research (QIMR) Berghofer Medical Research Institute, Brisbane, Australia); Guia Guffanti (Department of Psychiatry, Harvard Medical School); Rasha Hammamieh (US Army Center for Environmental Health Research, Army Medical Research and Materiel Command, Fort Detrick, Maryland); Supriya Harnal (Stanley Center for Psychiatric Research, Broad Institute); Michael A. Hauser (Department of Psychiatry and Behavioral Sciences, Duke University); Andrew C. Heath (Department of Genetics, Washington University in Saint Louis School of Medicine, St Louis, Missouri); Sian M.J. Hemmings (Department of Psychiatry, Stellenbosch University Faculty of Medicine and Health Sciences, Cape Town, South Africa); David Michael Hougaard (Department for Congenital Disorders, Statens Serum Institut); Miro Jakovljevic (Department of Psychiatry, University Hospital Center of Zagreb, Zagreb, Croatia); Marti Jett (US Army Center for Environmental Health Research, Army Medical Research and Materiel Command); Eric Otto Johnson (Behavioral Health and Criminal Justice Division, Research Triangle Institute International, Research Triangle Park, North Carolina); Ian Jones (Medical Research Council Centre for Psychiatric Genetics and Genomics, National Centre for Mental Health, Cardiff University); Tanja Jovanovic (Department of Psychiatry and Behavioral Sciences, Emory University); Xue-Jun Qin (Duke Molecular Physiology Institute, Duke University, Durham, North Carolina); Angela G. Junglen (Department of Psychological Sciences, Kent State University); Karen-Inge Karstoft (Research and Knowledge Centre, the Danish Veteran Center, Ringsted, Denmark); Milissa L. Kaufman (Department of Psychiatry, Harvard Medical School); Ronald C. Kessler (Department of Psychiatry, Harvard Medical School); Alaptagin Khan (Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts); Nathan A. Kimbrel (Duke Molecular Physiology Institute, Duke University); Anthony P. King (Department of Psychiatry, University of Michigan Medical School, Ann Arbor); Nastassja Koen (South Africa Medical Research Council Unit on Risk and Resilience in Mental Disorders, University of Cape Town); Henry R. Kranzler (Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia); William S. Kremen (Department of Psychiatry and Department of Family Medicine and Public Health, University of California, San Diego); Bruce R. Lawford (Queensland University of Technology, Institute of Health and Behavioral Innovation, Brisbane, Australia); Lauren A. M. Lebois (Department of Psychiatry, Harvard Medical School); Catrin E. Lewis (Medical Research Council Centre for Psychiatric Genetics and Genomics, National Centre for Mental Health, Cardiff University); Sarah D. Linnstaedt (Department of Anesthesiology, University of North Carolina Institute for Trauma Recovery, Chapel Hill); Adriana Lori (Department of Gynecology and Obstetrics, Emory University, Atlanta, Georgia); Bozo Lugonja (Medical Research Council Centre for Psychiatric Genetics and Genomics, National Centre for Mental Health, Cardiff University); Jurjen J. Luykx (Utrecht Brain Center Rudolf Magnus, Department of Psychiatry, University Medical Center); Michael J. Lyons (Dean's Office, Boston University, Boston, Massachusetts); Jessica Maples-Keller (Department of Psychiatry and Behavioral Sciences, Emory University); Charles Marmar (Department of Psychiatry, New York University, New York); Alicia R. Martin (Broad Institute, Stanley Center for Psychiatric Research); Nicholas G. Martin (Department of Genetics and Computational Biology, Queensland Institute of Medical Research [QIMR] Berghofer Medical Research Institute); Douglas Maurer (Command, US Army, Fort Sill, Oklahoma); Matig R. Mavissakalian (Department of Psychiatry, University Hospitals); Alexander McFarlane (Department of Psychiatry, University of Adelaide, Adelaide, Australia); Regina E. McGlinchey (GRECC/TRACTS, Veterans Affairs Boston Health Care System, Boston, Massachusetts); Katie A. McLaughlin (Department of Psychology, Harvard University); Samuel A. McLean (Department of Anesthesiology, University of North Carolina Institute for Trauma Recovery); Sarah McLeay (PTSD Initiative, Gallipoli Medical Research Institute, Greenslopes, Australia); Divya Mehta (Faculty of Health, Queensland University of Technology, Brisbane, Australia); William P. Milberg (GRECC/TRACTS, Veterans Affairs Boston Health Care System); Mark W. Miller (Veterans Affairs Boston Healthcare System, National Center for PTSD); Rajendra A. Morey (Duke Molecular Physiology Institute, Duke

University); Charles Phillip Morris (Faculty of Health, Queensland University of Technology); Ole Mors (Psychosis Research Unit, Aarhus University Hospital, Risskov, Denmark); Preben B. Mortensen (Centre for Integrated Register-Based Research, Aarhus University, Aarhus, Denmark); Benjamin M. Neale (Broad Institute, Stanley Center for Psychiatric Research); Elliot C. Nelson (Department of Psychiatry, Washington University in Saint Louis School of Medicine); Merete Nordentoft (The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Aarhus, Denmark); Sonya B. Norman (Executive Division, National Center for Posttraumatic Stress Disorder, White River Junction, Vermont); Meaghan O'Donnell (Department of Psychiatry, University of Melbourne, Melbourne, Australia); Holly K. Orcutt (Department of Psychology, Northern Illinois University, DeKalb); Matthew S. Panizzon (Department of Psychiatry, University of California, San Diego); Edward S. Peters (School of Public Health and Department of Epidemiology, Louisiana State University Health Sciences Center); Alan L. Peterson (Department of Psychology, University of Texas, San Antonio); Matthew Peverill (Department of Psychology, University of Washington, Seattle); Robert H. Pietrzak (US Department of Veterans Affairs National Center for Posttraumatic Stress Disorder, West Haven, Connecticut); Melissa A. Polusny (Department of Mental Health, Minneapolis Veterans Affairs Health Care System); John P. Rice (Department of Psychiatry, Washington University in Saint Louis School of Medicine); Stephan Ripke (Stanley Center for Psychiatric Research, Broad Institute); Victoria B. Risbrough (Department of Psychiatry, University of California, San Diego); Andrea L. Roberts (Department of Environmental Health, Harvard T.H. Chan School of Public Health, Harvard University); Alex O. Rothbaum (Department of Psychological Sciences, Case Western Reserve University); Barbara O. Rothbaum (Department of Psychiatry and Behavioral Sciences, Emory University); Peter Roy-Byrne (Department of Psychology, University of Washington); Ken Ruggiero (Department of Nursing and Department of Psychiatry, Medical University of South Carolina, Charleston); Ariane Rung (School of Medicine and Department of Physiology, Louisiana State University Health Sciences Center); Bart P. F. Rutten (School for Mental Health and Neuroscience, Department of Psychiatry and Neuropsychology, Maastricht Universitair Medisch Centrum, Maastricht, the Netherlands); Nancy L. Saccone (Department of Psychiatry, Washington University in Saint Louis School of Medicine); Sixto E. Sanchez (Facultad de Ciencias de la Salud, Department of Medicine, Universidad Peruana de Ciencias Aplicadas, Lima, Peru); Dick Schijven (Utrecht Brain Center Rudolf Magnus, Department of Psychiatry, University Medical Center); Soraya Seedat (Department of Psychiatry, Stellenbosch University Faculty of Medicine and Health Sciences); Antonia V. Seligowski (Department of Pediatrics, Harvard Medical School, Boston, Massachusetts); Julia S. Seng (School of Nursing, University of Michigan, Ann Arbor); Christina M. Sheerin (Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics); Derrick Silove (Department of Psychiatry, University of New South Wales, Sydney, Australia); Alicia K. Smith (Department of Gynecology and Obstetrics, Emory University); Jordan W. Smoller (Broad Institute, Stanley Center for Psychiatric Research); Nadia Solovieff (Massachusetts General Hospital, Boston); Scott R. Sponheim (Mental Health Service Line, Minneapolis Veterans Affairs Health Care System); Dan J. Stein (South African Medical Research Council Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry, University of Cape Town); Jennifer A. Sumner (Department of Medicine, Columbia University Medical Center, New York, New York); Martin H. Teicher (Department of Psychiatry, Harvard Medical School); Wesley K. Thompson (Institute of Biological Psychiatry, Mental Health Centre, Sankt Hans, Roskilde, Denmark); Edward Trapido (School of Public Health and Department of Epidemiology, Louisiana State University Health Sciences Center); Monica Uddin (Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign); Robert J. Ursano (Department of Psychiatry, Uniformed Services University, Bethesda, Maryland); Leigh Luella van den Heuvel (Department of Psychiatry, Stellenbosch University Faculty of Medicine and Health Sciences); Miranda van Hooff (Department of Psychiatry, University of Adelaide); Eric Vermetten (Arq Psychotrauma Research Expert Group, Diemen, the Netherlands); Christiaan H. Vinkers (Department of Anatomy and Neurosciences, Amsterdam Academic Medical Center, Amsterdam, the Netherlands); Joanne Voisey (Queensland University of Technology, Institute of Health and Behavioral Innovation); Yunpeng Wang (Institute of Biological Psychiatry, Mental Health Centre); Zhewu Wang (Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina); Thomas Werge (Institute of Biological Psychiatry, Mental Health Centre); Michelle A. Williams (Department of Epidemiology, Harvard T.H. Chan School of Public Health, Harvard University); Douglas E. Williamson (Department of Psychiatry and Behavioral Sciences, Duke University); Sherry Winternitz (Department of Psychiatry, Harvard Medical School); Christiane Wolf (Center of Mental Health, Psychiatry, Psychosomatics and Psychotherapy, University Hospital of Würzburg); Erika J. Wolf (Veterans Affairs Boston Healthcare System, National Center for PTSD); Jonathan D. Wolff (McLean Hospital, Belmont, Massachusetts); Rachel Yehuda (Department of Psychiatry, Icahn School of Medicine at Mount Sinai); Keith A. Young (Department of Psychiatry, Baylor Scott and White Central Texas, Temple); Ross McD Young (School of Psychology and Counseling, Queensland University of Technology, Brisbane, Australia); Hongyu Zhao (Department of Biostatistics, Yale University, New Haven, Connecticut); Lori A. Zoellner (Department of Psychiatry and Behavioral Sciences, University of Washington); Israel Liberzon (Department of Psychiatry, University of Michigan Medical School); Kerry J. Ressler (Department of Psychiatry and Behavioral Sciences, Emory University); Magali Haas (Cohen Veterans Bioscience); and Karestan C. Koenen (Department of Epidemiology, Harvard School of Public Health).

Additional Contributions: We thank study participants and research groups contributing to the Psychiatric Genomics Consortium Posttraumatic Stress Disorder Working Group for sharing their data and the members of the other cited consortia for making their data available.

REFERENCES

1. DiGangi JA, Gomez D, Mendoza L, Jason LA, Keys CB, Koenen KC. Pretrauma risk factors for posttraumatic stress disorder: a systematic review of the literature. *Clin Psychol Rev*. 2013;33(6):728-744. doi:10.1016/j.cpr.2013.05.002
2. Goldstein RB, Smith SM, Chou SP, et al. The epidemiology of DSM-5 posttraumatic stress disorder in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51(8):1137-1148. doi:10.1007/s00127-016-1208-5
3. Frankenberg E, Sikoki B, Sumantri C, Suriastini W, Thomas D. Education, vulnerability, and resilience after a natural disaster. *Ecol Soc*. 2013;18(2):16. doi:10.5751/ES-05377-180216
4. Gale CR, Deary IJ, Boyle SH, Barefoot J, Mortensen LH, Batty GD. Cognitive ability in early adulthood and risk of 5 specific psychiatric disorders in middle age: the Vietnam Experience Study. *Arch Gen Psychiatry*. 2008;65(12):1410-1418. doi:10.1001/archpsyc.65.12.1410
5. Kremen WS, Koenen KC, Boake C, et al. Pretrauma cognitive ability and risk for posttraumatic stress disorder: a twin study. *Arch Gen Psychiatry*. 2007;64(3):361-368. doi:10.1001/archpsyc.64.3.361
6. Bowling A, Ebrahim S. *Handbook of Health Research Methods*. Maidenhead, England: McGraw-Hill, Open University Press; 2005.
7. Szklo M, Nieto FJ. *Epidemiology: Beyond the Basics*. 4th ed. Sudbury, MA: Jones and Bartlett Publishers; 2006.
8. Rothman KJ, Greenland S, Lash TL. *Modern Epidemiology*. 3rd ed. Philadelphia, PA: Wolters Kluwer/Lippincott Williams & Wilkins; 2008.
9. Dick DM. Mapping risk from genes to behavior: the enduring and evolving influence of Irving Gottesman's endophenotype concept. *Twin Res Hum Genet*. 2018;21(4):306-309. doi:10.1017/thg.2018.35
10. Ravera S, Carrasco N, Gelernter J, Polimanti R. Phenomic impact of genetically-determined euthyroid function and molecular differences between thyroid disorders. *J Clin Med*. 2018;7(10):E296. doi:10.3390/jcm7100296
11. Polimanti R, Amstadter AB, Stein MB, et al; Psychiatric Genomics Consortium Posttraumatic Stress Disorder Workgroup. A putative causal relationship between genetically determined female body shape and posttraumatic stress disorder. *Genome Med*. 2017;9(1):99. doi:10.1186/s13073-017-0491-4
12. Polimanti R, Gelernter J, Stein DJ. Genetically determined schizophrenia is not associated with impaired glucose homeostasis. *Schizophr Res*. 2018;195:286-289. doi:10.1016/j.schres.2017.10.033
13. Polimanti R, Peterson RE, Ong JS, et al. Evidence of causal effect of major depression on alcohol dependence: findings from the Psychiatric Genomics Consortium [published online September 9, 2018]. *bioRxiv*. doi:10.1101/412098
14. Wendt FR, Carvalho C, Gelernter J, Polimanti R. DRD2 and FOXP2 are implicated in the associations between computerized device use and psychiatric disorders [published online December 17, 2018]. *bioRxiv*. doi:10.1101/497420
15. Davey Smith G, Ebrahim S. Mendelian randomization: genetic variants as instruments for strengthening causal inference in observational studies. In: Weinstein M, Vaupel JW, Wachter KW, eds. *Biosocial Surveys*. Washington, DC: The National Academies Press; 2008:428.
16. Lee JJ, Wedow R, Okbay A, et al; 23andMe Research Team; COGENT (Cognitive Genomics Consortium); Social Science Genetic Association Consortium. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet*. 2018;50(8):1112-1121. doi:10.1038/s41588-018-0147-3
17. Okbay A, Beauchamp JP, Fontana MA, et al; LifeLines Cohort Study. Genome-wide association study identifies 74 loci associated with educational attainment. *Nature*. 2016;533(7604):539-542. doi:10.1038/nature17671
18. Savage JE, Jansen PR, Stringer S, et al. Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet*. 2018;50(7):912-919. doi:10.1038/s41588-018-0152-6
19. Selzam S, Krapohl E, von Stumm S, et al. Predicting educational achievement from DNA. *Mol Psychiatry*. 2017;22(2):267-272. doi:10.1038/mp.2016.107
20. Anttila V, Bulik-Sullivan B, Finucane HK, et al; Brainstorm Consortium. Analysis of shared heritability in common disorders of the brain. *Science*. 2018;360(6395):eaap8757. doi:10.1126/science.aap8757

21. Nievergelt CM, Maihofer AX, Klengel T, et al. Largest genome-wide association study for PTSD identifies genetic risk loci in European and African ancestries and implicates novel biological pathways [published online November 1, 2018]. *bioRxiv*. doi:10.1101/458562
22. Little J, Higgins JP, Ioannidis JP, et al; Strengthening the Reporting of Genetic Association Studies. Strengthening the Reporting of Genetic Association Studies (STREGA): an extension of the STROBE statement. *PLoS Med*. 2009;6(2):e22. doi:10.1371/journal.pmed.1000022
23. Neale Lab. GWAS of the UK Biobank. <http://www.nealelab.is/uk-biobank/>. Accessed August 2, 2018.
24. Bulik-Sullivan B, Finucane HK, Anttila V, et al; ReproGen Consortium; Psychiatric Genomics Consortium; Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3. An atlas of genetic correlations across human diseases and traits. *Nat Genet*. 2015;47(11):1236-1241. doi:10.1038/ng.3406
25. Auton A, Brooks LD, Durbin RM, et al; 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*. 2015;526(7571):68-74.
26. Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics*. 2015;31(9):1466-1468. doi:10.1093/bioinformatics/btu848
27. Dastani Z, Hivert MF, Timpson N, et al; DIAGRAM+ Consortium; MAGIC Consortium; GLGC Investigators; MuTHER Consortium; DIAGRAM Consortium; GIANT Consortium; Global B Pgen Consortium; Procardis Consortium; MAGIC investigators; GLGC Consortium. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet*. 2012;8(3):e1002607. doi:10.1371/journal.pgen.1002607
28. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol*. 2013;37(7):658-665. doi:10.1002/gepi.21758
29. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife*. 2018;7:e4408. doi:10.7554/eLife.34408
30. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol*. 2015;44(2):512-525. doi:10.1093/ije/dyv080
31. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data mendelian randomization via the zero modal pleiotropy assumption. *Int J Epidemiol*. 2017;46(6):1985-1998. doi:10.1093/ije/dyx102
32. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in mendelian randomization with some invalid instruments using a weighted median estimator. *Genet Epidemiol*. 2016;40(4):304-314. doi:10.1002/gepi.21965
33. Choi KW, Chen CY, Stein MB, et al; Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium. Assessment of bidirectional relationships between physical activity and depression among adults: a 2-sample mendelian randomization study. *JAMA Psychiatry*. 2019;76(4):398-408.
34. Gage SH, Jones HJ, Burgess S, et al. Assessing causality in associations between cannabis use and schizophrenia risk: a two-sample mendelian randomization study. *Psychol Med*. 2017;47(5):971-980. doi:10.1017/S0033291716003172
35. Hartwig FP, Borges MC, Horta BL, Bowden J, Davey Smith G. Inflammatory biomarkers and risk of schizophrenia: a 2-sample mendelian randomization study. *JAMA Psychiatry*. 2017;74(12):1226-1233. doi:10.1001/jamapsychiatry.2017.3191
36. Zhao Q, Wang J, Hemani G, Bowden J, Small DS. Statistical inference in two-sample summary-data mendelian randomization using robust adjusted profile score. *arXiv*. 2018. <https://arxiv.org/abs/1801.09652>. Accessed March 25, 2019.
37. Zhang Q, Yang C, Wang J, Small DS. Powerful genome-wide design and robust statistical inference in two-sample summary-data mendelian randomization. *arXiv*. 2018. <https://arxiv.org/abs/1804.07371>. Accessed March 25, 2019.
38. Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in mendelian randomization studies. *Hum Mol Genet*. 2018;27(R2):R195-R208. doi:10.1093/hmg/ddy163
39. Jordan DM, Verbanck M, Do R. The landscape of pervasive horizontal pleiotropy in human genetic variation is driven by extreme polygenicity of human traits and diseases [published online April 30, 2018]. *bioRxiv*. doi:10.1101/311332
40. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from mendelian randomization between complex traits and diseases. *Nat Genet*. 2018;50(5):693-698. doi:10.1038/s41588-018-0099-7
41. Burgess S, Thompson SG. Multivariable mendelian randomization: the use of pleiotropic genetic variants to estimate causal effects. *Am J Epidemiol*. 2015;181(4):251-260. doi:10.1093/aje/kwu283

42. Sanderson E, Davey Smith G, Windmeijer F, Bowden J. An examination of multivariable mendelian randomization in the single-sample and two-sample summary data settings [published online December 10, 2018]. *Int J Epidemiol*. doi:10.1093/ije/dyy262
43. Yavorska OO, Burgess S. MendelianRandomization: an R package for performing mendelian randomization analyses using summarized data. *Int J Epidemiol*. 2017;46(6):1734-1739. doi:10.1093/ije/dyx034
44. Battle A, Brown CD, Engelhardt BE, Montgomery SB; GTEx Consortium; Laboratory, Data Analysis & Coordinating Center (LDACC)—Analysis Working Group; Statistical Methods groups—Analysis Working Group; Enhancing GTEx (eGTEx) groups; NIH Common Fund; NIH/NCI; NIH/NHGRI; NIH/NIMH; NIH/NIDA; Biospecimen Collection Source Site—NDRI; Biospecimen Collection Source Site—RPCI; Biospecimen Core Resource—VARI; Brain Bank Repository—University of Miami Brain Endowment Bank; Leidos Biomedical—Project Management; ELSI Study; Genome Browser Data Integration & Visualization—EBI; Genome Browser Data Integration & Visualization—UCSC Genomics Institute, University of California Santa Cruz; Lead analysts; Laboratory, Data Analysis & Coordinating Center (LDACC); NIH program management; Biospecimen collection; Pathology; eQTL manuscript working group. Genetic effects on gene expression across human tissues. *Nature*. 2017;550(7675):204-213. doi:10.1038/nature24277
45. Darmanis S, Sloan SA, Zhang Y, et al. A survey of human brain transcriptome diversity at the single cell level. *Proc Natl Acad Sci U S A*. 2015;112(23):7285-7290. doi:10.1073/pnas.1507125112
46. La Manno G, Gyllborg D, Codeluppi S, et al. Molecular diversity of midbrain development in mouse, human, and stem cells. *Cell*. 2016;167(2):566-580.e19.
47. Zhong S, Zhang S, Fan X, et al. A single-cell RNA-seq survey of the developmental landscape of the human prefrontal cortex. *Nature*. 2018;555(7697):524-528. doi:10.1038/nature25980
48. Habib N, Avraham-Davidi I, Basu A, et al. Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nat Methods*. 2017;14(10):955-958. doi:10.1038/nmeth.4407
49. de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol*. 2015;11(4):e1004219. doi:10.1371/journal.pcbi.1004219
50. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun*. 2017;8(1):1826. doi:10.1038/s41467-017-01261-5
51. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLoS Genet*. 2017;13(11):e1007081. doi:10.1371/journal.pgen.1007081
52. Robinson MR, Kleinman A, Graff M, et al. Genetic evidence of assortative mating in humans. *Nature Human Behaviour*. 2017;1(1). <https://www.nature.com/articles/s41562-016-0016>. Accessed April 1, 2019.
53. Betancourt TS, Newnham EA, Birman D, Lee R, Ellis BH, Layne CM. Comparing trauma exposure, mental health needs, and service utilization across clinical samples of refugee, immigrant, and US-origin children. *J Trauma Stress*. 2017;30(3):209-218. doi:10.1002/jts.22186
54. James LM, Strom TQ, Leskela J. Risk-taking behaviors and impulsivity among veterans with and without PTSD and mild TBI. *Mil Med*. 2014;179(4):357-363. doi:10.7205/MILMED-D-13-00241
55. Lowe SR, Galea S, Uddin M, Koenen KC. Trajectories of posttraumatic stress among urban residents. *Am J Community Psychol*. 2014;53(1-2):159-172. doi:10.1007/s10464-014-9634-6
56. Abdellaoui A, Hugh-Jones D, Kemper KE, et al. Genetic consequences of social stratification in Great Britain [published online October 30, 2018]. *bioRxiv*. doi:10.1101/457515
57. Kong A, Thorleifsson G, Frigge ML, et al. The nature of nurture: effects of parental genotypes. *Science*. 2018; 359(6374):424-428. doi:10.1126/science.aan6877
58. Trejo S, Domingue BW. Genetic nature or genetic nurture? quantifying bias in analyses using polygenic scores [published online January 18, 2019]. *bioRxiv*. doi:10.1101/524850
59. Pietrzak RH, Feder A, Singh R, et al. Trajectories of PTSD risk and resilience in World Trade Center responders: an 8-year prospective cohort study. *Psychol Med*. 2014;44(1):205-219. doi:10.1017/S0033291713000597
60. Martschenko D, Trejo S, Domingue BW. Genetics and education: recent developments in the context of an ugly history and an uncertain future. *AERA Open*. 2019;5(1). doi:10.1177/2332858418810516

SUPPLEMENT.

eAppendix. Phenotype Definitions, Sample Overlap, Palindromic Variants, and Assortative Mating

eReferences.

eTable 1. Traumatic Experiences Assessed in the UK Biobank

eTable 2. Results of the Sensitivity Analyses Conducted With Respect to the MathClass→PTSD Test With and Without the Outlier Variants in the MathClass Genetic Instrument

- eTable 3.** Results of the IVW Analyses Considering Genetic Instruments With and Without Palindromic Variants With Ambiguous Allele Frequencies (PAL and noPAL, Respectively)
- eTable 4.** MR-RAPS Analysis Considering Various Adjustments Based on Genome-Wide Genetic Instruments
- eTable 5.** Results of the Sensitivity Analyses Conducted With Respect to the Income→PTSD and Risk-Tak→PTSD Tests
- eTable 6.** Genetic Correlation Among Trauma Experiences Assessed in the UK Biobank
- eTable 7.** Results (Causal Effects and Sensitivity Analyses) of the MR Test Conducted Using Trauma-Related Genetic Instruments With Respect to PTSD
- eTable 8.** Results of the Enrichment Analysis Based on Tissue-Specific and Cell Type-Specific Transcriptomic Data
- eFigure 1.** Schematic Workflow of the Analyses Conducted
- eFigure 2.** Genetic Correlations Estimated Between Traits Related to Cognitive Ability and 2 Versions of the Posttraumatic Stress Disorder Data Set, PGC-PTSD Freeze-2 (2) and PGC-PTSD Freeze-1.5 (1.5)
- eFigure 3.** Effect of the PTSD PRS on Educational Attainment (Yellow) and Cognitive Performance (Green) Considering Different Inclusion Thresholds
- eFigure 4.** Leave-1-Out Analysis Conducted With Respect to the MathClass→PTSD2 Result
- eFigure 5.** Identification of Potential Outliers (in Red) in MathClass Genetic Instrument Based on IVW Heterogeneity Test and MR-RAPS Standardized Residuals
- eFigure 6.** Results of the MathClass→PTSD2 Analysis After the Removal of the Potential Outliers From the Genetic Instrument
- eFigure 7.** Results of the Sensitivity Analyses With Respect to All MR Analyses Conducted
- eFigure 8.** Multivariable Mendelian Randomization Analysis Considering the Effects of EdAtt and MathClass on PTSD
- eFigure 9.** Effect of Income and RiskTak PRS (Green and Blue, Respectively) on PTSD Considering Different Inclusion Thresholds
- eFigure 10.** Genetic Correlations Between Traumatic Experiences and the Other Traits of Interest
- eFigure 11.** Effect of Trauma-Related PRS on PTSD Considering Different Inclusion Thresholds
- eFigure 12.** Multivariable Mendelian Randomization Analysis Considering the Effects of Physically Abused by Family as a Child and Belittlement by Partner or Ex-Partner as an Adult on PTSD