

Genome-Wide Association Studies in Alzheimer Disease

Stephen C. Waring, DVM, PhD; Roger N. Rosenberg, MD

The genetics of Alzheimer disease (AD) to date support an age-dependent dichotomous model whereby earlier age of disease onset (< 60 years) is explained by 3 fully penetrant genes (*APP* [NCBI Entrez gene 351], *PSEN1* [NCBI Entrez gene 5663], and *PSEN2* [NCBI Entrez gene 5664]), whereas later age of disease onset (≥ 65 years) representing most cases of AD has yet to be explained by a purely genetic model. The *APOE* gene (NCBI Entrez gene 348) is the strongest genetic risk factor for later onset, although it is neither sufficient nor necessary to explain all occurrences of disease. Numerous putative genetic risk alleles and genetic variants have been reported. Although all have relevance to biological mechanisms that may be associated with AD pathogenesis, they await replication in large representative populations. Genome-wide association studies have emerged as an increasingly effective tool for identifying genetic contributions to complex diseases and represent the next frontier for furthering our understanding of the underlying etiologic, biological, and pathologic mechanisms associated with chronic complex disorders. There have already been success stories for diseases such as macular degeneration and diabetes mellitus. Whether this will hold true for a genetically complex and heterogeneous disease such as AD is not known, although early reports are encouraging. This review considers recent publications from studies that have successfully applied genome-wide association methods to investigations of AD by taking advantage of the currently available high-throughput arrays, bioinformatics, and software advances. The inherent strengths, limitations, and challenges associated with study design issues in the context of AD are presented herein.

Arch Neurol. 2008;65(3):329-334

Alzheimer disease (AD) is the most common cause of dementia and the most prevalent neurodegenerative disorder associated with aging.¹ Alzheimer disease is

For editorial comment see page 307

a heterogeneous disorder with a complex etiology owing to genetic and environmental influences as causal or risk modifiers. The neuropathologic hallmarks of

disease are extracellular amyloid plaques and intracellular neurofibrillary tangles of hyperphosphorylated tau protein.² Only 10% of AD cases occurring before 60 years of age (early-onset AD) are due to rare, fully penetrant (autosomal dominant) mutations in 3 genes: $A\beta$ precursor protein (*APP*) on chromosome 21,³ presenilin 1 (*PSEN1*) on chromosome 14,⁴ and presenilin 2 (*PSEN2*) on chromosome 1.^{5,6} In contrast, most cases of AD are later in onset (≥ 65 years of age) (late-onset AD), are nonfamilial, and are likely the result of highly prevalent genetic variants with low penetrance.⁷ To date, the only genetic risk factor for late-onset AD remains the apolipoprotein E gene (*APOE*), specifically the $\epsilon 4$ allele, which is moderately penetrant, accounting for up to 50% of cases.⁸

Author Affiliations: Department of Epidemiology, The University of Texas School of Public Health, The University of Texas Health Science Center at Houston (Dr Waring); Department of Neurology, The University of Texas Southwestern Medical Center at Dallas (Dr Rosenberg); and Texas Alzheimer's Research Consortium (University of Texas Southwestern Medical Center, Dallas; University of North Texas Health Science Center, Fort Worth; Texas Tech University Health Science Center, Lubbock; and Baylor College of Medicine, Houston) (Drs Waring and Rosenberg).

Table 1. Genetics of AD

Putative Genetic Variants (Selected Individual Source)	Chromosome	Consequences
<i>DAPK1</i> (Li et al, ¹² 2006)	9q34.1	Neuronal apoptosis
<i>UBQLN</i> (Bertram et al, ¹³ 2005)	9q21.32	Modulates accumulation of presenilin proteins
<i>ABCA1</i> (Sundar et al, ¹⁴ 2007)	9q22-q31	Possible role in lipid regulation
<i>SORL1</i> (Rogaeva et al, ⁹ 2007)	11q23	Involved with APP trafficking
<i>GAB2</i> (Li et al, ¹¹ 2004)	11q14.1	Scaffolding protein possibly affecting tau, amyloid, and other AD-related pathologic mechanism
<i>LRP6</i> (De Ferrari et al, ¹⁵ 2007)	12q13-q14	Altered Wnt/ β -catenin signaling

Abbreviations: AD, Alzheimer disease; APP, amyloid precursor protein.

However, a robust literature reports numerous putative genetic risk alleles and promising genetic variants. Recent reports from individual studies reveal significant associations with the sortilin-related receptor (*SORL1* [NCBI Entrez gene 6653])^{9,10} and glycine-rich protein 2-associated binding protein 2 (*GAB2* [NCBI Entrez gene 9846])¹¹ on chromosome 11; death-associated protein kinase 1 (*DAPK1* [NCBI Entrez gene 1612]),¹² ubiquilin 1 (*UBQLN1* [NCBI Entrez gene 299798]),¹³ and adenosine triphosphate-binding cassette transporter 1, subfamily A (*ABCA1* [NCBI Entrez gene 19]), on chromosome 9¹⁴; and low-density lipoprotein receptor-related protein 6 (*LRP6* [NCBI Entrez gene 4040]) on chromosome 12.¹⁵ All of these putative variants still lack replication in large representative populations but have relevance to neuropathologic mechanisms and pathways that may be associated with AD pathogenesis (**Table 1**).

A large meta-analysis from the AlzGene database¹⁶ representing 1055 polymorphisms and 355 genes reported in the literature as of August 2006 revealed the following 13 additional potential AD-susceptibility genes: angiotensin I converting enzyme (*ACE* [NCBI Entrez gene 1636]); cholinergic receptor, nicotinic, beta 2 (*CHRNA2* [NCBI Entrez gene 1141]); cystatin C (*CST3* [NCBI Entrez gene 1471]); estrogen receptor 1 (*ESR1* [NCBI Entrez gene 2099]); glyceraldehyde-3-phosphate dehydrogenase, spermatogenic (*GAPDH* [NCBI Entrez gene 26330]); insulin-degrading enzyme (*IDE* [NCBI Entrez gene 3416]); 5,10-methylenetetrahydrofolate reductase (*MTHFR* [NCBI Entrez gene 4524]); nicastrin (*NCSTN* [NCBI Entrez gene 23385]); prion protein (*PRNP* [NCBI Entrez gene 5621]); presenilin 1 (*PSEN1*); transferrin (*TF* [NCBI Entrez gene 7018]); mitochondrial transcription factor A (*TFAM* [NCBI Entrez gene 7019]); and tumor necrosis factor (*TNF* [NCBI Entrez gene 7124]).¹⁷ All are associated with relevant biological mechanisms and pathways but await replication to further elucidate their utility as significant markers for AD.

Understanding the role that genetic defects play in the pathogenesis of AD has been a major focus of investigation for several years. These collective efforts have provided valuable insights into the genetic and molecular mechanisms associated with AD. Until recently, most reports have been from linkage analyses and studies that have examined the association of single-nucleotide polymorphisms (SNPs), usually in a single candidate gene.¹⁶ However, the completion of the Human Genome Project,¹⁸ along with the development of public databases to cata-

log variation between individuals (SNPs) and advances in high-throughput, high-density genotyping technology have led to a sharp increase in the number of studies now examining a large number of SNPs simultaneously in hypothesis-independent designs. Indeed, genome-wide association studies (GWASs) have emerged as an increasingly effective tool for identifying genetic contributions to complex diseases and represent the next frontier for furthering our understanding of the underlying etiologic, biological, and pathologic mechanisms associated with chronic complex disorders. There have been early success stories for diseases such as macular degeneration¹⁹⁻²¹ and diabetes mellitus.^{22,23} Whether this will hold true for a genetically complex and heterogeneous disease such as AD is not known, although indications from early reports are encouraging.

This review considers recent publications from studies that have successfully applied genome-wide association methods to investigations of AD by taking advantage of the currently available high-throughput arrays, bioinformatics, and software advances. The inherent strengths, limitations, and challenges associated with study design issues are presented herein to help guide future GWASs in AD.

GWASs IN AD

The recent decline in the cost of genotyping coupled with technological advances that now allow simultaneous examination of up to 1 000 000 SNPs in a single high-throughput assay has led to an explosion of GWASs of common chronic diseases, including AD. Initial studies focusing on AD have taken advantage of already amassed robust data from case-control and longitudinal cohorts with reliable information on the phenotype of interest (eg, age at onset, diagnosis, and cognitive profile) and DNA available for genotyping²⁴⁻²⁷ (**Table 2**).

A GWAS of neuropathologically confirmed AD cases and control subjects from the United States and the Netherlands revealed a significant SNP (rs4420638) in linkage disequilibrium to the *APOE* ϵ 4 variant on chromosome 19, thus providing support for the *APOE* locus as the major susceptibility gene for late-onset AD, with an odds ratio significantly greater than that for any other locus in the human genome.²⁴ That study also supported the feasibility of using the most recently available ultrahigh-density GWAS (502 627 SNPs) for AD and other heritable phenotypes. A subsequent study on the

Table 2. High-Density Genome-Wide Association Studies on AD

Source	Population	Total Markers (SNPs) Examined	Significant Markers	Relevance
Coon et al, ²⁴ 2007	Neuropathologically confirmed AD (n=664) or healthy (n=422) from United States and the Netherlands)	502 627 (Affy 500k) ^a	rs4420638 on chromosome 19 in LD to <i>APOE</i> ϵ 4 variant	Confirms strength of association of <i>APOE</i> and development of AD; would have been missed except for 1 SNP; importance of increased SNP density for more complete coverage
Reiman et al, ²⁵ 2007	Total: 1146 subjects from the United States and the Netherlands Discovery: neuropathologically confirmed cases of AD (n=446) and controls (n=290) Replication: neuropathologic sample of 197 cases of AD and 114 controls; clinical sample of 218 cases of late-onset AD and 146 controls	312 316 (Affy 500K) ^a	rs2373115 (<i>GAB2</i>) on chromosome 11q14.1	rs2373115 genotype GG inferred significantly higher risk (OR, 2.36) of AD among <i>APOE</i> ϵ 4 carriers compared with genotypes GT or TT but not among <i>APOE</i> ϵ 4 noncarriers; <i>GAB2</i> modifies risk of AD in <i>APOE</i> ϵ 4 carriers
Liu et al, ²⁷ 2007	103 Patients with late-onset AD and 170 first-degree relatives from pedigree of 4645 members in the Netherlands	262 000 (Affy 250 Nsp arrays of Affy 500k) ^a	Chromosomes 1q25 (<i>RGSL2</i> , <i>RALGPS2</i> , and <i>C1orf49</i>), 3q22-24 (<i>NMNAT3</i> and <i>CLSTN2</i>), and 10Q22-24 (<i>HTR7</i> , <i>MPHOSPH1</i> , and <i>CYP2C</i>)	Confirmatory for previous linkage to 1q21-25 and 10q22-24; with cognitive function as endophenotype, authors conclude that <i>RGSL2</i> on chromosome 1 and <i>NMNAT3</i> on chromosome 3 likely to be the relevant genes; first to show significant linkage to 3q23 markers
Grube et al, ²⁶ 2007	5 Case-control sample sets from the United Kingdom and the United States, including cases of late-onset AD (n=1808) and healthy controls (n=2062)	17 343 (Celera cSNP) ^b	3 SNPs on chromosome 19 in LD to <i>APOE</i> gene; rs3745833 (<i>GALP</i>) on chromosome 19q13.42; rs1554948 (<i>TNK1</i>) on chromosome 17p13.1; rs8192708 (<i>PCK1</i>) on chromosome 20q13.31	Three most significant SNPs in LD to <i>APOE</i> lends to validity of design; galaninlike peptide precursor (<i>GALP</i>); nonreceptor tyrosine kinase (<i>TNK1</i>) and phosphoenolpyruvate carboxykinase (<i>PCK1</i>) show direct or indirect effect on metabolic, trafficking, or signaling pathways that may have relevance to AD

Abbreviations: AD, Alzheimer disease; LD, linkage disequilibrium; OR, odds ratio; SNP, single-nucleotide polymorphism.

^aDetermined using Affymetrix GeneChip Human Mapping 500K Array set (Affymetrix, Inc, Santa Clara, California).

^bAvailable from Celera Diagnostics (Alameda, California).

same population enriched with clinical samples divided into *APOE* ϵ 4 carriers and noncarriers for discovery and replication demonstrated a modifying effect on the risk of AD associated with the *GAB2* among *APOE* ϵ 4 carriers but not among noncarriers.²⁵

The first GWAS to appear in the literature, and which focused on putative functional variants (SNPs) in AD, examined 17 343 markers in a robust sample of 1808 late-onset cases of AD and 2062 controls from the United States and the United Kingdom. Results revealed 19 significant markers associated with the risk of AD. Three of these were galaninlike peptide precursor (*GALP* [NCBI Entrez gene 85569]) on chromosome 19q13.42, nonreceptor tyrosine kinase (*TNK1* [NCBI Entrez gene 8711]) on chromosome 17p13.1, and phosphoenolpyruvate carboxykinase (*PCK1* [NCBI Entrez gene 5105]) on chromosome 20q13.31, all with potential relevance to AD.²⁶ The fact that 3 of the 19 SNPs were in linkage disequi-

librium to the *APOE* ϵ 4 locus lends validity to the study design. However, replication in large appropriately designed studies is necessary for confirmation and further elucidation of their relevance in AD pathogenesis.

A GWAS from a genetically isolated population in the Netherlands of 103 patients with late-onset AD and 170 first-degree relatives from a complex pedigree of 4645 members focused on genetic variants associated with cognitive endophenotypes in addition to risk of AD.²⁷ This was the first study to show significant linkage to 3q23 markers associated with AD. Findings also confirmed previous linkage to 1q25, 10q22-24, and 11q25 associated with AD. Using cognitive function as the endophenotype, the authors²⁷ concluded that *RGSL2* (NCBI Entrez gene 84227), *RALGPS2* (NCBI Entrez gene 55103), and *C1orf49* (NCBI Entrez gene 84066) on chromosome 1; *HTR7* (NCBI Entrez gene 3363), *MPHOSPH1* (NCBI Entrez gene 9585), and *CYP2C* (NCBI Entrez gene 1559)

on chromosome 10; and *NMNAT3* (NCBI Entrez gene 349565) and *CLSTN2* (NCBI Entrez gene 64084) on chromosome 3 are likely to be the relevant genes. They failed to confirm *SORL1* on chromosome 11 but reported *OPCMS* (NCBI Entrez gene 4978) and *HNT* (NCBI Entrez gene 50863) to be the relevant genes. Another GWAS of endophenotypes with relevance to AD from the Framingham group focused on changes in brain volumetric measures on magnetic resonance imaging and cognitive performance in 705 individuals free of dementia and stroke.²⁸ They found significant associations between *SORL1* (rs1131497) and abstract reasoning and between *CDH4* (rs1970546) and total cerebral brain volume. They also reported polymorphisms with 28 of 163 candidate genes for AD, stroke, and memory impairment for the endophenotypes studied. Overall, these results indicate that genetic variants may be detectable with subclinical phenotypes known to increase the risk of developing clinical neurological disease.²⁸

Although these studies represent an excellent start, with several new associations already detected within the past year, they constitute only the beginning, as many of these putative associations await validation in sufficiently large and representative populations before their utility as genetic markers for disease is completely understood. An important consideration is that the results thus far show very modest effect sizes in the order of relative risks (or odds ratios) of less than 2.0, and most in the range of 1.1 to 1.5. As expected with most common complex diseases with onset later in life,²⁹ late-onset AD likely results from genetic variants with modest effect size that, while statistically significant, are neither sufficient nor necessary to explain all occurrences of disease.³⁰ It is therefore incumbent that the synthesis and interpretation of weak but consistent associations that are biologically plausible afford distinction from spurious associations that may result from bias and inferior study design.

GWAS DESIGN CONSIDERATIONS IN AD

Sample Size and Power

One of the most important and challenging steps in designing large-scale GWASs on currently available high-density genotyping platforms is assembling sufficiently large study populations that will allow robust gene discovery and replication. Sample size and power calculations for GWASs are statistically rigorous and beyond the scope of this review. Interested readers are referred to recent publications on the subject that take into account the density and coverage of newer available genotyping platforms and software.^{31,32} These reports and others point to the importance of issues that must be taken into account when planning a study.

Power to detect a causal variant increases with sample size and magnitude of effect (relative risk or odds ratio) and can vary by genotyping platform and the population studied. The effect of stratification on *APOE* and the effect of population stratification need to be taken into account. Owing to the large number of genetic variants (SNPs) examined and the lack of independence with SNPs in linkage disequilibrium to each other, multiple testing

and type I error are even greater threats to validity for the modest effect sizes expected. In addition, power calculations should take into account misclassification error, not just of the genotype but of clinical misclassification error associated with the phenotype, particularly when the trait of interest is disease status in the case of AD (as discussed in the “Phenotype” subsection).

To overcome these inherent issues, large sample sizes of well-characterized phenotypes and tiered strategies for discovery and replication have proved to be the most successful strategy for studies appearing in the recent literature. Most studies, such as those funded through the Genetic Association Information Network³³ and the Wellcome Trust Case Control Consortium,³⁴ have included 2000 to more than 10 000 cases and controls to identify truly significant results. Considering the current cost of genotyping large-scale samples and the availability of large numbers of well-characterized research subjects, some of which have already been included in GWASs, a similar approach is both timely and feasible for AD, albeit not without challenges and limitations.

Phenotype

Although the initial analyses for most GWASs of AD may use disease status as the phenotypic trait of interest, testing for associations with secondary traits such as clinical and cognitive features, biomarkers including neuroimaging, and neuropathological features has considerable merit. Because of established criteria for a diagnosis of AD and the need for standardization across sites for large multicenter studies, considerable effort goes into making a clinical diagnosis of AD. However, establishing an appropriate control continues to be a challenge. The presence of AD neuropathologic changes in brains of nondemented individuals at autopsy is not uncommon, and a number of patients with a clinical diagnosis of probable AD also have cerebrovascular comorbidities that may influence cognition. This has important implications for designing GWASs, because most genetic defects will be associated with neurobiological and biochemical processes that lead directly to clinical AD through neurodegenerative changes (eg, formation of neurofibrillary tangles and amyloid deposition) or indirectly through other mechanisms (eg, cerebrovascular disease). Indeed, the potential for biological heterogeneity among individuals with clinical AD and among nondemented controls presents a challenge in the design of a GWAS and in the interpretation of the findings. Therefore, strategies to minimize the effects of misclassification error associated with the phenotype of interest need to be developed for implementation in future large-scale studies.

Discovery and Replication Samples

Ideally, a GWAS for AD would begin with a discovery sample that is sufficiently large with a wealth of quantitative and qualitative information available for phenotypic structuring. Another large sample with similar phenotypic structure would then be used to further narrow the most significant SNPs resulting from discovery genotyping to identify a much smaller number of significant

SNPs. Those SNPs would then be examined in a larger sample with similar phenotypic structure for replication (**Figure**). Finally, the most significant SNPs identified would then need to be subjected to further analyses (gene expression, sequencing, fine mapping, and functional studies) for final confirmation of relevant putative causal variants. Guidelines for replication of genotype-phenotype associations and GWASs have been published recently.^{35,36}

Although the numbers are arbitrary, because of the number of traits to be examined and the modest effect size expected for most traits, it is clear that multiple large samples for discovery and replication are necessary to ensure a successful GWAS.^{37,38} This will require forming consortia that will afford pooling across studies because it is unlikely that a single study will have adequate power to unequivocally identify novel genetic risk factors for AD. To take advantage of available data on a large number of patients with AD and controls, plans to form consortia such as the National Institute on Aging (NIA) Genome-Wide Association Study Consortium are already under way. These collective efforts will attempt to design and complete GWASs in the near future to provide an excellent and timely foundation for numerous studies to follow. These initial efforts will likely include GWASs already completed or in progress such as the Translational Genomics Research Initiative study of AD, the National Institute of Mental Health AD sample, the Framingham Study, and the Texas Alzheimer's Research Consortium among other publicly funded and public-private initiatives in progress or planned. Valuable resources such as the National Cell Repository of AD, the NIA National Alzheimer Coordinating Center, and ongoing studies such as the Adult Changes in Thought, the Washington Heights–Inwood Columbia Aging Project, and the Chicago Healthy Aging Project should also be considered, along with ongoing biomarker and neuroimaging studies such as the Alzheimer Disease Neuroimaging Initiative. All of these resources would be expected to have readily available DNA from a large sample of well-characterized research participants and could provide a robust pooled sample that is immediately available to initiate a large-scale GWAS.

The potential contribution from large family studies such as the NIA Late Onset Alzheimer Disease centers, particularly for discovery, should also be evaluated. Although the “common disease–common variant” hypothesis³⁹ provided the conceptual framework for the HapMap Project on which the commercially available high-density genotyping platforms are based, multiple rare variants would not be identified by the tagging SNPs. However, their role in susceptibility to common complex diseases may be significant.^{40,41}

This discussion focuses on replication-based multistage design analysis. However, some groups recommend joint analysis as an alternative strategy for GWASs, whereby data from both stages are analyzed together. This strategy may be even more efficient and achieve greater power when a large proportion of the study sample is genotyped in stage 1 and a relatively large proportion of markers is selected for stage 2. An excellent review comparing joint analysis with replication-based analysis is provided by Skol et al.⁴² The choice of analysis for multistage design is an impor-




Figure. Hypothetical tiered design strategy for a large-scale genome-wide association study of Alzheimer disease (AD). SNP indicates single-nucleotide polymorphism.

tant study design issue that must be taken into account to minimize genotyping cost and maximize statistical power within the context of the available sample size.

CHALLENGES

Assembling large representative samples for discovery and replication is certainly not without challenges. The extent to which individual institutional review board–approved protocols allow sharing of banked DNA and genetic information must be verified. Logistical and technical issues associated with locating and assembling large DNA sample sets and subsequent genotyping is a large and expensive undertaking. Cataloguing all available data required to support the various endophenotypes to be tested initially and transforming these data into a standardized format for entry into a unified database require considerable planning, technical support, and funding. As evidenced by the success of studies already completed, these initial large-scale studies are expected to produce a number of candidate polymorphisms and haplotypes that will require replication and confirmation in other samples. Expanding current studies now to include comparably collected information on anticipated significant endophenotypes and collection of biospecimens and neuroimaging will require tremendous resources and planning. Research teams populated with subject matter experts representing the entire spectrum of expertise required must be in place to ensure success of such an undertaking over the next 5 years and beyond and therefore becomes a matter of funding and prioritization of research agendas among all respective stakeholders.

CONCLUSIONS

Although there are challenges to assembling a large representative population for discovery and replication studies, the prospect of new insights that will further our understanding of the role of genetic defects in the neuropathogenic mechanisms that lead to AD make the GWAS an attractive and powerful tool worth the effort and investment. However, there is tremendous work yet to be done, not only in designing appropriately large and representative studies that will lead to numerous new gene discoveries but also in translating these discoveries to ad-

vance clinical practice. Indeed, the time has come to leverage the wealth of advancing methods in genetics, neuroscience, bioinformatics, statistics, and genetic epidemiology that will lead to better treatment outcomes, drug discovery, and prognosis for the millions of individuals already affected by or at risk of developing AD.

Accepted for Publication: November 24, 2007.

Correspondence: Stephen C. Waring, DVM, PhD, Department of Epidemiology, The University of Texas School of Public Health, 1200 Herman Pressler, RAS-E629, Houston, TX 77030 (stephen.c.waring@uth.tmc.edu)

Author Contributions: *Study concept and design:* Waring and Rosenberg. *Drafting of the manuscript:* Waring. *Critical revision of the manuscript for important intellectual content:* Waring and Rosenberg.

Financial Disclosure: None reported.

Funding/Support: This study was supported by P30AG12300-14 from the NIA, National Institutes of Health (Alzheimer's Disease Center, The University of Texas Southwestern Medical Center [Dr Rosenberg]), and by funds from the state of Texas through the Texas Council on Alzheimer's Disease and Related Disorders (Drs Waring and Rosenberg).

Additional Contributions: The NIA Genetics of Alzheimer's Disease Workgroup discussants provided some of the conceptual framework presented herein.

REFERENCES

- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics [published correction appears in *Science*. 2002;297(5590):2209]. *Science*. 2002;297(5580):353-356.
- Rosenberg CK, Pericak-Vance MA, Saunders AM, Gilbert JR, Gaskell PC, Hulette CM. Lewy body and Alzheimer pathology in a family with the amyloid-beta precursor protein APP717 gene mutation. *Acta Neuropathol*. 2000;100(2):145-152.
- Goate A, Chartier-Harlin MC, Mullan M, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature*. 1991;349(6311):704-706.
- Sherrington R, Rogaev EI, Liang Y, et al. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*. 1995;375(6534):754-760.
- Levy-Lahad E, Wasco W, Poorkaj P, et al. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science*. 1995;269(5226):973-977.
- Rogaev EI, Sherrington R, Rogaeva EA, et al. Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature*. 1995;376(6543):775-778.
- Tanzi RE. A genetic dichotomy model for the inheritance of Alzheimer's disease and common age-related disorders. *J Clin Invest*. 1999;104(9):1175-1179.
- Strittmatter WJ, Saunders AM, Schmechel D, et al. Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci U S A*. 1993;90(5):1977-1981.
- Rogaeva E, Meng Y, Lee JH, et al. The neuronal sortilin-related receptor SORL1 is genetically associated with Alzheimer disease. *Nat Genet*. 2007;39(2):168-177.
- Lee JH, Cheng R, Schupf N, et al. The association between genetic variants in SORL1 and Alzheimer disease in an urban, multiethnic, community-based cohort. *Arch Neurol*. 2007;64(4):501-506.
- Li Y, Nowotny P, Holmans P, et al. Association of late-onset Alzheimer's disease with genetic variation in multiple members of the GAPD gene family [published correction appears in *Proc Natl Acad Sci U S A*. 2006;103(16):6411]. *Proc Natl Acad Sci U S A*. 2004;101(44):15688-15693.
- Li Y, Grupe A, Rowland C, et al. DAPK1 variants are associated with Alzheimer's disease and allele-specific expression. *Hum Mol Genet*. 2006;15(17):2560-2568.
- Bertram L, Hiltunen M, Parkinson M, et al. Family-based association between Alzheimer's disease and variants in UBQLN1. *N Engl J Med*. 2005;352(9):884-894.
- Sundar PD, Feingold E, Minster RL, DeKosky ST, Kambou MI. Gender-specific association of ATP-binding cassette transporter 1 (ABCA1) polymorphisms with the risk of late-onset Alzheimer's disease. *Neurobiol Aging*. 2007;28(6):856-862.
- De Ferrari GV, Papassotiropoulos A, Biechele T, et al. Common genetic variation within the low-density lipoprotein receptor-related protein 6 and late-onset Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2007;104(22):9434-9439.
- AlzGene: published AD candidate genes. Alzheimer Research Forum Web site. <http://www.alzforum.org/res/com/gen/alzgene/>. Accessed November 16, 2007.
- Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE. Systematic meta-analyses of Alzheimer disease genetic association studies: the AlzGene database. *Nat Genet*. 2007;39(1):17-23.
- International HapMap Consortium. A haplotype map of the human genome. *Nature*. 2005;437(7063):1299-1320.
- Edwards AO, Ritter R III, Abel KJ, Manning A, Panhuysen C, Farrer LA. Complement factor H polymorphism and age-related macular degeneration. *Science*. 2005;308(5720):421-424.
- Haines JL, Hauser MA, Schmidt S, et al. Complement factor H variant increases the risk of age-related macular degeneration. *Science*. 2005;308(5720):419-421.
- Klein RJ, Zeiss C, Chew EY, et al. Complement factor H polymorphism in age-related macular degeneration. *Science*. 2005;308(5720):385-389.
- Wellcome Trust Case Control Consortium. Genome-wide association study of 14 000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007;447(7145):661-678.
- Zeggini E, Weedon MN, Lindgren CM, et al. Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes [published correction appears in *Science*. 2007;317(5841):1035-1036]. *Science*. 2007;316(5829):1336-1341.
- Coon KD, Myers AJ, Craig DW, et al. A high-density whole-genome association study reveals that APOE is the major susceptibility gene for sporadic late-onset Alzheimer's disease. *J Clin Psychiatry*. 2007;68(4):613-618.
- Reiman EM, Webster JA, Myers AJ, et al. GAB2 alleles modify Alzheimer's risk in APOE epsilon4 carriers. *Neuron*. 2007;54(5):713-720.
- Grupe A, Abraham R, Li Y, et al. Evidence for novel susceptibility genes for late-onset Alzheimer's disease from a genome-wide association study of putative functional variants. *Hum Mol Genet*. 2007;16(8):865-873.
- Liu F, Arias-Vasquez A, Sleegers K, et al. A genomewide screen for late-onset Alzheimer disease in a genetically isolated Dutch population. *Am J Hum Genet*. 2007;81(1):17-31.
- Seshadri S, DeStefano AL, Au R, et al. Genetic correlates of brain aging on MRI and cognitive test measures. *BMC Med Genet*. 2007;8(suppl 1):S15.
- Wright A, Charlesworth B, Rudan I, Carothers A, Campbell H. A polygenic basis for late-onset disease. *Trends Genet*. 2003;19(2):97-106.
- Bertram L, Tanzi RE. The genetic epidemiology of neurodegenerative disease. *J Clin Invest*. 2005;115(6):1449-1457.
- Klein RJ. Power analysis for genome-wide association studies. *BMC Genet*. 2007;8(1):58.
- Pe'er I, de Bakker PI, Maller J, Yelensky R, Altshuler D, Daly MJ. Evaluating and improving power in whole-genome association studies using fixed marker sets. *Nat Genet*. 2006;38(6):663-667.
- Foundation for the National Institutes of Health. Genetic Association Information Network (GAIN). http://www.fnih.org/GAIN2/home_new.shtml. Accessed October 25, 2007.
- Wellcome Trust Case Control Consortium. WTCCC Web site. <http://www.wtccc.org.uk/>. Accessed October 25, 2007.
- Chanock SJ, Manolio T, Boehnke M, et al. Replicating genotype-phenotype associations. *Nature*. 2007;447(7145):655-660.
- Ioannidis JP, Boffetta P, Little J, et al. Assessment of cumulative evidence on genetic associations: interim guidelines [published online ahead of print September 26, 2007]. *Int J Epidemiol*. doi:10.1093/ije/dym159.
- Amos CI. Successful design and conduct of genome-wide association studies. *Hum Mol Genet*. 2007;16(special issue 2):R220-R225.
- Traynor BJ, Singleton A. Genome-wide association studies and ALS: are we there yet? *Lancet Neurol*. 2007;6(10):841-843.
- Lander ES. The new genomics: global views of biology. *Science*. 1996;274(5287):536-539.
- Barrett JC, Cardon LR. Evaluating coverage of genome-wide association studies. *Nat Genet*. 2006;38(6):659-662.
- Di Rienzo A. Population genetics models of common diseases. *Curr Opin Genet Dev*. 2006;16(6):630-636.
- Skol AD, Scott LJ, Abecasis GR, Boehnke M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet*. 2006;38(2):209-213.