


Original Investigation

Phenotypic Overlap Between Familial Exudative Vitreoretinopathy and Microcephaly, Lymphedema, and Chorioretinal Dysplasia Caused by *KIF11* Mutations

Johane M. Robitaille, MD; Roxanne M. Gillett, PhD; Marissa A. LeBlanc, PhD; Daniel Gaston, PhD; Mathew Nightingale, HND; Michael P. Mackley, BSc; Sandhya Parkash, MD; Julie Hathaway, MSc; Aidan Thomas, MSc; Anna Ellis, MD; Elias I. Traboulsi, MD; Elise Héon, MD; Mélanie Roy, MD; Stavit Shalev, MD; Conrad V. Fernandez, MD; Christine MacGillivray, BSc; Karin Wallace, BSc; Somayyeh Fahiminiya, PhD; Jacek Majewski, PhD; Christopher R. McMaster, PhD; Karen Bedard, PhD

 Supplemental content at jamaophthalmology.com

IMPORTANCE Retinal detachment with avascularity of the peripheral retina, typically associated with familial exudative vitreoretinopathy (FEVR), can result from mutations in *KIF11*, a gene recently identified to cause microcephaly, lymphedema, and chorioretinal dysplasia (MLCRD) as well as chorioretinal dysplasia, microcephaly, and mental retardation (CDMMR). Ophthalmologists should be aware of the range of presentations for mutations in *KIF11* because the phenotypic distinction between FEVR and MLCRD/CDMMR portends management implications in patients with these conditions.

OBJECTIVE To identify gene mutations in patients who present with a FEVR phenotype and explore the spectrum of ocular and systemic abnormalities caused by *KIF11* mutations in a cohort of patients with FEVR or microcephaly in conjunction with chorioretinopathy or FEVR.

DESIGN, SETTING, AND PARTICIPANTS Clinical data and DNA were collected from each participant between 1998 and 2013 from the clinical practices of ophthalmologists and clinical geneticists internationally. Twenty-eight FEVR probands with diagnoses made by the referring physician and without a known FEVR gene mutation, and 3 with microcephaly and chorioretinopathy, were included. At least 1 patient in each pedigree manifested 1 or more of the following: macular dragging, partial retinal detachment, falciform folds, or total retinal detachment.

EXPOSURES Whole-exome sequencing was conducted on affected members in multiplex pedigrees, and Sanger sequencing of the 22 exons of the *KIF11* gene was performed on singletons. Clinical data and history were collected and reviewed.

MAIN OUTCOMES AND MEASURES Identification of mutations in *KIF11*.

RESULTS Four novel heterozygous *KIF11* mutations and 1 previously published mutation were identified in probands with FEVR: p.A218Gfs*15, p.E470X, p.R221G, c.790-1G>T, and the previously described heterozygous p.R47X. Documentation of peripheral avascular areas on intravenous fluorescein angiography was possible in 2 probands with fibrovascular proliferation demonstrating phenotypic overlap with FEVR.

CONCLUSIONS AND RELEVANCE Mutations in *KIF11* cause a broader spectrum of ocular disease than previously reported, including retinal detachment. The *KIF11* gene likely plays a role in retinal vascular development and mutations in this gene can lead to clinical overlap with FEVR. Cases of FEVR should be carefully inspected for the presence of microcephaly as a marker for *KIF11*-related disease to enhance the accuracy of the prognosis and genetic counseling.

JAMA Ophthalmol. 2014;132(12):1393-1399. doi:10.1001/jamaophthalmol.2014.2814
Published online August 14, 2014.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Johane M. Robitaille, MD, IWK Health Centre, Eye Care Team, 5850/5980 University Ave, PO Box 9700, Halifax, NS B3K 6R8, Canada (jrobitai@dal.ca).

Microcephaly, lymphedema, and chorioretinal dysplasia (MLCRD), as well as chorioretinal dysplasia, microcephaly, and mental retardation (CDMMR), were recently reported by Ostergaard et al¹ to result from mutations in the *KIF11* gene (NCBI NM_004523), demonstrating that these conditions are allelic (OMIM 152950) with highly variable phenotypic expression. None of the patients with *KIF11* mutations in that study had retinal folds or microphthalmia. Of the 27 mutation carriers identified, unilateral total retinal detachment was present in only 1 individual.

Familial exudative vitreoretinopathy (FEVR) is a hereditary developmental disorder characterized by the failure of peripheral retinal vascularization at birth. Complications include partial and total retinal detachment, and the disease may present with retinal folds that mimic persistent fetal vasculature.² To date, 5 genes have been identified that account for approximately 50% of cases of FEVR: *NDP* (OMIM 300658),³ *FZD4* (OMIM 604579),⁴ *LRP5* (OMIM 603506),^{5,6} *TSPAN12* (OMIM 613138),^{7,8} and, more recently, *ZNF408* (NCBI 79797).⁹

We performed whole-exome sequencing on DNA from affected members of multiplex FEVR pedigrees with no known causative FEVR gene mutations and identified 1 pedigree with a heterozygous *KIF11* mutation (p.E470X). To test the hypothesis that more cases of FEVR are caused by *KIF11* mutations, we screened the *KIF11* gene in a cohort of FEVR probands who did not have a mutation in any of the known FEVR genes. We also screened the *KIF11* gene in typical and atypical cases of microcephaly with chorioretinopathy and/or retinal detachment. In addition, 2 of the 5 pedigrees with *KIF11* mutations were investigated further to identify a mechanism for the phenotypic overlap and differences between MLCRD and FEVR.

Methods

Participants and Clinical Data Collection

The study was approved by the research ethics board of the IWK Health Centre and written informed consent was obtained according to Canadian Tri-Council guidelines. No stipend was offered.

Participants were identified from a database of individuals in a study aiming to identify novel FEVR genes and describe the phenotypic spectrum of the disease. For recruitment in this database, the laboratory was listed on the GeneTests website¹⁰ and potential participants were recruited at the treating physician's request. Patients were included if they manifested signs compatible with one of the following diagnoses: FEVR, atypical/typical persistent fetal vasculature, Coats disease, and congenital retinal folds or detachment. Pedigrees with microcephaly and bilateral or unilateral chorioretinopathy were recruited separately for specific *KIF11* screening.

Clinical data were collected prospectively and retrospectively by the referring physician or the study investigator (J.M.R.). These data included results from eye examinations and, whenever possible, ultrasonography, electrophysiologic testing, intravenous fluorescein angiography (IVFA), and fun-

dus photography. Relatives at risk were also invited to participate by the referring physician or study investigator, and an eye examination was performed including best-corrected visual acuity, ocular alignment, slitlamp examination, dilated fundus examination, and in some cases, IVFA. Ethnicity was recorded for each participant by the referring physician who completed the study history questionnaire. This information was used in the analysis of novel mutation screening in a random population and in available databases.

Genetic Testing

Exome Capture and Sequencing, Read Mapping, and Variant Annotation

Blood and/or saliva samples were obtained from each participant for genomic DNA extraction using standard protocols. Whole-exome capture and sequencing were performed at the McGill University and Genome Quebec Innovation Centre as previously described.¹¹ In brief, a total of 3 µg of DNA was used for exome capture (SureSelect Human All Exon Kit, version 3; Agilent Technologies Inc). The captured DNA was sequenced with 100 base pair paired-end reads (HiSeq 2000 sequencer; Illumina). High-quality reads were mapped against the human genomic reference sequence (NCBI37/hg19) using the Burrows-Wheeler Aligner software.¹² Genomic variants were called using the Genome Analysis Toolkit pipeline¹³ and annotated with ANNOVAR.¹⁴

All variants were compared against the Cardiff University Human Gene Mutation Database,¹⁵ dbSNP,¹⁶ 1000 Genomes Project,¹⁷ Exome Variant Server,¹⁸ and a pool of more than 300 in-house exomes from unrelated projects. Variants considered potentially damaging included nonsynonymous mutations (missense and nonsense), splice-site variants, and frameshift changes due to short insertions and/or deletions (indels). Depending on the type of mutation discovered, novel mutations were evaluated for functional significance using various software programs including SIFT,¹⁹⁻²³ PolyPhen-2,²⁴ PROVEAN (version 1.13),^{25,26} CRYPP-SKIP,²⁷⁻³⁰ and Spliceman.^{31,32}

Sanger Sequencing and Mutation Identification

Direct automated Sanger sequencing was performed (ABI PRISM 3100 Genetic Analyzer; Applied Biosystems). Regions sequenced included the 2 coding exons of the *FZD4* gene, the 23 coding exons of the *LRP5* gene, the 3 exons of the *NDP* gene (including the 5' untranslated and promoter regions), the 8 coding exons of the *TSPAN12* gene, and the 22 coding exons of the *KIF11* gene (primer pairs available upon request). Because all *KIF11* mutations were detected before discovery of the *ZNF408* gene, *ZNF408* was not screened for variants in this study.

Mutation calling was performed (Mutation Surveyor, version 3.97; SoftGenetics LLC). Sequence changes were confirmed either in the forward and reverse direction or twice in the same direction. The variants were compared as stated above using the University of California, Santa Cruz, Genome Browser with the Human February 2009 (GRCh37/hg19) Assembly (<http://genome.ucsc.edu/>).³³ Novel missense mutations were screened using Sanger sequencing in 95 random population samples (180 chromosomes) of white race origin.

Results

Cohort Description and Mutation Detection Rate

Participants were recruited between April 1998 and September 2013. Seventy-two probands were enrolled to screen for a *FEVR* gene mutation because of ocular features compatible with a diagnosis of *FEVR*. Of these, 28 probands were found to not have a mutation in any of the following genes: *FZD4*, *LRP5*, *TSPAN12*, and *NDP*. A mutation in the *KIF11* gene was identified in 4 of 72 probands (5.6% mutation detection rate) with a phenotype mimicking *FEVR*. Two of these 4 probands were not reported to have microcephaly at the time of recruitment. Three of 6 probands with microcephaly associated with unilateral or bilateral chorioretinopathy and/or retinal folds or retinal detachment had a *KIF11* mutation. Clinical details for each participant are summarized in the eTable in the Supplement.

Genetic Analysis

Whole-exome sequencing was initially performed on an affected male and his affected sister (pedigree I) to identify a shared disease-causing variant. There were 267 shared missense or nonsense variants after filtering for suspected sequencing artifacts and with a multiple allele frequency greater than 5%. The siblings shared 3 truncating mutations and 1 frame shift deletion: the frame shift deletion was seen in 12 other sequenced exomes; 1 truncating mutation (*CORO7*) had been identified in an additional in-house exome and another was in the olfactory receptor gene *OR52N4*. The remaining heterozygous truncating mutation, *KIF11* p.E470X (c.1408 G>T), was selected as the candidate, because heterozygous truncating mutations in *KIF11* had previously been reported¹ as causative for *MLCRD*. There were no other siblings and no family history of a condition compatible with a diagnosis of *FEVR* or *MLCRD*. The mutation was absent in both parents, suggesting mosaicism in one of the parents.

Sanger sequencing identified additional novel heterozygous mutations in 3 probands (Figure 1): p.A218Gfs*15 (c.652DupG), p.R221G (c.661 A>G), and *KIF11* c.790-1G>T that alters an acceptor splice site. A previously reported³⁴ heterozygous mutation, *KIF11* p.R47X (c.139C>T), was found in a fifth proband. No participant harbored compound heterozygous mutations.

Clinical Description


The first mutation, *KIF11* p. E470X, identified in 2 siblings (1 male and 1 female) was associated with a *FEVR* phenotype and with microcephaly that was noted prior to enrollment. The family was recruited in the *FEVR* project specifically because of the fundus appearance mimicking *FEVR*. The proband was an 8-year-old boy at the time of recruitment. He had been noted to have intermittent esotropia since age 1½ years and had amblyopia that did not respond to patching or atropine treatment. The retinal abnormalities were first noted at age 3 years. He was born full-term with a birth weight of 3.0 kg. His past medical history was significant for microcephaly (head circumference measures not available). His best-corrected vi-

sual acuity was 20/50 OD and 20/210 OS, with a correction of +1.50 + 1.50X85 in the right eye and -1.25 + 1.75X100 in the left eye. Chorioretinal atrophy without retinal detachment was present in both eyes (Figure 2A). The younger sister was examined at age 5½ years. She had a history of eyes drifting in or out intermittently and had received patching and atropine treatment that had been discontinued at age 2 years. She also was noted to have microcephaly. On eye examination, her best-corrected visual acuity was 20/170 OD (-2.25 + 2.25X70) and 20/120 OS (-4.25 + 5.25X110). She had a unilateral retinal fold in addition to bilateral areas of chorioretinopathy (Figure 2B).

The *KIF11* p.A218Gfs*15 mutation was discovered in a boy who was born full-term and had a history of vision problems since age 6 months. There was no family history of *FEVR* or congenital retina problems. In addition, there was no family history of microcephaly or lymphedema, and the family noted large head sizes on the paternal side. The boy was not noted to have an anomalous head size on review of his medical records. At age 4 years, he had noncentral, unsteady, and non-maintained fixation in either eye (cycloplegic refraction +1.00 in the right eye and +0.50 in the left eye). He had an alternating esotropia of 40 prism diopters. On fundus examination, retinal folds were present in both eyes without evidence of chorioretinopathy typically seen in *MLCRD* (Figure 2C). Intravenous fluorescein angiography revealed peripheral avascular areas of retina typical of *FEVR* (Figure 2D). Results of a head computed tomography scan were reported as normal. Following the discovery of the *KIF11* mutation, the head circumference was measured and found to be below the second percentile. The parents and a sibling did not share the *KIF11* mutation, suggesting that the mutation occurred de novo in the proband.

The *KIF11* p.R47X mutation was identified in a 6-year-old boy who was born full-term. The visual acuity was light perception with both eyes open. The retina was completely detached, forming a fibrous mass behind the lens on the right side. The left eye manifested a falciform fold. The anterior segment was normal. The child was lost to follow-up, precluding further investigation of head circumference measurements and formal systemic evaluation. The parents' fundus examination including IVFA showed peripheral vascular tortuosity in both. Tissue samples were not collected from the parents.

The intronic sequence mutation *KIF11* c.790-1G>T was identified in a singleton with recognized microcephaly and diagnosed with *FEVR* by the ophthalmologist. There was no family history of health problems compatible with a diagnosis of *FEVR* or *MLCRD*, and the mutation was not found in either parent or sister. The boy was born full-term. Vision problems were noted soon after birth at which time he received a diagnosis of *FEVR*. At age 2, his development appeared normal, but his head circumference was 5 SDs below the mean. On eye examination, he had a large-angle esotropia with leukocoria on the right side. At age 15 months, he was able to fix and follow objects with his left eye (+3.25-2.00X180) but had no light perception on the right side. The right eye had a complete retinal detachment with a cloudy cornea. The left fundus showed peripheral vitreous condensation with subretinal fibrosis inferiorly after prior treatment with cryotherapy and scleral buckle.

Figure 1. Novel *KIF11* Mutations

A-D, Location of novel *KIF11* mutations with respect to the genomic organization of *KIF11*. Three of the novel mutations, *KIF11* p.E470X (c.1408G>T), p.A218Gfs*15 (c.652DupG), and *KIF11* c.790-1G>T, lead to a truncation (asterisks) of the *KIF11* protein coding sequence. *KIF11* c.790-1G>T alters an acceptor splice site. The mutation occurs in a nucleotide that is highly conserved across a variety of taxa. The mutation was evaluated for functional significance using the software CRYP-SKIP, and Spliceman. Both programs predicted that the mutation has a high probability of affecting splicing. The nonsynonymous *KIF11*


p.R221G (c.661 A>G) mutation segregated with the disease in a pedigree with 4 affected individuals. The mutation, located in the kinesin motor domain, was evaluated for functional significance using the software programs PolyPhen-2, PROVEAN, and SIFT. All programs predicted that the mutation is damaging. The mutation was absent in the 300 in-house exomes and the Exome Variant Server, as well as in 180 chromosomes from a random white population. E, Sequence comparison of the *KIF11* protein demonstrating the evolutionary conservation of the mutated *KIF11* arginine 221 residue across a variety of taxa.

Three probands were recruited specifically to investigate a genetic origin for likely or possible MLCRD. None had retinal detachment. Of the 3 pedigrees, one family of Acadian descent with 4 affected members was identified with *KIF11* p.R221G. The proband was born full-term. She was noted to have microcephaly in infancy, but according to the family, this was deemed to represent uncomplicated hereditary microcephaly before discovering the eye problems, because she was meeting her developmental milestones and her mother was also microcephalic but otherwise healthy. There were no siblings, and a maternal uncle was known to have significant developmental delays, having lived most of his life in a special

care facility, and he had a long-standing history of blindness. The family was further investigated by medical genetics (eTable in the Supplement). Although not examined, the maternal grandmother's head size was described as normal but the maternal uncle with significant developmental delays was known to have a small head size. Medical ocular records of the maternal uncle described retinal lesions similar to those found in the proband.

At age 20 months, the proband had a right esotropia and subnormal visual behavior. The corneal diameter was reduced at 10 mm with mild sclerocornea over 360° in both eyes. The axial length was normal at 20.4 mm in the right eye and

Figure 2. Fundus Photographs and Intravenous Fluorescein Angiography (IVFA)



A, E470X proband: the right fundus shows peripheral atrophic changes inferotemporally and an area of mild elevation. The left fundus manifests an inferior peripheral area of chorioretinal atrophy with retinal pigment epithelium clumping. B, E470X sister: a peripheral retinal fold courses inferotemporally with areas of atrophy in the right eye and peripheral retinal pigmentary changes only on the left. C, A218Gfs*15 proband: fibrovascular tissue with retinal folds involving the macula was present in both eyes without evidence of chorioretinopathy typically seen in microcephaly, lymphedema, and chorioretinal dysplasia. Retinal pigment epithelial changes were evident in keeping with retinal detachment and ischemic stress. D, IVFA revealed

peripheral areas of an avascular retina. E, R221G proband: multiple inferior chorioretinal atrophic lesions are present with moderate, diffuse atrophy of the optic nerve. The macula is featureless and the retinal vessels are fine and straightened, in keeping with severe retinal dysplasia. The vascular arcade is dragged temporally toward an atrophic scar with a temporal fibrous mass. F, IVFA revealed an area of avascular retina peripheral to the chorioretinal scars and the fibrous mass as well as retinal vascular anomalous formation in the areas without chorioretinal atrophy, resembling what is typically seen in familial exudative vitreoretinopathy with the presence of a fibrovascular mass at the junction of the avascular area.

20.16 mm in the left eye, and the cycloplegic refraction was +1.50 spherical equivalent hyperopic astigmatism in both eyes. Multiple bilateral inferior chorioretinal atrophic lesions were present. The macula was dragged peripherally and there was retinal detachment in both eyes (Figure 2E). Intravenous fluorescein angiography revealed bilateral areas of peripheral avascular retina that was otherwise difficult to detect with indirect ophthalmoscopy alone (Figure 2F). The proband's mother underwent a full eye examination including IVFA that failed to show vascular abnormalities or chorioretinopathy.

Discussion

In this study, we have extended the description of the spectrum of ocular manifestation resulting from mutations in *KIF11* to include retinal detachment that can mimic FEVR. Ostergaard et al¹ recently reported on cases of MLCRD/CDMMR associated with *KIF11* mutations (OMIM 152950). The major ocular phenotype described in 27 patients consisted of cho-

rioretinopathy characterized by peripheral areas of atrophic patches involving the choroid and retina. Other ocular characteristics included refractive errors, nystagmus, strabismus, and only a single eye with a total retinal detachment. Because previous reports³⁵⁻³⁷ of MLCRD and CDMMR have described the presence of retinal folds, persistent fetal vasculature, and microphthalmia, none of which were present in their patients with *KIF11* mutations, the authors¹ suggested that microcephaly with retinal folds and microphthalmia could be caused by mutations in a different gene. Consistent with this hypothesis, the only other patient with MLCRD who had a *KIF11* mutation also did not have a retinal fold or detachment.³⁴

The present study suggests that retinal folds and detachments are more common in the *KIF11*-related MLCRD spectrum of conditions than was previously reported, with the identification of *KIF11* mutations in 5.6% of all FEVR probands and 14.3% (4 of 28) of probands without a known FEVR gene mutation. The difference in prevalence could be attributed to ascertainment bias. Children with FEVR presenting with partial or complete retinal detachments at a very young age should be examined for mild to moderate microcephaly that could

point to a more precise diagnosis. Microcephaly that ranges from mild to severe,¹ especially in the absence of developmental delays, can be easily overlooked as unrelated to the FEVR phenotype. Patients with FEVR should be carefully examined for the presence of microcephaly as a marker for *KIF11*-related disease, because differentiating FEVR from MLCRD/CDMMR is important for the accuracy of genetic counseling. Of particular significance is the possibility of variable degrees of mental retardation, as is the case for our pedigree carrying the *KIF11* p.R221G mutation.

An ocular feature that can help distinguish FEVR from MLCRD/CDMMR is the presence of chorioretinopathy typically affecting the inferior retina. However, this feature may be subtle or absent, as was the case for the patient with *KIF11* p.A218Gfs*15. We have shown that the identification of a peripheral avascular retina on examination and/or IVFA does not exclude *KIF11*-related disease. Our data do not enable us to determine the exact frequency of *KIF11* mutations in cases with such a presentation at a young age, but we hypothesize that the frequency is likely to be higher in those patients than the 5.6% of all FEVR cases in our cohort.

In FEVR, the primary abnormality is variable incomplete vascularization of the peripheral retina, a process that should be largely completed by term birth. Secondary fibrovascular proliferation is responsible for retinal detachment that can range from partial to complete. More severe cases usually present in infancy or early childhood. Until now, the cause of retinal detachment in some cases of MLCRD and CDMMR had been unresolved. To our knowledge, we have demonstrated for the first time that *KIF11* mutations can result in abnormal peripheral retinal vascular development that can present with or without retinal detachment, which is identical to what occurs in patients with FEVR. This feature is different from retinal vascular attenuation or paucity associated with retinal dysplasia. The identification of peripheral avascular retina is impor-

tant to minimize vision loss from secondary complications. Our findings suggest that newly diagnosed cases and newborns at risk for MLCRD should receive IVFA as part of their workup with consideration to treat the avascular zones with laser in selected cases to avoid tractional detachment and prevent vision loss. At a minimum, such patients will require closer follow-up, especially at a young age, to detect potential complications early and improve the visual outcome.

A kinesin family member motor protein, *KIF11* localizes to spindle microtubules during mitosis and is required for mitotic progression. Several genes with roles in spindle development, including *KIF11*, result in microcephaly when mutated.¹ An insertional mutagenesis screen in zebrafish identified the *kif11* gene as being required for normal development of secondary motor neurons and oligodendrocytes, as well as for glial cell viability. Glial cells from the *kif11* mutants arrested in mitosis and subsequently underwent apoptotic cell death.³⁸ However, the role of *KIF11* in retinal vascular development is less clear. Of interest is the report of a case of congenital retinal folds, microcephaly, and mild mental retardation associated with a chromosomal abnormality disrupting *CDK19*,³⁹ suggesting that genes other than *KIF11* that can cause microcephaly and that retinal folds likely play a role in the development of retinal vasculature. Further research will clarify the role of these genes in the development of retinal vessels.

Conclusions

Mutations of *KIF11* are associated with a broad spectrum of phenotypes ranging from apparently classical FEVR to MLCRD and CDMMR. Careful clinical examination is necessary to identify associated microcephaly. Genotyping patients with newly diagnosed FEVR should include *KIF11* in addition to classical FEVR genes.

ARTICLE INFORMATION

Submitted for Publication: March 5, 2014; final revision received April 29, 2014; accepted April 29, 2014.

Published Online: August 14, 2014.
doi:10.1001/jamaophthalmol.2014.2814.

Author Affiliations: IWK Health Centre Eye Care Team, Halifax, Nova Scotia, Canada (Robitaille, Wallace); Department of Ophthalmology and Visual Sciences, Dalhousie University, Halifax, Nova Scotia, Canada (Robitaille, Wallace); Department of Pathology, Dalhousie University, Halifax, Nova Scotia, Canada (Robitaille, Gillett, LeBlanc, Gaston, Nightingale, Mackley, Bedard); Department of Pediatrics, Maritime Medical Genetics Service, Dalhousie University, Halifax, Nova Scotia, Canada (Parkash); Providence Health Care Heart Centre, St. Paul's Hospital, Vancouver Coastal Health, Vancouver, British Columbia, Canada (Hathaway); Maritime Medical Genetics Service, IWK Health Centre, Halifax, Nova Scotia, Canada (Thomas); Department of Surgery, University of Calgary, Alberta Children's Hospital Research Institute, Calgary, Alberta, Canada (Ells); Cole Eye Institute, Cleveland Clinic, Cleveland, Ohio (Traboulsi); Department of Ophthalmology, The Hospital for

Sick Children, University of Toronto, Toronto, Ontario, Canada (Héon); Réseau de santé Vitalité Health Network, Hôpital régional Chaleur Regional Hospital, Bathurst, New Brunswick, Canada (Roy); Genetic Institute, Emek Medical Center, Afula, Rappaport School of Medicine, Technion, Haifa, Israel (Shalev); Department of Pediatrics, Pediatric Oncology, Dalhousie University, Halifax, Nova Scotia, Canada (Fernandez); Department of Ophthalmology, Capital Health, Halifax, Nova Scotia, Canada (MacGillivray); Department of Human Genetics, Faculty of Medicine, McGill University, Montreal, Quebec, Canada (Fahiminiya, Majewski); Genome Quebec Innovation Center, Montreal, Quebec, Canada (Fahiminiya, Majewski); Department of Pharmacology, Dalhousie University, Halifax, Nova Scotia, Canada (McMaster).

Author Contributions: Drs Robitaille and Bedard had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Robitaille, LeBlanc, Ells, Shalev, Fernandez, McMaster, Bedard.

Acquisition, analysis, or interpretation of data: Robitaille, Gillett, LeBlanc, Gaston, Nightingale, Mackley, Parkash, Hathaway, Thomas, Ells, Traboulsi, Héon, Roy, MacGillivray, Wallace,

Fahiminiya, Majewski, McMaster, Bedard.

Drafting of the manuscript: Robitaille, Gillett, LeBlanc, Gaston, Nightingale, Mackley, Shalev, McMaster.

Critical revision of the manuscript for important intellectual content: Robitaille, Gillett, LeBlanc, Gaston, Parkash, Hathaway, Thomas, Ells, Traboulsi, Héon, Roy, Fernandez, MacGillivray, Wallace, Fahiminiya, Majewski, McMaster, Bedard.

Statistical analysis: Robitaille, Gaston, Héon, Bedard.

Obtained funding: Robitaille, Fernandez, Majewski, McMaster, Bedard.

Administrative, technical, or material support: Robitaille, Gillett, Gaston, Nightingale, Mackley, Thomas, Ells, Traboulsi, MacGillivray, Wallace, McMaster, Bedard.

Study supervision: Robitaille, Gillett, Nightingale, Ells, McMaster, Bedard.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Funding/Support: Orphan Diseases: Identifying Genes and Novel Therapeutics to Enhance Treatment Project is funded under the Large Scale

Applied Research Program of Genome Canada with matching funding provided by Health Canada; Departments of Medicine and Ophthalmology, Dalhousie University Faculty of Medicine; the Dalhousie Medical Research Fund; the Nova Scotia Health Research Foundation; the Dalhousie University Industry and Liaison Office; Genome British Columbia; the Center for Drug Research and Development, Vancouver, British Columbia; the Capital District Health Authority; and the Department of Health and Wellness of Nova Scotia.

Role of the Funder/Sponsor: The sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We wish to acknowledge the support and effort of the participating families.

REFERENCES

- Ostergaard P, Simpson MA, Mendola A, et al. Mutations in *KIF11* cause autosomal-dominant microcephaly variably associated with congenital lymphedema and chorioretinopathy. *Am J Hum Genet*. 2012;90(2):356-362.
- Robitaille JM, Wallace K, Zheng B, et al. Phenotypic overlap of familial exudative vitreoretinopathy (FEVR) with persistent fetal vasculature (PFV) caused by *FZD4* mutations in two distinct pedigrees. *Ophthalmic Genet*. 2009;30(1):23-30.
- Chen ZY, Battinelli EM, Fielder A, et al. A mutation in the Norrie disease gene (*NDP*) associated with X-linked familial exudative vitreoretinopathy. *Nat Genet*. 1993;5(2):180-183.
- Robitaille J, MacDonald ML, Kaykas A, et al. Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy. *Nat Genet*. 2002;32(2):326-330.
- Toomes C, Bottomley HM, Jackson RM, et al. Mutations in *LRP5* or *FZD4* underlie the common familial exudative vitreoretinopathy locus on chromosome 11q. *Am J Hum Genet*. 2004;74(4):721-730.
- Jiao X, Ventruito V, Trese MT, Shastri BS, Hejtmancik JF. Autosomal recessive familial exudative vitreoretinopathy is associated with mutations in *LRP5*. *Am J Hum Genet*. 2004;75(5):878-884.
- Poulter JA, Ali M, Gilmour DF, et al. Mutations in *TSPAN12* cause autosomal-dominant familial exudative vitreoretinopathy. *Am J Hum Genet*. 2010;86(2):248-253.
- Poulter JA, Davidson AE, Ali M, et al. Recessive mutations in *TSPAN12* cause retinal dysplasia and severe familial exudative vitreoretinopathy (FEVR). *Invest Ophthalmol Vis Sci*. 2012;53(6):2873-2879.
- Collin RW, Nikopoulos K, Dona M, et al. *ZNF408* is mutated in familial exudative vitreoretinopathy and is crucial for the development of zebrafish retinal vasculature. *Proc Natl Acad Sci U S A*. 2013;110(24):9856-9861.
- GeneTests. <http://www.genetests.org>. Accessed April 29, 2014.
- Fahiminiya S, Almuriekh M, Nawaz Z, et al. Whole exome sequencing unravels disease-causing genes in consanguineous families in Qatar [published online September 14, 2013]. *Clin Genet*. doi:10.1111/cge.12280.
- Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009;25(14):1754-1760.
- McKenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010;20(9):1297-1303.
- Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res*. 2010;38(16):e164.
- Stenson PD, Mort M, Ball EV, Shaw K, Phillips AD, Cooper DN. The Human Gene Mutation Database: building a comprehensive mutation repository for clinical and molecular genetics, diagnostic testing and personalized genomic medicine. *Hum Genet*. 2014;133(1):1-9.
- Sherry ST, Ward MH, Kholodov M, et al. dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res*. 2001;29(1):308-311.
- Abecasis GR, Auton A, Brooks LD, et al; 1000 Genomes Project Consortium. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012;491(7422):56-65.
- Exome Variant Server. NHLBI GO Exome Sequencing Project (ESP). <http://evs.gs.washington.edu/EVS/>. Accessed April 1, 2014.
- Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc*. 2009;4(7):1073-1081.
- Ng PC, Henikoff S. Accounting for human polymorphisms predicted to affect protein function. *Genome Res*. 2002;12(3):436-446.
- Ng PC, Henikoff S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res*. 2003;31(13):3812-3814.
- Ng PC, Henikoff S. Predicting the effects of amino acid substitutions on protein function. *Annu Rev Genomics Hum Genet*. 2006;7:61-80.
- Ng PC, Henikoff S. Predicting deleterious amino acid substitutions. *Genome Res*. 2001;11(5):863-874.
- Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7(4):248-249.
- Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. *PLoS One*. 2012;7(10):e46688. doi:10.1371/journal.pone.0046688.
- Choi Y. A fast computation of pairwise sequence alignment scores between a protein and a set of single-locus variants of another protein. In: *Proceedings of the ACM Conference on Bioinformatics, Computational Biology and Biomedicine*. New York, NY: Association for Computing Machinery; 2012:414-417.
- Buratti E, Chivers M, Královicová J, et al. Aberrant 5' splice sites in human disease genes: mutation pattern, nucleotide structure and comparison of computational tools that predict their utilization. *Nucleic Acids Res*. 2007;35(13):4250-4263.
- Divina P, Kvitekovicova A, Buratti E, Vorechovsky I. Ab initio prediction of mutation-induced cryptic splice-site activation and exon skipping. *Eur J Hum Genet*. 2009;17(6):759-765.
- Královicová J, Vorechovsky I. Global control of aberrant splice-site activation by auxiliary splicing sequences: evidence for a gradient in exon and intron definition. *Nucleic Acids Res*. 2007;35(19):6399-6413.
- Vorechovsky I. Aberrant 3' splice sites in human disease genes: mutation pattern, nucleotide structure and comparison of computational tools that predict their utilization. *Nucleic Acids Res*. 2006;34(16):4630-4641.
- Lim KH, Fairbrother WG. Spliceman—a computational web server that predicts sequence variations in pre-mRNA splicing. *Bioinformatics*. 2012;28(7):1031-1032.
- Lim KH, Ferraris L, Filloux ME, Raphael BJ, Fairbrother WG. Using positional distribution to identify splicing elements and predict pre-mRNA processing defects in human genes. *Proc Natl Acad Sci U S A*. 2011;108(27):11093-11098.
- Kent WJ, Sugnet CW, Furey TS, et al. The human genome browser at UCSC. *Genome Res*. 2002;12(6):996-1006.
- Hazan F, Ostergaard P, Ozturk T, et al. A novel *KIF11* mutation in a Turkish patient with microcephaly, lymphedema, and chorioretinal dysplasia from a consanguineous family. *Am J Med Genet A*. 2012;158A(7):1686-1689.
- Casteels I, Devriendt K, Van Cleynenbreugel H, Demareel P, De Tavernier F, Fryns JP. Autosomal dominant microcephaly-lymphoedema-chorioretinal dysplasia syndrome. *Br J Ophthalmol*. 2001;85(4):499-500.
- Fryns JP, Smeets E, Van den Berghe H. On the nosology of the "primary true microcephaly, chorioretinal dysplasia, lymphoedema" association. *Clin Genet*. 1995;48(3):131-133.
- Trzupek KM, Falk RE, Demer JL, Weleber RG. Microcephaly with chorioretinopathy in a brother-sister pair: evidence for germ line mosaicism and further delineation of the ocular phenotype. *Am J Med Genet A*. 2007;143A(11):1218-1222.
- Barresi MJ, Burton S, Dipietrantonio K, Amsterdam A, Hopkins N, Karlstrom RO. Essential genes for astroglial development and axon pathfinding during zebrafish embryogenesis. *Dev Dyn*. 2010;239(10):2603-2618.
- Mukhopadhyay A, Kramer JM, Merckx G, et al. *CDK19* is disrupted in a female patient with bilateral congenital retinal folds, microcephaly and mild mental retardation. *Hum Genet*. 2010;128(3):281-291.