

National Trends in Hospitalizations for Opioid Poisonings Among Children and Adolescents, 1997 to 2012

Julie R. Gaither, PhD, MPH, RN; John M. Leventhal, MD; Sheryl A. Ryan, MD; Deepa R. Camenga, MD, MHS

IMPORTANCE National data show a parallel relationship between recent trends in opioid prescribing practices and hospitalizations for opioid poisonings in adults. No similar estimates exist describing hospitalizations for opioid poisonings in children and adolescents.

OBJECTIVE To describe the incidence and characteristics of hospitalizations attributed to opioid poisonings in children and adolescents.

DESIGN, SETTING, AND PARTICIPANTS Retrospective analysis of serial cross-sectional data from a nationally representative sample of US pediatric hospital discharge records collected every 3 years from January 1, 1997, through December 31, 2012. The Kids' Inpatient Database was used to identify 13 052 discharge records for patients aged 1 to 19 years who were hospitalized for opioid poisonings. Data were analyzed within the collection time frame.

MAIN OUTCOMES AND MEASURES Poisonings attributed to prescription opioids were identified by codes from the *International Classification of Diseases, Ninth Revision, Clinical Modification*. In adolescents aged 15 to 19 years, poisonings attributed to heroin were also identified. Census estimates were used to calculate incidence per 100 000 population. The Cochran-Armitage test for trend was used to assess for changes in incidence over time.

RESULTS From 1997 to 2012, a total of 13 052 (95% CI, 12 500-13 604) hospitalizations for prescription opioid poisonings were identified. The annual incidence of hospitalizations for opioid poisonings per 100 000 children aged 1 to 19 years rose from 1.40 (95% CI, 1.24-1.56) to 3.71 (95% CI, 3.44-3.98), an increase of 165% (*P* for trend, <.001). Among children 1 to 4 years of age, the incidence increased from 0.86 (95% CI, 0.60-1.12) to 2.62 (95% CI, 2.17-3.08), an increase of 205% (*P* for trend, <.001). For adolescents aged 15 to 19 years, the incidence increased from 3.69 (95% CI, 3.20-4.17) to 10.17 (95% CI, 9.48-10.85), an increase of 176% (*P* for trend, <.001). In this age group, poisonings from heroin increased from 0.96 (95% CI, 0.75-1.18) to 2.51 (95% CI, 2.21-2.80), an increase of 161% (*P* for trend, <.001); poisonings involving methadone increased from 0.10 (95% CI, 0.03-0.16) to 1.05 (95% CI, 0.87-1.23), an increase of 950% (*P* for trend, <.001).

CONCLUSIONS AND RELEVANCE During the course of 16 years, hospitalizations attributed to opioid poisonings rose nearly 2-fold in the pediatric population. Hospitalizations increased across all age groups, yet young children and older adolescents were most vulnerable to the risks of opioid exposure. Mitigating these risks will require comprehensive strategies that target opioid storage, packaging, and misuse.

JAMA Pediatr. 2016;170(12):1195-1201. doi:10.1001/jamapediatrics.2016.2154
Published online October 31, 2016.

[+ Author Video Interview and JAMA Report Video](#)

[+ Supplemental content](#)

Author Affiliations: Department of Epidemiology and Public Health, Yale School of Medicine, New Haven, Connecticut (Gaither); Yale Center for Medical Informatics, Yale School of Medicine, New Haven, Connecticut (Gaither); Biomedical Informatics/Research Service, Veterans Affairs Connecticut Healthcare System, Department of Veterans Affairs, West Haven, Connecticut (Gaither); Department of Pediatrics, Yale School of Medicine, New Haven, Connecticut (Leventhal, Ryan, Camenga); Department of Emergency Medicine, Yale School of Medicine, New Haven, Connecticut (Camenga).

Corresponding Author: Julie R. Gaither, PhD, MPH, RN, Department of Epidemiology and Public Health, Yale School of Medicine, 333 Cedar St, PO Box 208064, New Haven, CT 06520 (julie.gaither@yale.edu).

Poisonings attributed to prescription drugs are now the leading cause of injury-related mortality in the United States^{1,2}; this epidemic is driven by the increased use of opioid analgesics to treat chronic pain.³⁻⁶ From 1999 to 2010, retail sales of prescription opioids quadrupled,⁴ and during this time, deaths attributed to opioid overdose rose 4-fold in those 15 to 64 years of age and 6-fold in those 15 to 24 years of age.⁷ In 2014 alone, opioids were blamed for 18 893 deaths in the United States.⁸

These mortality data, however, represent only a fraction of the serious adverse events attributed to prescription opioids.^{5,9-16} The increased availability of these medications has resulted in an unprecedented rise in opioid addiction and nonfatal overdoses.⁵ Emergency department (ED) visits for opioid-related indications have risen substantially in children and adults during the past 2 decades; ED visits for prescription opioid overdose, abuse, and misuse now rival those for illicit drugs, including heroin and cocaine.¹⁷⁻²¹ Even in children younger than 6 years, opioids, followed closely by benzodiazepines, now account for most of the drug poisonings in this age group¹⁸; in nearly all these poisonings, the child was exposed to a prescription intended for an adult in the household.²²

How many children and adolescents are hospitalized each year for opioid poisonings and how hospitalization rates have changed over time remain unknown. In young adults and those middle-aged and older, inpatient hospitalizations for opioid-related indications, such as for opioid dependence, abuse, and overdose, have increased 2- and 5-fold, respectively, in less than a decade.^{23,24} We hypothesized that similar trends in opioid poisonings have occurred in children and adolescents.

Therefore, the purpose of this study was to examine trends in hospitalizations for opioid poisonings in children and adolescents (aged 1-19 years). Moreover, because of the association between prescription opioid misuse and progression to heroin use,^{3,25-28} we examined trends in heroin overdose among those 15 to 19 years of age. In this same age group, we also examined poisonings attributed to methadone, a prescription opioid used to treat illicit and prescription opioid addiction and severe intractable pain. Recent analyses²⁹ have shown that despite the low frequency with which methadone is prescribed, it is responsible for a disproportionate number of drug poisonings. Given that opioids are already among the most widely prescribed medications in the United States—available in millions of US households^{6,30}—and that the US Food and Drug Administration recently approved the use of oxycodone hydrochloride (OxyContin) for children meeting certain criteria,³¹ a clearer understanding of pediatric opioid-related morbidity and mortality is needed.

Methods

Study Overview

From a national database of pediatric hospitalizations, we identified children and adolescents admitted to US hospitals for opioid poisonings in 1997, 2000, 2003, 2006, 2009, and 2012. We conducted a retrospective analysis of serial cross-sectional data abstracted from the Kids' Inpatient Database (KID).³² The KID, created by the Agency for Healthcare Research and Quality as part

Key Points

Question How has the prescription opioid epidemic affected pediatric hospitalization rates in the United States?

Findings This retrospective analysis of 13 052 national hospital discharge records found that pediatric hospitalizations for opioid poisonings increased nearly 2-fold from 1997 to 2012. Hospitalization rates were highest in older adolescents, but the largest percentage increase in hospitalizations over time occurred among the youngest children (toddlers and preschoolers).

Meaning Reducing pediatric opioid exposure and misuse will require a combination of public health interventions, policy initiatives, and consumer-product regulations.

of the Healthcare Cost and Utilization Project, is the only US database to provide all-payer data on pediatric services and clinical outcomes for children admitted as inpatients, routinely or through the ED.^{32,33} This study involved deidentified data and was determined to be exempt from approval and the need for informed consent by the Yale School of Medicine's institutional review board.

The KID has been released every 3 years since 1997 and is compiled from a random sample of pediatric discharge records collected from acute-care hospitals across the United States (the unit of observation is the inpatient record, regardless of the length of stay).³² For the most recent year of data, 2012, approximately 3.2 million records were selected from 4179 hospitals across 44 states. For a given year, 10% of uncomplicated hospital births and 80% of other pediatric hospitalizations (ie, patients younger than 19 years in 1997 and younger than 20 years beginning in 2000) are then sampled from among these records. Each discharge record is then weighted to facilitate national estimates. Weighting is accomplished by stratifying hospitals according to the following 6 characteristics: number of beds, teaching status, ownership or control (eg, private), hospital type (eg, children's hospital), rural or urban location, and US region. For each year of the KID, the unweighted database contains information on approximately 2 to 3 million annual pediatric discharges. When weighted, the KID provides estimates on approximately 6 to 7 million discharges.

Identification of the Sample

We identified hospitalizations attributed to opioid poisonings using diagnostic codes from the *International Classification of Disease, Ninth Revision, Clinical Modification (ICD-9-CM)*.³⁴ Admissions for prescription opioid poisonings were identified by the following ICD-9-CM poisoning codes: 965.00 (opium), 965.02 (methadone), and 965.09 (other opiates and narcotics). Among those 15 to 19 years of age, we also identified admissions for illicit opioid poisonings using the ICD-9-CM code 965.01 (heroin). We restricted the heroin and methadone analyses to those 15 to 19 years of age because most of these poisonings occurred in this age group.

To further characterize poisonings, we used E codes for external causes of injury to establish intent. These codes included E850.x to E858.x (unintentional [ie, accidental]), E950.0 to E950.5 (intentional [ie, suicide or self-inflicted]), and E980.x (undetermined).

Table. Weighted National Estimates of Demographic and Clinical Characteristics According to Year of Hospitalization for Prescription Opioid Poisonings

Characteristic	Study Year, No. (%) of Patients						P Value ^a
	1997	2000	2003	2006	2009	2012	
Weighted No. of hospitalizations (95% CI)	1049 (931-1167)	1474 (1333-1615)	2090 (1941-2238)	2436 (2267-2605)	3084 (2884-3283)	2918 (2706-3130)	NA
Age group, y							
1-4	133 (12.7)	146 (9.9)	210 (10.0)	244 (10.0)	377 (12.2)	421 (14.4)	<.001
5-9 ^b	NA	NA	NA	NA	NA	NA	
10-14	171 (16.3)	176 (11.9)	247 (11.8)	216 (8.9)	240 (7.8)	271 (9.3)	
15-19	715 (68.2)	1116 (75.7)	1595 (76.3)	1950 (80.0)	2427 (78.7)	2171 (74.4)	
Male sex	364 (34.7)	661 (44.8)	976 (46.7)	1305 (53.5)	1636 (53.0)	1382 (47.4)	<.001
Race or ethnicity							
White	550 (72.0)	938 (76.5)	1056 (72.3)	1270 (75.2)	1907 (75.0)	1886 (70.6)	<.001
Black	118 (15.4)	117 (9.5)	184 (12.6)	152 (9.0)	251 (9.9)	315 (11.8)	
Hispanic	64 (8.4)	118 (9.6)	141 (9.7)	156 (9.2)	210 (8.3)	271 (10.1)	
Other	33 (4.3)	54 (4.4)	80 (5.5)	111 (6.6)	173 (6.8)	198 (7.4)	
Insurance							
Medicaid	253 (24.1)	392 (26.6)	651 (31.1)	816 (33.5)	1164 (38.0)	1280 (44.0)	<.001
Private	643 (61.4)	824 (56.4)	1107 (53.0)	1163 (47.9)	1362 (44.4)	1249 (43.0)	
Self-pay	103 (9.8)	190 (13.0)	224 (10.7)	312 (12.8)	356 (11.6)	243 (8.4)	
Other	49 (4.7)	56 (3.8)	107 (5.1)	140 (5.8)	184 (6.0)	133 (4.6)	

Abbreviation: NA, not applicable.

^a Calculated using the χ^2 test.^b Estimates that do not meet the criteria for statistical reliability are not shown.

We restricted the sample to those 1 to 19 (inclusive) years of age. Infants were excluded to primarily capture those poisonings related to self-administration. To be consistent with the age categories used by the Centers for Disease Control and Prevention WISQARS (Web-based Injury Statistics Query and Reporting System) for reporting fatal and nonfatal injuries,³⁵ we grouped children into the following age categories: 1 to 4, 5 to 9, 10 to 14, and 15 to 19 years. The estimates for those aged 5 to 9 years (n = 135, unweighted), however, did not meet the criteria³⁶ for statistical reliability, and thus are not shown.

Statistical Analysis

Data were collected and analyzed from January 1, 1997, to December 31, 2012. Descriptive statistics were used to characterize the sample according to each year of the KID. Differences according to demographic and clinical characteristics were assessed with χ^2 tests for categorical variables and, as appropriate, 2-tailed *t* tests, analysis of variance, or Wilcoxon rank sum tests for continuous variables.

Nationally representative estimates were calculated using methods that take into account the complex survey design of the KID. Specifically, we used weights, strata, and cluster variables provided by the Healthcare Cost and Utilization Project to produce weighted estimates and SEs, which were then used to establish 95% CIs.³⁷ Unless otherwise noted, all counts, proportions, and incidence rates are based on weighted estimates.

To calculate the annual incidence of hospitalizations, we divided the weighted number of hospitalizations for opioid poisonings for a given year by the intercensal number of children at risk in that year.³⁸ The Cochran-Armitage test for trend³⁹ was used to assess changes in incidence over time.

All analyses were performed using SAS software (version 9.3; SAS Institute Inc). We applied a 2-sided statistical significance level of .05 to all analyses.

Results


Demographic Characteristics

Among children and adolescents aged 1 to 19 years admitted to US hospitals every 3 years from 1997 through 2012, we identified a total of 13 052 (95% CI, 12 500-13 604) hospitalizations for prescription opioid poisonings. Demographic characteristics for each of the 6 assessment periods are shown in the **Table**. Those in the oldest group accounted for the largest proportion of poisonings, whereas those aged 5 to 9 years accounted for the smallest. In 1997, males accounted for 34.7% (95% CI, 31%-39%) of hospitalizations; by 2012, this proportion was 47.4% (95% CI, 45%-50%). Children were predominately white (73.5%; 95% CI, 72%-75%) and covered by private insurance (48.8%; 95% CI, 48%-50%); however, the proportion insured by Medicaid increased from 24.1% (95% CI, 20%-28%) in 1997 to 44.0% (95% CI, 42%-46%) in 2012 (*P* for trend, <.001). Across the 6 years of the KID, 176 children (1.3%; 95% CI, 1.1%-1.6%) died during hospitalization (in compliance with Healthcare Cost and Utilization Project privacy protections, we do not present these data stratified by year because of small cell frequencies).

Trends in Hospitalizations for Prescription Opioid Poisonings


From 1997 to 2012, the annual incidence of hospitalizations for opioid poisonings per 100 000 children rose from 1.40 (95%

Figure 1. Weighted National Estimates of Temporal Trends in Hospitalizations for Prescription Opioid Poisonings Stratified by Age Category



Error bars indicate 95% CI (*P* for trend, <.001 for all ages). Estimates for 5- to 9-year-olds do not meet the criteria for statistical reliability and thus are not shown.

Figure 2. Weighted National Estimates of Temporal Trends in Prescription Opioid Poisonings by Intent in the Group Aged 15 to 19 Years



Error bars indicate 95% CI (*P* for trend, <.001 for poisonings attributed to accidental and suicidal intent).

CI, 1.24-1.56) to 3.71 (95% CI, 3.44-3.98), an increase of 165% (*P* for trend, <.001). As shown in **Figure 1** and eTable 1 in the **Supplement**, when examined by age category, the greatest change occurred among the group aged 1 to 4 years, among whom the incidence rose from 0.86 (95% CI, 0.60-1.12) in 1997

to 2.62 (95% CI, 2.17-3.08) in 2012, an increase of 205% (*P* for trend, <.001). For adolescents aged 15 to 19 years, the incidence rose from 3.69 (95% CI, 3.20-4.17) in 1997 to 10.17 (95% CI, 9.48-10.85) in 2012, an increase of 176% (*P* for trend, <.001).

When prescription opioid poisonings were examined by intent, among children younger than 10 years, 16 poisonings (0.92%; 95% CI, 0.37%-1.47%) were attributed to suicide or self-inflicted injury from 1997 to 2012. In children 10 to 14 years of age, the incidence of poisonings attributed to suicide or self-inflicted injury increased from 0.62 (95% CI, 0.44-0.79) in 1997 to 0.85 (95% CI, 0.68-1.02) in 2012, an increase of 37% (*P* for trend, <.001). The incidence of poisonings attributed to accidental intent increased from 0.17 (95% CI, -0.04 to 0.39) in 1997 to 0.31 (95% CI, 0.21 to 0.42) in 2012, an increase of 82% (*P* for trend, <.001).

These trends were more marked in adolescents 15 to 19 years of age. Poisonings attributed to suicide or self-inflicted injury increased by 140% (*P* for trend, <.001), whereas those attributed to accidental intent increased 303% (*P* for trend, <.001). Incidence rates, stratified by year, for this age group are shown in **Figure 2** and eTable 2 in the **Supplement**.

Trends in Hospitalizations for Heroin and Methadone Poisonings Among 15- to 19-Year-Olds

As shown in **Figure 3** and eTable 2 in the **Supplement**, prescription opioids accounted for most opioid poisonings across all time points. However, poisonings involving heroin increased from 0.96 (95% CI, 0.75-1.18) in 1997 to 2.51 (95% CI, 2.21-2.80) in 2012, an increase of 161% (*P* for trend, <.001). Poisonings involving methadone increased from 0.10 (95% CI,

0.03-0.16) in 1997 to 1.05 (95% CI, 0.87-1.23) in 2012, an increase of 950% (P for trend, $<.001$).

Discussion

Using a nationally representative sample of pediatric hospital discharge records, we found that the incidence of hospitalizations for prescription opioid poisonings among children and adolescents 1 to 19 years of age increased nearly 2-fold from 1997 to 2012. Although rates increased across all age groups, the largest percentage increase occurred among the youngest children aged 1 to 4 years. The second largest increase occurred among adolescents 15 to 19 years of age, among whom hospitalizations for prescription opioid and heroin poisoning increased. Adolescents in this age group had the highest incidence overall for each of the 6 years examined.


In contrast to poisoning intent in younger children, poisonings in those older than 10 years were primarily attributed to suicide or self-inflicted injury. During the 16-year period, poisonings attributed to suicide or self-inflicted injury among those aged 15 to 19 years increased by 140%. These data underscore the dangers associated with the widespread availability of prescription opioids, particularly for adolescents at risk for depression (many adolescents are also thought to self-medicate with prescription opioids).^{16,40}

Although poisonings attributed to suicide or self-inflicted injury among those aged 15 to 19 years exceeded those attributed to accidental poisonings across all time points, accidental poisonings increased at a more rapid rate, 3-fold from 1997 to 2012. Increasing rates of prescription opioid misuse and abuse, rather than therapeutic errors or adverse effects among adolescents prescribed opioids for pain management, are likely driving these trends in accidental poisonings.¹⁹

Mirroring demographic trends seen in adults,^{23,24} we found that approximately three-quarters of hospitalizations occurred in white individuals. Similarly, private insurers were found to be the largest payer. However, the percentage of hospitalizations covered by private payers decreased over time, whereas the percentage paid by Medicaid increased. In 2012, Medicaid covered 44.0% of hospitalizations for pediatric opioid poisonings, up from 24.1% in 1997. This change may reflect trends in Medicaid payments for all hospitalizations; from 1997 to 2011, the percentage of hospitalizations where Medicaid was the primary payer increased by 34%.⁴¹ This increase, however, has largely been attributed to a rise in hospitalizations for a small number of conditions. Thus, the shift over time in payers for opioid hospitalizations may instead be indicative of the widening effect of opioid use across sociodemographic groups.

This study—to our knowledge, the first to extensively examine pediatric hospitalizations due to opioid poisonings—contributes to a broader understanding of the public health crisis caused by the now widespread availability of prescription opioids in the United States. Our findings are consistent with previous research demonstrating that as physicians have increasingly relied on prescription opioids to treat chronic pain during the past 2 decades, rates for opioid poisonings have risen

Figure 3. Weighted National Estimates of Temporal Trends in Hospitalizations for Illicit vs Prescription Opioid Poisonings in the Group Aged 15 to 19 Years



Error bars indicate 95% CI (P for trend, $<.001$ for poisonings attributed to heroin and prescription opioid drug use).

in tandem.^{1-4,9} For young children, calls to poisoning control centers for opioid ingestion have increased markedly in recent years,^{19,20,22} as have ED visits across all age groups.¹⁷⁻²¹ In addition, a number of studies have examined trends in ED visits and subsequent hospitalizations for pharmaceutical poisonings and found that prescription opioids were among the most commonly implicated medications and that these poisonings frequently resulted in hospital admission.^{17,19,20}

Emerging data suggest that in recent years physicians have been prescribing opioids less frequently. From 2011 to 2013, the number of opioid prescriptions dispensed in the United States slightly decreased.^{25,42} Our data show a similar marginal decrease in the overall incidence of hospitalizations for prescription opioid poisonings from 2009 to 2012. This change, however, was driven entirely by a slight (7%) but statistically significant decrease in hospitalizations among those aged 15 to 19 years. In contrast, in this age group, hospitalizations involving heroin overdose continued to rise. This finding is consistent with research suggesting that prescription opioid abuse may be a precursor to initiation of heroin use,^{3,25-28} which can be explained in part by the low cost of heroin compared with opioid analgesics such as oxycodone.²⁶⁻²⁸

Notably, we found that the incidence of hospitalizations for methadone poisonings increased 950% among 15- to 19-year-olds from 1997 to 2012. Methadone is among the most misused of prescription drugs and is commonly diverted for illicit (nonmedical) purposes, such as getting high or to enhance the effects of alcohol or other drugs.³ Our data suggest that the incidence of poisonings attributed to methadone misuse may be slowing. From 2009 to 2012, hospitalization rates for methadone misuse decreased slightly among those aged 15 to 19 years. These findings are consistent with recent data showing that rates for

prescription opioid abuse, misuse, and diversion for nonmedical use have plateaued or decreased in the general adult population.⁴³ Declining rates for abuse and diversion have also been seen among those aged 12 to 17 years.⁴⁴

Nonmedical use of prescription opioids, nevertheless, remains a substantial problem, particularly for adolescents. In 1 study,^{44,45} nearly 10% of high school seniors reported using opioids nonmedically. Although most acquired these medications from family and friends, the source for nearly 40% of students was their own prior prescription.⁴⁶ This information is in keeping with findings that adolescents are frequently prescribed opioids for common injury- and non-injury-related indications.⁴⁶ A recent study of ambulatory care visits in the United States, for example, reports that from 2005 to 2007, opioids were prescribed at 22% of visits among adolescents presenting with back pain.⁴⁷ A separate study of commercial medical and pharmacy claims data from 2007 to 2008 found that nearly 50% of patients aged 13 to 17 years presenting with first-time complaints of headache received an opioid prescription during the 2-year follow-up, and nearly one-third received 3 or more prescriptions.³¹ Of note, the US Food and Drug Administration recently extended the off-label use of oxycodone for children as young as 11 years with certain types of intractable pain—namely cancer pain.⁴⁸ Despite these trends, however, the new national opioid prescribing guidelines⁴⁹ by the Centers for Disease Control and Prevention fail to include recommendations for patients younger than 18 years.⁵⁰

Limitations

This study has at least 4 limitations. The major limitation is that the national estimates are based on ICD-9-CM codes, which are subject to miscoding and errors of omission. Although limited data are available regarding the accuracy of opioid poisoning codes, validation studies involving other common pediatric medications (eg, acetaminophen) support the use of ICD-9-CM codes for identifying poisonings in children.⁵¹ Second, we were unable to provide a full clinical picture or psychosocial profile (eg, the circumstances that led to the poisoning) of the children admitted for opioid poisonings or to validate the poisoning codes with toxicologic results. Similarly, we

were not able to distinguish between opioids prescribed for chronic pain and those, such as buprenorphine hydrochloride and methadone, prescribed for opioid dependence (ie, methadone maintenance treatment). Third, each record within the KID represents a hospitalization; thus, children admitted within a given year for 1 or more repeated opioid poisonings would be counted multiple times. We believe, however, that this limitation is minor because such instances are likely rare. Finally, similar to other recent studies^{17,19,20} examining trends in opioid poisonings in the young, we lacked the data to determine whether the trends we report continued beyond 2012.

Conclusions

The current public health crisis caused by prescription opioids is a systemic issue that affects individuals across the age spectrum. To date, numerous initiatives at the local, state, and federal levels have been implemented to combat prescription opioid poisonings and overdose; most of these efforts, however, have focused on modifying physician prescribing practices for those treating adults with chronic pain.⁵² The full effect of these initiatives remains to be seen, but recent data indicate that the United States is making progress in gaining control of the epidemic caused by opioid analgesics.³

Our research, however, suggests that poisonings by prescription and illicit opioids are likely to remain a persistent and growing problem in the young unless greater attention is directed toward the pediatric community, who make up nearly one-quarter of the US population. A combination of public health interventions (eg, parental education), policy initiatives, and consumer-product regulations is needed to reduce pediatric exposure to opioids.^{50,52} In addition, further resources should be directed toward addressing opioid misuse and abuse during adolescence. Of particular importance are prevention programs that address the overlap in opioid misuse and depression among adolescents.¹⁶ Finally, for clinicians who treat acute and chronic pain in children, national clinical practice guidelines for opioid prescribing that include pediatric-specific recommendations are urgently needed.⁵⁰

ARTICLE INFORMATION

Accepted for Publication: June 21, 2016.

Published Online: October 31, 2016.

doi:10.1001/jamapediatrics.2016.2154

Author Contributions: Dr Gaither had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: All authors.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Gaither, Leventhal, Camenga.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Gaither.

Obtained funding: Gaither, Camenga.

Administrative, technical, or material support: Gaither.

Study supervision: Gaither, Leventhal.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study was supported by grants F31DA035567 (Dr Gaither) and K12DA033012 (Dr Camenga) from the National Institute on Drug Abuse.

Role of the Funder/Sponsor: The funding source had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the Yale School of Medicine.

REFERENCES

1. Warner M, Chen LH, Makuc DM. Increase in fatal poisonings involving opioid analgesics in the United States, 1999-2006. *NCHS Data Brief*. 2009;(22):1-8.

2. Chen LH, Hedegaard H, Warner M.

Drug-poisoning deaths involving opioid analgesics: United States, 1999-2011. *NCHS Data Brief*. 2014;(166):1-8.

3. Dart RC, Surratt HL, Cicero TJ, et al. Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med*. 2015;372(3):241-248.

4. Manchikanti L, Helm S II, Fellows B, et al. Opioid epidemic in the United States. *Pain Physician*. 2012;15(3)(suppl):ES9-ES38.

5. Von Korff M, Kolodny A, Deyo RA, Chou R. Long-term opioid therapy reconsidered. *Ann Intern Med*. 2011;155(5):325-328.

6. Von Korff M, Saunders K, Thomas Ray G, et al. De facto long-term opioid therapy for noncancer pain [published correction appears in *Clin J Pain*. 2014;30(9):830]. *Clin J Pain*. 2008;24(6):521-527.

7. Calcaterra S, Glanz J, Binswanger IA. National trends in pharmaceutical opioid related overdose deaths compared to other substance related

overdose deaths: 1999-2009. *Drug Alcohol Depend.* 2013;131(3):263-270.

8. National Center for Health Statistics. Number and age-adjusted rates of drug-poisoning deaths involving opioid analgesics and heroin: United States 2000-2014. https://www.cdc.gov/nchs/data/health_policy/AADR_drug_poisoning_involving_OA_Heroin_US_2000-2014.pdf. Published 2014. Accessed May 31, 2016.

9. Edlund MJ, Steffick D, Hudson T, Harris KM, Sullivan M. Risk factors for clinically recognized opioid abuse and dependence among veterans using opioids for chronic non-cancer pain. *Pain.* 2007;129(3):355-362.

10. Morasco BJ, Turk DC, Donovan DM, Dobscha SK. Risk for prescription opioid misuse among patients with a history of substance use disorder. *Drug Alcohol Depend.* 2013;127(1-3):193-199.

11. Tetrault JM, Desai RA, Becker WC, Fiellin DA, Concato J, Sullivan LE. Gender and non-medical use of prescription opioids: results from a national US survey. *Addiction.* 2008;103(2):258-268.

12. Miech R, Johnston L, O'Malley PM, Keyes KM, Heard K. Prescription opioids in adolescence and future opioid misuse. *Pediatrics.* 2015;136(5):e1169-e1177.

13. Li L, Setoguchi S, Cabral H, Jick S. Opioid use for noncancer pain and risk of myocardial infarction amongst adults. *J Intern Med.* 2013;273(5):511-526.

14. Saunders KW, Dunn KM, Merrill JO, et al. Relationship of opioid use and dosage levels to fractures in older chronic pain patients. *J Gen Intern Med.* 2010;25(4):310-315.

15. Miller M, Barber CW, Leatherman S, et al. Prescription opioid duration of action and the risk of unintentional overdose among patients receiving opioid therapy. *JAMA Intern Med.* 2015;175(4):608-615.

16. Edlund MJ, Forman-Hoffman VL, Winder CR, et al. Opioid abuse and depression in adolescents: results from the National Survey on Drug Use and Health. *Drug Alcohol Depend.* 2015;152:131-138.

17. Lovegrove MC, Mathew J, Hamp C, Governale L, Wysowski DK, Budnitz DS. Emergency hospitalizations for unsupervised prescription medication ingestions by young children. *Pediatrics.* 2014;134(4):e1009-e1016.

18. Lovegrove MC, Weidle NJ, Budnitz DS. Trends in emergency department visits for unsupervised pediatric medication exposures, 2004-2013. *Pediatrics.* 2015;136(4):e821-e829.

19. Bond GR, Woodward RW, Ho M. The growing impact of pediatric pharmaceutical poisoning [published correction appears in *J Pediatr.* 2012;160(5):888-889]. *J Pediatr.* 2012;160(2):265-270.e1.

20. Burghardt LC, Ayers JW, Brownstein JS, Bronstein AC, Ewald MB, Bourgeois FT. Adult prescription drug use and pediatric medication exposures and poisonings. *Pediatrics.* 2013;132(1):18-27.

21. Centers for Disease Control and Prevention (CDC). Emergency department visits involving nonmedical use of selected prescription drugs: United States, 2004-2008. *MMWR Morb Mortal Wkly Rep.* 2010;59(23):705-709.

22. Bailey JE, Campagna E, Dart RC; RADARS System Poison Center Investigators. The

underrecognized toll of prescription opioid abuse on young children. *Ann Emerg Med.* 2009;53(4):419-424.

23. Coben JH, Davis SM, Furbee PM, Sikora RD, Tillotson RD, Bossarte RM. Hospitalizations for poisoning by prescription opioids, sedatives, and tranquilizers. *Am J Prev Med.* 2010;38(5):517-524.

24. Rubin R. Analysis reveals large increase in hospitalizations in recent years among older patients prescribed opioids. *JAMA.* 2014;312(16):1621-1623.

25. Meiman J, Tomasallo C, Paulozzi L. Trends and characteristics of heroin overdoses in Wisconsin, 2003-2012. *Drug Alcohol Depend.* 2015;152:177-184.

26. Cicero TJ, Ellis MS, Surratt HL. Effect of abuse-deterrent formulation of OxyContin. *N Engl J Med.* 2012;367(2):187-189.

27. Cicero TJ, Ellis MS. Abuse-deterrent formulations and the prescription opioid abuse epidemic in the United States: lessons learned from OxyContin. *JAMA Psychiatry.* 2015;72(5):424-430.

28. Pollini RA, Banta-Green CJ, Cuevas-Mota J, Metzner M, Teshale E, Garfein RS. Problematic use of prescription-type opioids prior to heroin use among young heroin injectors. *Subst Abuse Rehabil.* 2011;2(1):173-180.

29. Centers for Disease Control and Prevention. Vital signs: risk for overdose from methadone used for pain relief—United States, 1999-2010. Vital Signs Factsheet 2012. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6126a5.htm>. Published July 6, 2012. Accessed June 7, 2016.

30. Centers for Disease Control and Prevention. Opioid painkiller prescribing. Vital Signs Factsheet. <http://www.cdc.gov/vitalsigns/opioid-prescribing/>. Updated July 1, 2014. Accessed November 18, 2015.

31. US Food and Drug Administration. CDER conversation: pediatric pain management options. <http://www.fda.gov/Drugs/NewsEvents/ucm456973.htm>. Updated August 14, 2015. Accessed November 17, 2015.

32. Healthcare Cost and Utilization Project. Introduction to the HCUP KIDS' Inpatient Database (KID). <https://www.hcup-us.ahrq.gov>. Modified September 20, 2016. Accessed June 7, 2016.

33. Agency for Healthcare Research and Quality. Care of children and adolescents in US hospitals. <http://archive.ahrq.gov/data/hcup/factbk4/factbk4.htm>. Modified September 14, 2016. Accessed June 7, 2016.

34. World Health Organization. *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*. Geneva, Switzerland: World Health Organization; 2010.

35. Centers for Disease Control and Prevention. WISQARS (Web-based Injury Statistics Query and Reporting System). <http://www.cdc.gov/injury/wisqars/>. Updated May 4, 2016. Accessed November 18, 2015.

36. Healthcare Cost and Utilization Project. HCUPnet methodology. <http://hcupnet.ahrq.gov/HCUPnet.jsp?GoTo=HCUPnetMethodology>. Published 2014. Accessed May 12, 2016.

37. Healthcare Cost and Utilization Project. Calculating Kids' Inpatient Database (KID) variances. <https://www.hcup-us.ahrq.gov/db/nation/kid/reports/CalculatingKIDVariances.pdf>. Published December 16, 2005. Accessed November 19, 2015.

38. Centers for Disease Control and Prevention NCHS. Bridged-race population estimates: 1990-2011 request. <http://wonder.cdc.gov/bridged-race-v2011.html>. Published 2015. Accessed November 19, 2015.

39. Base SAS. 9.2 Procedures Guide: Statistical Procedures, Third Edition. https://support.sas.com/documentation/cdl/en/statug/63033/HTML/default/viewer.htm#statug_freq_a0000000645.htm. Updated May 19, 2010. Accessed September 20, 2016.

40. Curtin SC, Warner M, Hedegaard H. Increase in suicide in the United States, 1999-2014. *NCHS Data Brief.* 2016;(241):1-8.

41. Pfuntner A, Wier LM, Stocks C. *Most Frequent Conditions in US Hospitals, 2011: Healthcare Cost and Utilization Project (HCUP) Statistical Briefs*. Rockville, MD: Healthcare Cost and Utilization Project; 2006. Statistical brief 62.

42. Unick GJ, Rosenblum D, Mars S, Ciccarone D. Intertwined epidemics: national demographic trends in hospitalizations for heroin- and opioid-related overdoses, 1993-2009. *PLoS One.* 2013;8(2):e54496.

43. Center for Behavioral Health Statistics and Quality. Behavioral health trends in the United States: results from the 2014 National Survey on Drug Use and Health. Health and Human Services publication SMA 15-4927, NSDUH Series H-50. <http://www.samhsa.gov/data>. Published 2015. Accessed March 29, 2016.

44. McCabe SE, Boyd CJ, Cranford JA, Teter CJ. Motives for nonmedical use of prescription opioids among high school seniors in the United States: self-treatment and beyond. *Arch Pediatr Adolesc Med.* 2009;163(8):739-744.

45. McCabe SE, West BT, Boyd CJ. Leftover prescription opioids and nonmedical use among high school seniors: a multi-cohort national study. *J Adolesc Health.* 2013;52(4):480-485.

46. Fortuna RJ, Robbins BW, Caiola E, Joynt M, Halterman JS. Prescribing of controlled medications to adolescents and young adults in the United States. *Pediatrics.* 2010;126(6):1108-1116.

47. DeVries A, Koch T, Wall E, Getchius T, Chi W, Rosenberg A. Opioid use among adolescent patients treated for headache. *J Adolesc Health.* 2014;55(1):128-133.

48. Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain—United States, 2016. *MMWR Recomm Rep.* 2016;65(1):1-49.

49. Olsen Y. The CDC guideline on opioid prescribing: rising to the challenge. *JAMA.* 2016;315(15):1577-1579.

50. Schechter NL, Walco GA. The potential impact on children of the CDC guideline for prescribing opioids for chronic pain: above all, do no harm. *JAMA Pediatr.* 2016;170(5):425-426.

51. de Achaval S, Feudtner C, Palla S, Suarez-Almazor ME. Validation of ICD-9-CM codes for identification of acetaminophen-related emergency department visits in a large pediatric hospital. *BMC Health Serv Res.* 2013;13:72.

52. Califf RM, Woodcock J, Ostroff S. A proactive response to prescription opioid abuse. *N Engl J Med.* 2016;374(15):1480-1485.