

Psychiatric Disorders in Youth in Juvenile Detention

Linda A. Teplin, PhD; Karen M. Abram, PhD; Gary M. McClelland, PhD; Mina K. Dulcan, MD; Amy A. Mericle, PhD

Background: Given the growth of juvenile detainee populations, epidemiologic data on their psychiatric disorders are increasingly important. Yet, there are few empirical studies. Until we have better epidemiologic data, we cannot know how best to use the system's scarce mental health resources.

Methods: Using the Diagnostic Interview Schedule for Children version 2.3, interviewers assessed a randomly selected, stratified sample of 1829 African American, non-Hispanic white, and Hispanic youth (1172 males, 657 females, ages 10-18 years) who were arrested and detained in Cook County, Illinois (which includes Chicago and surrounding suburbs). We present 6-month prevalence estimates by demographic subgroups (sex, race/ethnicity, and age) for the following disorders: affective disorders (major depressive episode, dysthymia, manic episode), anxiety (panic, separation anxiety, overanxious, generalized anxiety, and obsessive-compulsive disorders), psychosis, attention-deficit/hyperactivity disorder, disruptive behavior disorders (oppositional defiant disorder, conduct disorder), and substance use disorders (alcohol and other drugs).

Results: Nearly two thirds of males and nearly three quarters of females met diagnostic criteria for one or more psychiatric disorders. Excluding conduct disorder (common among detained youth), nearly 60% of males and more than two thirds of females met diagnostic criteria and had diagnosis-specific impairment for one or more psychiatric disorders. Half of males and almost half of females had a substance use disorder, and more than 40% of males and females met criteria for disruptive behavior disorders. Affective disorders were also prevalent, especially among females; more than 20% of females met criteria for a major depressive episode. Rates of many disorders were higher among females, non-Hispanic whites, and older adolescents.

Conclusions: These results suggest substantial psychiatric morbidity among juvenile detainees. Youth with psychiatric disorders pose a challenge for the juvenile justice system and, after their release, for the larger mental health system.

Arch Gen Psychiatry. 2002;59:1133-1143

From The Feinberg School of Medicine, Northwestern University Department of Psychiatry and Behavioral Sciences (Drs Teplin, Abram, McClelland, and Dulcan); Psycho-Legal Studies Program (Drs Teplin, Abram, and McClelland); Children's Memorial Hospital (Dr Dulcan); and the School of Social Service Administration, University of Chicago (Dr Mericle), Chicago, Ill.

A GREAT PROPORTION of this country's young people are now involved in the juvenile justice system. In 1999, the Federal Bureau of Investigation estimated that there were 2.5 million arrests of juveniles.¹ In 1997, juvenile courts handled almost 1 800 000 delinquency cases.² On an average day, more than 106 000 youth are in custody in juvenile facilities.³ Almost 60% of detained youth are African American or Hispanic.³ Moreover, recent changes in the laws, such as mandatory penalties for drug crimes and lowering the age that juveniles can be tried as adults, have resulted in more juveniles serving time than ever before. There are currently 163 200 cases per year of juveniles convicted and serving sentences.² Many are incarcerated in adult prisons, which do not have psychi-

atric services designed for juveniles. The number of females in the juvenile justice system is increasing at an even faster rate than the number of males³ and is at an all time high.²

Given the growth of juvenile detainee populations,⁴ epidemiologic data on their psychiatric disorders are increasingly important. Like adult detainees, juvenile detainees with serious mental disorders have a constitutional right (under the Eighth and Fourteenth Amendments) to receive needed treatment.⁵ Mental health professionals believe that providing psychiatric services to juvenile detainees could improve their quality of life and help reduce recidivism.⁶⁻⁸ Until we have better data, we cannot know how best to use the system's scarce mental health resources.^{9,10}

Despite the importance of psychiatric epidemiologic data on juvenile detain-

ees, there are few empirical studies¹⁰ and little consistency in results. Among studies^{7,11-28} published since 1980 (summary table available from authors), rates for affective disorder varied from 2%¹⁵ to 88%.⁷ Rates of substance use disorders ranged from 13%¹⁴ to 88%.⁷ This disparity in findings may be because youth were sampled at various points in the juvenile justice system (eg, at admission, after conviction). In addition, there are 3 methodologic problems.

1. *Biased Samples.* Previous studies¹¹ used disparate exclusion criteria (eg, excluding juveniles with psychotic symptoms, mental retardation, or physical handicaps). Many studies excluded females entirely^{16,21} or sampled too few to analyze them.²⁵
2. *Small Samples.* Some severe disorders have low base rates, between 1% and 4%.^{29,30} Low base rates require large sample sizes to generate reliable estimates.³¹ Some studies sampled too few subjects to generate reliable rates even for the more common disorders.^{18,21}
3. *Problems in Measurement.* Some studies did not specify the diagnostic criteria,¹⁸ used nonstandard or untested instruments,¹⁶ or extracted diagnoses from case records.¹⁷

This study overcomes these methodologic limitations. We have a large, random sample of juvenile detainees and used a reliable measure, version 2.3 of the Diagnostic Interview Schedule for Children (DISC),³² to determine psychiatric diagnoses.

METHODS

PARTICIPANTS AND SAMPLING PROCEDURES

Participants in the Northwestern Juvenile Project included 1829 male and female youth (aged 10-18 years) who were randomly sampled from intake into the Cook County Juvenile Temporary Detention Center (CCJTDC) from November 1995 through June 1998. The sample was stratified by sex, race/ethnicity (African American, non-Hispanic white, Hispanic), age (10-13 years or ≥ 14 years), and legal status (processed as a juvenile or as an adult) to obtain enough participants to compare key subgroups (eg, females, Hispanics, and younger children).

The CCJTDC receives approximately 8500 admissions each year (John Howard Association, unpublished data, 1992) and is used solely for pretrial detention and for offenders sentenced for fewer than 30 days. All detainees younger than 17 years are held at CCJTDC, including youth processed as adults (automatic transfers to adult court). Juveniles up to age 21 years may be detained in the CCJTDC if they are still being prosecuted for an arrest that occurred when they were younger than 17 years.

Like juvenile detainees nationwide, approximately 90% of the CCJTDC detainees are males, and most are racial/ethnic minorities.³ The CCJTDC population is 77.9% African American, 5.6% non-Hispanic white, 16.0% Hispanic, and 0.5% other racial or ethnic groups. The age and offense distributions of CCJTDC detainees are also similar to detained juveniles nationwide.³

We chose the detention center in Cook County (which includes Chicago and surrounding suburbs) for 3 reasons. First, nationwide, most juvenile detainees live in and are detained in urban areas.³³ Second, Cook County is ethnically diverse and

has the third largest Hispanic population in the United States.³⁴ Studying Hispanics is important because they are the largest minority group in the United States³⁵ and they are overrepresented in the justice systems.³ Finally, the detention center's size (daily census of approximately 650 youth and intake of 20 youth per day) ensured that enough participants would be available.

No single site can represent the entire country because jurisdictions may have different options for diversion.^{36,37} Nevertheless, Illinois' criteria for detaining juveniles are similar to the criteria of other states.³⁶ All states allow pretrial detention if the juvenile needs protection, is likely to flee, or is considered a danger to the community.^{36,37}

Detainees were eligible to participate, regardless of their psychiatric morbidity, state of alcohol or other drug intoxication, or fitness to stand trial. Within each stratum, we used a random-numbers table to select names from the CCJTDC's intake log. Throughout the study, we tracked how many participants were still needed to fill each stratum. Project staff sampled the rarest cells first. When more than one participant was available for a stratum, a random-numbers table was used. The final sampling fractions ranged from 0.018 to 0.689. (Additional information on the sample is available from the authors.)

Studying detained youth requires special procedures because they are minors, they are detained, and many do not have a parent or guardian who can provide appropriate consent.³⁸ Project staff approached participants on their units, explained the project, and assured them that anything they told us (except comments implying imminent danger to self or others) would be confidential. Detainees who chose to participate signed an assent form (if they were younger than 18 years) or consent form (if they were 18 years or older). Federal regulations allow parental consent to be waived if the research involves minimal risk (45 CFR [Code of Federal Regulations] 46.116[c], 45 CFR 46.116[d], and 45 CFR 46.408[c]).^{38,39} The Northwestern University Institutional Review Board, the Centers for Disease Control and Prevention Institutional Review Board, and the US Office of Protection From Research Risks waived parental consent. However, as ethicists recommend, we nevertheless tried to contact parents to provide them an opportunity to decline participation and to offer them additional information (45 CFR 46.116[d][4]).^{40,41} Despite repeated attempts to contact the parent or guardian, for 43.8% of participants, none could be found. In lieu of parental consent, assent from the juvenile subject was overseen by a participant advocate who represented the interests of the participants. Federal regulations allow for a participant advocate if parental consent is not feasible (45 CFR 46.116[d]).⁴⁰ Of the 2275 names selected, 4.2% (34 youth and 62 parents or guardians) refused to participate. There were no significant differences in refusal rates by sex, race/ethnicity, or age. Some youth processed as adults (automatic transfers) were counseled by their lawyers to refuse participation; in this stratum, the refusal rate was 7.1% (26 of 368 youth). Twenty-seven youth left the detention center before we could schedule an interview; 312 were not interviewed because they left while we were locating their caretakers for consent. Eleven others were excluded: 9 participants who became physically ill during the interview and could not finish it, 1 participant who was too cognitively impaired to be interviewed, and 1 participant who seemed to be lying. The final sample size was 1829. This sample size allows us to reliably detect disorders (ie, distinguish them from zero) that have a base rate in the general population of 1.0% or greater with a power of 0.80.³¹

Participants were interviewed in a private area, almost always within 2 days of intake. Most interviews lasted 2 to 3 hours, depending on how many symptoms were reported. We used

both male and female interviewers. Female participants were always interviewed by female interviewers. Interviewers were trained for at least a month; most had a master's degree in psychology or an associated field and experience interviewing high-risk youth. One third of our interviewers were fluent in Spanish. We maintained consistency throughout the study by monitoring scripted interviews with mock subjects.

PSYCHIATRIC DIAGNOSES

We used version 2.3 of the DISC,^{32,42} the most recent English and Spanish versions then available. The DISC assesses the presence of disorders in the past 6 months. The DISC is highly structured, contains detailed symptom probes, has acceptable reliability and validity,^{32,43-46} and requires relatively brief training.

Two diagnoses required special management. The psychosis module, a broad symptom screen, does not generate a specific diagnosis. Instead, this module flags participants if they endorse any "possible" or "probable" pathognomonic symptoms or at least 3 nonpathognomonic symptoms. More than one quarter of our participants scored positive on the screen. To be conservative, we counted these participants as psychotic only if (1) their symptoms persisted for at least 1 week; (2) they had not used alcohol, illicit drugs, or medication during this time; and (3) a project clinician (a psychiatrist or clinical psychologist) reviewed the case and judged that the symptoms were "probably indicative of psychosis." Twelve participants met these criteria. Project clinicians also included another 8 participants as psychotic who, although they denied symptoms, appeared to have auditory hallucinations, thought disorders, or delusions during the interview.

Attention-deficit/hyperactivity disorder (ADHD) is difficult to assess via self-report⁴⁷ and is even more challenging to diagnose among delinquent youth.⁴⁸ In addition, *DSM-III-R* requires that symptoms of ADHD be present before the age of 7 years. Age of onset is usually reported by the caretaker. Most of our participants, even if they reported symptoms of ADHD, could not remember when their symptoms began. To avoid underreporting ADHD, we calculated rates in 2 ways: in the conventional manner (requiring that the subject report that symptoms were present before the age of 7 years) and counting the disorder as present regardless of the reported age of onset, as long as the duration criterion was met. (We present only the latter; the former rates are available from the authors.)

We determined rates of disorders in 2 ways. First, as most investigators have done, we used the DISC standard computer algorithms to calculate rates using *DSM-III-R* criteria. We then calculated more conservative (less inclusive) rates for diagnoses that met both *DSM-III-R* criteria and diagnosis-specific impairment criteria, reported by participants.³² Although young people are poor reporters of their own impairment,^{32,49} we calculated these latter rates because recent reviews^{32,50,51} suggest that psychiatric diagnoses are more accurately determined by the presence of both symptoms and functional impairment. (We also examined rates using *DSM-III-R* criteria and a global measure of functional impairment, the Children's Global Assessment Scale.^{52,53} These rates, substantially similar to those reported herein, are available from the authors.)

STATISTICAL ANALYSIS

Because we stratified our sample by sex, race/ethnicity, age, and legal status, we weighted all prevalence estimates to reflect the distributions of these variables in the detention center's population. All reported SEs and tests of significance have been corrected for design characteristics with Taylor series linearization.^{54,55} We used 2-tailed tests; our level of significance for all

Table 1. Unweighted Sample Characteristics*

Characteristic	No. (%) of Participants (N = 1829)
Race/ethnicity	
African American	1005 (54.9)
Non-Hispanic white	296 (16.2)
Hispanic	524 (28.7)
Other	4 (0.2)
Sex	
Male	1172 (64.1)
Female	657 (35.9)
Age, y	
Mean	14.9
Median	15
Mode	16
Specific ages, y	
10	7 (0.4)
11	20 (1.1)
12	87 (4.8)
13	258 (14.1)
14	217 (11.9)
15	498 (27.2)
16	644 (35.2)
17	89 (4.9)
18	9 (0.5)
Education, grade	
≤6th	89 (4.9)
7th	171 (9.3)
8th	306 (16.7)
9th	568 (31.1)
10th	455 (24.9)
11th	172 (9.4)
12th	27 (1.5)
Currently in general equivalency diploma classes	31 (1.7)
Alternative or home schooling	5 (0.3)
Unknown	5 (0.3)
Legal status	
Processed in adult court (automatic transfer)	275 (15.0)
Processed in juvenile court	1554 (85.0)

*Percentages may not sum to 100% because of rounding error.

tests was .05. We report all disorders for males and females separately because combining them masks important differences.

RESULTS

Table 1 presents unweighted demographic characteristics of our sample. **Table 2** provides data showing that nearly two thirds of the males and nearly three quarters of females met diagnostic criteria for 1 or more of the disorders listed. The more conservative estimates using the diagnosis-specific impairment criteria are only slightly lower. We also calculated overall rates excluding conduct disorder because many symptoms are related to delinquent behaviors; **Table 2** also shows overall rates, excluding conduct disorder (with and without diagnosis-specific impairment criteria); rates decreased only slightly.

The most common disorders among both males and females were substance use disorders and disruptive behavior disorders (oppositional defiant disorder and conduct disorder). Half of males and almost half of females met criteria for a substance use disorder, and more than 40% of males and females met criteria for disruptive behavior

Table 2. Six-Month Prevalence and Odds Ratios (ORs) of *DSM-III-R* Diagnoses by Sex With and Without Diagnosis-Specific Impairment Criteria*

Disorder	Male, % (95% CI) (n = 1170)		Female, % (95% CI) (n = 656)		Female-Male ORs (95% CI)	
	Diagnosis	Diagnosis With Impairment	Diagnosis	Diagnosis With Impairment	Diagnosis	Diagnosis With Impairment
Any of the listed disorders	66.3 (61.6-70.7)	63.3 (58.6-67.8)	73.8 (70.1-77.1)	71.2 (67.5-74.7)	1.43 (1.09-1.88)	1.43 (1.10-1.87)
Any except conduct disorder	60.9 (56.2-65.5)	59.7 (54.9-64.3)	70.0 (66.2-73.5)	68.2 (64.4-71.8)	1.49 (1.15-1.94)	1.45 (1.12-1.88)
Any affective disorder	18.7 (15.2-22.8)	16.1 (12.8-20.0)	27.6 (23.6-32.0)	22.9 (19.0-27.2)	1.66 (1.20-2.29)	1.55 (1.09-2.20)
Major depressive episode	13.0 (10.0-16.6)	11.0 (8.3-14.5)	21.6 (17.8-25.9)	18.9 (15.2-23.2)	1.85 (1.27-2.70)	1.88 (1.25-2.82)
Dysthymia	12.2 (9.3-15.8)	9.9 (7.3-13.2)	15.8 (13.1-18.8)	12.5 (10.2-15.3)	1.34 (0.93-1.95)	1.31 (0.87-1.96)
Manic episode	2.2 (1.1-4.3)	2.0 (1.0-4.1)	1.8 (1.0-3.2)	1.2 (0.6-2.4)	0.81 (0.33-1.99)	0.58 (0.21-1.63)
Psychotic disorders	1.0 (0.4-2.6)	...	1.0 (0.5-2.1)	...	0.98 (0.30-3.25)	...
Any anxiety disorder	21.3 (17.6-25.6)	20.7 (17.0-24.9)	30.8 (27.2-34.6)	28.9 (25.5-32.7)	1.64 (1.22-2.20)	1.56 (1.16-2.10)
Panic disorder	0.3 (0.1-0.6)	0.1 (0.0-0.4)	1.5 (0.8-2.7)	1.0 (0.5-2.0)	5.65 (2.04-15.65)	8.13 (2.01-32.85)
Separation anxiety disorder	12.9 (9.9-16.5)	10.8 (8.1-14.2)	18.6 (15.7-21.9)	16.3 (13.6-19.4)	1.55 (1.08-2.21)	1.61 (1.10-2.34)
Overanxious disorder	6.7 (4.6-9.7)	5.9 (4.0-8.7)	12.3 (9.9-15.1)	11.5 (9.2-14.2)	1.95 (1.23-3.10)	2.06 (1.27-3.35)
Generalized anxiety disorder	7.1 (4.9-10.2)	6.4 (4.3-9.4)	7.3 (5.6-9.6)	6.8 (5.1-9.0)	1.03 (0.63-1.69)	1.07 (0.64-1.79)
Obsessive-compulsive disorder	8.3 (6.1-11.3)	...	10.6 (8.4-13.2)	...	1.31 (0.86-2.00)	...
Attention-deficit/hyperactivity disorder†	16.6 (13.3-20.5)	11.2 (8.5-14.6)	21.4 (18.4-24.8)	16.4 (13.7-19.5)	1.37 (0.99-1.89)	1.55 (1.07-2.25)
Any disruptive behavior disorder	41.4 (36.8-46.2)	31.4 (27.2-36.0)	45.6 (41.4-49.8)	38.0 (33.9-42.2)	1.19 (0.92-1.53)	1.33 (1.02-1.75)
Oppositional-defiant disorder	14.5 (11.4-18.2)	12.6 (9.8-16.2)	17.5 (14.7-20.6)	15.1 (12.5-18.1)	1.25 (0.89-1.76)	1.23 (0.86-1.76)
Conduct disorder	37.8 (33.3-42.6)	24.3 (20.5-28.5)	40.6 (36.5-44.8)	28.5 (24.6-32.8)	1.12 (0.86-1.46)	1.24 (0.92-1.67)
Any substance use disorder	50.7 (45.9-55.5)	...	46.8 (42.6-51.1)	...	0.86 (0.66-1.11)	...
Alcohol use disorder	25.9 (21.9-30.4)	...	26.5 (22.6-30.9)	...	1.03 (0.76-1.40)	...
Marijuana use disorder	44.8 (40.1-49.6)	...	40.5 (36.8-44.4)	...	0.84 (0.65-1.08)	...
Other substance use disorder	2.4 (1.7-3.4)	...	6.9 (4.1-11.4)	...	3.00 (1.57-5.74)	...
Both alcohol and other drug use disorders	20.7 (17.0-24.9)	...	20.9 (18.0-24.2)	...	1.01 (0.75-1.38)	...

*CI indicates confidence interval. Ellipses indicate that diagnosis and diagnosis with impairment are identical because the diagnostic criteria for psychotic disorders, obsessive-compulsive disorder, and substance use disorders include impairment.

†Attention-deficit/hyperactivity disorder is reported without the criterion of onset before age 7 years because caretaker information is not available and self-report of symptoms before age 7 years is unreliable.

disorders. Rates of disorder using diagnosis-specific impairment criteria for conduct disorder are more than 10% lower than conduct disorder without impairment. More than one fourth of females and almost one fifth of males met criteria for 1 or more affective disorders.

Table 2 also reports the female-male odds ratios. Odds ratios greater than 1.0 indicate that females had higher odds of having the disorder than males; those less than 1.0 show that females had lower odds of having the disorder. Females had significantly higher odds than males of having any disorder, any disorder except conduct disorder, any affective disorder, major depressive episode, any anxiety disorder, panic disorder, separation anxiety disorder, overanxious disorder, and substance use disorder other than alcohol or marijuana.

Table 3 and **Table 4** show the prevalence rates of disorders for males and females by race/ethnicity. Cases in these and subsequent tables met *DSM-III-R* criteria. (Tables of disorders that meet diagnosis-specific impairment criteria also are available from the authors.) We report protected tests of significance for specific racial/ethnic contrasts only when the overall test was significant. Table 3 shows that among males, non-Hispanic whites had the highest rates of many disorders and African Americans the lowest. Specifically, compared with African Americans, non-Hispanic whites had significantly higher rates of any disorder, any disorder except conduct disorder, any disruptive behavior disorder, conduct disorder, any substance use disorder, and substance use disorder other than

alcohol or marijuana. The only disorder for which African Americans had significantly higher rates than non-Hispanic whites was separation anxiety disorder. Hispanics had significantly higher rates than non-Hispanic whites of any anxiety disorder and separation anxiety disorder. Hispanics had higher rates than African Americans of panic disorder, obsessive-compulsive disorder, and substance use other than alcohol or marijuana disorders. Non-Hispanic whites had higher rates than Hispanics of any disorder, any disruptive behavior disorder, conduct disorder, and substance use disorder other than alcohol or marijuana.

Table 4 compares rates by race/ethnicity for females. Non-Hispanic white females had significantly higher rates than African Americans of any disorder, any disorder except conduct disorder, any disruptive behavior disorder, conduct disorder, and all substance use disorders and higher rates than Hispanics of any disorder except conduct disorder. Hispanic females had higher rates of generalized anxiety disorder than either African American or white females. Compared with African Americans, Hispanic females had higher rates of all disruptive behavior disorders, conduct disorder, alcohol use disorder, substance use disorder other than alcohol or marijuana, and both alcohol and drug use disorder.

Table 5 and **Table 6** show the prevalence rates of disorders for males and females by age. Among males, Table 5 indicates that the youngest age group had the lowest rates of many disorders. They had significantly lower rates than both older age groups of any disorder, any dis-

Table 3. Six-Month Prevalence of *DSM-III-R* Diagnoses for Males by Race/Ethnicity*

Disorder	Prevalence, % (95% CI)			Overall Significance	Protected Tests†
	African American (n = 574)	Non-Hispanic White (n = 207)	Hispanic (n = 386)		
Any of the listed disorders	64.6 (58.8-69.9)	82.0 (76.2-86.7)	70.4 (63.3-76.7)	<.001	White > African American; white > Hispanic
Any except conduct disorder	59.4 (53.5-65.0)	72.9 (66.5-78.6)	65.3 (58.1-71.9)	.009	White > African American
Any affective disorder	18.6 (14.4-23.6)	13.8 (9.6-19.5)	21.5 (15.3-29.3)	.19	
Major depressive episode	12.5 (9.1-17.0)	9.5 (6.0-14.6)	16.6 (10.8-24.7)	.20	
Dysthymia	12.2 (8.8-16.7)	9.5 (6.1-14.5)	13.3 (8.4-20.6)	.53	
Manic episode	2.5 (1.2-5.2)	0.5 (0.1-3.7)	1.4 (0.6-3.2)	.27	
Psychotic disorders	1.0 (0.3-3.2)	2.6 (1.1-6.2)	0.7 (0.2-2.6)	.19	
Any anxiety disorder	20.9 (16.5-26.1)	14.4 (10.1-20.2)	25.5 (18.7-33.7)	.046	Hispanic > white
Panic disorder	0.1 (0.0-0.4)	0.5 (0.1-3.7)	1.0 (0.3-3.1)	.04	Hispanic > African American
Separation anxiety disorder	12.7 (9.3-17.2)	5.9 (3.3-10.3)	15.5 (9.8-23.6)	.02	African American > white; Hispanic > white
Overanxious disorder	6.9 (4.4-10.7)	2.9 (1.3-6.6)	7.0 (3.6-13.0)	.16	
Generalized anxiety disorder	7.5 (4.8-11.4)	2.5 (1.0-5.9)	7.2 (3.7-13.3)	.08	
Obsessive-compulsive disorder	6.5 (4.2-10.0)	9.3 (5.8-14.4)	17.0 (10.7-25.9)	.01	Hispanic > African American
Attention-deficit/hyperactivity disorder‡	17.0 (13.0-21.9)	20.9 (15.8-27.3)	13.7 (9.4-19.5)	.18	
Any disruptive behavior disorder	39.8 (34.2-45.7)	60.3 (53.3-66.9)	43.3 (36.1-50.8)	<.001	White > African American; white > Hispanic
Oppositional-defiant disorder	14.4 (10.7-19.1)	19.4 (14.4-25.6)	13.6 (9.3-19.5)	.23	
Conduct disorder	35.6 (30.1-41.5)	59.9 (53.0-66.5)	41.7 (34.5-49.2)	<.001	White > African American; white > Hispanic
Any substance use disorder	49.1 (43.2-55.0)	62.6 (55.7-69.0)	55.4 (47.8-62.7)	.01	White > African American
Alcohol use disorder	24.6 (19.8-30.2)	30.1 (24.0-36.9)	30.8 (24.1-38.5)	.28	
Marijuana use disorder	44.4 (38.6-50.4)	53.8 (46.8-60.6)	45.4 (38.0-52.9)	.11	
Other substance use disorder	0.5 (0.1-2.8)	21.1 (15.9-27.4)	6.0 (3.9-9.1)	<.001	White > African American; white > Hispanic; Hispanic > African American
Both alcohol and other drug use disorders	20.4 (16.0-25.7)	24.0 (18.5-30.6)	21.7 (16.5-28.0)	.65	

*CI indicates confidence interval. Two cases of "other" race/ethnicity are excluded from this table.

†Protected tests are performed only if the α for the overall test is less than .05.

‡Attention-deficit/hyperactivity disorder is reported without the criterion of onset before the age of 7 years because caretaker information is not available and self-report of symptoms before the age of 7 years is unreliable.

order except conduct disorder, generalized anxiety disorder, and all the substance use disorders. The 14- to 15-year-old group had higher rates of psychotic disorders than the 16 years or older age group.

Table 6 shows somewhat different patterns of disorder for females. The oldest age group had significantly lower rates of oppositional defiant disorder than the younger age groups. The youngest age group had significantly lower rates of any substance use disorder and marijuana use disorder than either of the older age groups.

COMMENT

Our study shows that youth with psychiatric disorders pose a challenge for the juvenile justice system and, after their release, for the larger mental health system. Even after excluding conduct disorder, nearly 60% of male juvenile detainees and more than two thirds of females met diagnostic criteria and had diagnosis-specific impairment for one or more psychiatric disorders.

These rates may underestimate the true prevalence among youth entering the juvenile justice system for 2 reasons. First, our sample included only detainees; it excluded youth who were not detained because their charges were less serious, they were immediately released, or they

were referred directly into the mental health system. Second, underreporting of symptoms and impairments by youth is common, especially for disruptive behavior disorders.⁴⁷

It is difficult to compare our findings with studies of general population youth because rates vary widely, depending on the sample, the method, the source of data (participant or collaterals), and whether functional impairment was required for diagnosis.⁵⁰ Despite these differences, our overall rates are substantially higher than the median rate reported in a major review article (15%)⁵⁰ and other more recent investigations: the Great Smoky Mountains Study (20.3%),⁵⁶ the Virginia Twin Study of Adolescent Behavioral Development (142 cases per 1000 persons),⁵⁷ the Methods for the Epidemiology of Child and Adolescent Mental Disorders (6.1%),³² and the Miami-Dade County Public School Study (38%).⁵⁸ We are especially concerned about the high rates of depression and dysthymia among detained youth (17.2% of males, 26.3% of females), which are also higher than general population rates.^{51,56-61} Depressive disorders are difficult to detect (and treat) in the chaos of the corrections milieu. Overall, our prevalence rates are comparable to rates in other high-risk populations (eg, maltreated or runaway youth).^{62,63}

Table 4. Six-Month Prevalence of *DSM-III-R* Diagnoses for Females by Race/Ethnicity*

Disorder	Prevalence, % (95% CI)			Overall Significance	Protected Tests†
	African American (n = 430)	Non-Hispanic White (n = 89)	Hispanic (n = 136)		
Any of the listed disorders	70.9 (66.4-75.0)	86.1 (77.1-92.0)	75.9 (67.9-82.5)	.01	White > African American
Any except conduct disorders	67.4 (62.8-71.6)	83.9 (74.6-90.3)	69.5 (61.2-76.7)	.01	White > African American; white > Hispanic
Any affective disorder	26.2 (22.2-30.5)	23.4 (15.8-33.4)	28.7 (21.8-36.9)	.68	
Major depressive episode	19.7 (16.2-23.7)	19.0 (12.1-28.5)	22.8 (16.5-30.5)	.70	
Dysthymia	15.5 (12.4-19.2)	17.9 (11.2-27.3)	17.2 (11.8-24.5)	.80	
Manic episode	1.9 (0.9-3.7)	1.1 (0.2-7.5)	2.1 (0.7-6.4)	.85	
Psychotic disorders	0.9 (0.4-2.5)	0.0	2.1 (0.7-6.3)	.29‡	
Any anxiety disorder	31.2 (27.0-35.8)	30.0 (21.4-40.3)	32.6 (25.2-40.9)	.92	
Panic disorder	0.9 (0.4-2.5)	3.4 (1.1-10.0)	2.8 (1.0-7.1)	.17	
Separation anxiety disorder	18.9 (15.5-22.9)	14.5 (8.6-23.4)	21.7 (15.5-29.4)	.41	
Overanxious disorder	12.5 (9.7-16.0)	11.1 (6.1-19.5)	13.2 (8.4-20.1)	.90	
Generalized anxiety disorder	6.6 (4.6-9.4)	4.4 (1.7-11.3)	13.1 (8.4-19.9)	.03	Hispanic > African American; Hispanic > white
Obsessive-compulsive disorder	10.3 (7.8-13.6)	12.4 (7.0-21.1)	10.6 (6.5-16.9)	.84	
Attention-deficit/hyperactivity disorder§	20.0 (16.5-24.1)	22.2 (14.7-32.0)	29.3 (22.2-37.5)	.08	
Any disruptive behavior disorder	39.4 (34.9-44.1)	61.6 (51.0-71.1)	56.5 (47.9-64.6)	<.001	White > African American; Hispanic > African American
Oppositional-defiant disorder	15.8 (12.7-19.6)	17.8 (11.1-27.1)	26.2 (19.5-34.3)	.03	Hispanic > African American
Conduct disorder	34.3 (29.9-38.9)	58.9 (48.3-68.7)	50.2 (41.8-58.6)	<.001	White > African American; Hispanic > African American
Any substance use disorder	42.3 (37.6-47.1)	61.9 (51.2-71.6)	51.7 (43.1-60.1)	.002	White > African American
Alcohol use disorder	21.2 (17.5-25.3)	39.2 (29.5-49.9)	34.0 (26.4-42.5)	<.001	White > African American; Hispanic > African American
Marijuana use disorder	37.8 (33.3-42.5)	53.4 (42.9-63.6)	44.7 (36.3-53.3)	.02	White > African American
Other substance use disorder	0.9 (0.4-2.5)	20.0 (12.9-29.6)	14.7 (9.7-21.5)	<.001	White > African American; Hispanic > African American
Both alcohol and other drug use disorders	17.2 (13.9-21.1)	35.1 (25.7-45.8)	28.3 (21.2-36.7)	<.001	White > African American; Hispanic > African American

*CI indicates confidence interval. Two cases of "other" race/ethnicity are excluded from this table.

†Protected tests are performed only if the α for the overall test is less than .05.

‡Test computed with 1 *df* because of empty cells.

§Attention-deficit/hyperactivity disorder is reported without the criterion of onset before the age of 7 years because caretaker information is not available and self-report of symptoms before the age of 7 years is unreliable.

Our data highlight an important paradox regarding race/ethnicity. More than half of the youth in our juvenile justice system are African American or Hispanic. Therefore, most detained youth with psychiatric disorders are minorities. The prevalence, however, of many disorders is highest among non-Hispanic whites. Thus, white youth in the juvenile justice system may, on average, be more dysfunctional (have greater psychiatric morbidity) than minority youth.

Females had higher rates than males of many psychiatric disorders, including major depressive episode, some anxiety disorders, and "other substance use disorders" (eg, cocaine and hallucinogens). Our findings confirm those of prior studies of adult female detainees and females with conduct disorders, which found that females have higher rates of psychiatric disorders than males.^{64,65}

Overall, the youngest age group (≤ 13 years) had the lowest prevalence rates of most disorders, confirming studies^{57,66-68} of general population youth. Many youth in the juvenile justice system may develop new or additional disorders as they age.

LIMITATIONS

Our study provides only a "snapshot" of our participants' psychiatric disorders immediately after arrest and

detention. We cannot know whether mental disorder causes delinquency, increases the likelihood of arrest and detention, or is merely a frequent trait among delinquent youth. Some symptoms could be a reaction to incarceration. Moreover, our rates might differ somewhat if we had been able to use *DSM-IV* instead of *DSM-III-R* criteria. Our findings, drawn from only one site, may pertain only to youth in urban detention centers with similar demographic composition. Finally, because it was not feasible to interview caretakers, our data are subject to the limitations of self-report. Despite these limitations, our study has important implications for research on delinquent youth and mental health policy.

FUTURE RESEARCH

We suggest 3 directions for future research.

1. *Studies of Patterns and Sequences of Comorbidity.* Examining comorbidity is critical because it is so prevalent among juveniles in the general population,^{69,70} adult jail detainees,⁷¹ and adults who have high arrest rates, such as substance abusers,⁷² young, long-term psychiatric patients,⁷³ and homeless, mentally ill persons.⁷⁴ Moreover, studies⁷¹ of adults suggest that juveniles with comorbid disorders may be especially vulnerable to arrest, particularly if they are poor and

Table 5. Six-Month Prevalence of *DSM-III-R* Diagnosis for Males by Age*

Disorder	Prevalence, % (95% CI)			Overall Significance	Protected Tests†
	Age ≤13 Years (n = 315)	Age 14 and 15 Years (n = 361)	Age ≥16 Years (n = 494)		
Any of the listed disorders	52.7 (46.5-58.8)	68.0 (60.3-74.8)	67.3 (60.3-73.7)	.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Any except conduct disorder	44.9 (38.9-51.0)	63.4 (55.6-70.6)	61.8 (54.7-68.5)	<.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Any affective disorder	13.0 (9.4-17.6)	21.2 (15.4-28.4)	17.7 (12.9-23.7)	.09	
Major depressive episode	7.5 (4.9-11.4)	14.8 (10.0-21.5)	12.4 (8.5-17.8)	.06	
Dysthymia	7.3 (4.7-11.3)	14.5 (9.7-21.1)	11.2 (7.4-16.4)	.08	
Manic episode	1.6 (0.7-4.0)	2.6 (0.9-7.2)	2.0 (0.7-5.1)	.80	
Psychotic disorders	0.0	2.1 (0.7-6.0)	0.3 (0.2-0.8)	.01‡	14 and 15 years > 16 years and older
Any anxiety disorder	17.7 (13.6-22.9)	23.0 (16.9-30.4)	20.6 (15.5-26.7)	.42	
Panic disorder	0.8 (0.2-3.3)	0.1 (0.0-0.9)	0.3 (0.1-0.9)	.25	
Separation anxiety disorder	10.0 (6.9-14.3)	14.5 (9.7-21.1)	12.0 (8.1-17.5)	.40	
Overanxious disorder	4.8 (2.8-8.0)	5.1 (2.6-9.9)	8.4 (5.1-13.5)	.25	
Generalized anxiety disorder	1.3 (0.5-3.4)	5.9 (3.1-11.0)	9.2 (5.8-14.4)	.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Obsessive-compulsive disorder	6.0 (3.7-9.7)	9.4 (5.7-15.0)	7.8 (4.9-12.2)	.43	
Attention-deficit/hyperactivity disorder§	12.5 (9.1-16.9)	20.9 (15.1-28.0)	13.8 (9.7-19.2)	.06	
Any disruptive behavior disorder	32.9 (27.5-38.8)	43.5 (35.9-51.3)	41.2 (34.5-48.2)	.06	
Oppositional-defiant disorder	10.7 (7.5-14.9)	18.2 (12.8-25.1)	12.1 (8.3-17.3)	.08	
Conduct disorder	30.8 (25.6-36.6)	41.1 (33.6-49.1)	36.4 (30.0-43.3)	.10	
Any substance use disorder	28.3 (23.1-34.0)	51.3 (43.5-59.1)	54.4 (47.3-61.3)	<.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Alcohol use disorder	12.9 (9.5-17.4)	25.6 (19.3-33.0)	28.7 (22.8-35.4)	<.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Marijuana use disorder	25.1 (20.3-30.5)	46.9 (39.1-54.8)	46.8 (39.8-53.9)	<.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Other substance use disorder	0.8 (0.4-1.7)	2.5 (1.2-5.0)	2.6 (1.9-3.6)	.01	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Both alcohol and other drug use disorders	10.2 (7.2-14.3)	21.5 (15.7-28.7)	22.0 (16.7-28.3)	<.001	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger

*CI indicates confidence interval.

†Protected tests are performed only if the α for the overall test is less than .05.

‡Test computed with 1 *df* because of empty cells.

§Attention-deficit/hyperactivity disorder is reported without the criterion of onset before the age of 7 years because caretaker information is not available and self-report of symptoms before the age of 7 years is unreliable.

cannot afford treatment. Data on the sequences of comorbidity would help provide the foundation for innovative treatments and tailor services for special populations such as females and minorities.

2. *Studies of Females in the Juvenile Justice System.* Females are increasingly arrested for crimes against persons and violent crimes⁷⁵ and make up an increasingly large proportion of delinquent youth.^{1,2} Prior studies^{76,77} of youth with conduct disorders (many of whom will become delinquent) suggest that females have greater persistence of emotional disorder and worse outcomes than males. Moreover, their problem behaviors often persist beyond adolescence. As they age, they may become suicidal, addicted to alcohol or other drugs, enmeshed in violent relationships, and unable to care for their children.^{64,76} De-

linquent females also engage in sexual activity at an earlier age than nonoffenders, placing them at greater risk for unwanted pregnancy and human immunodeficiency virus.⁷⁸ Understanding psychiatric morbidity and associated risk factors among delinquent females could help us to improve treatment and reduce the cycle of disorder and dysfunction.

3. *Longitudinal Studies.* Many youth in the juvenile justice population may develop new disorders as they age. Risk factors for the development of disorders⁷⁹ are common among delinquent youth, including physical and sexual abuse, a troubled family environment, parental substance abuse, poverty, poor education, neighborhood disintegration, and neglect.⁸⁰⁻⁸⁴ Delinquent youth have few protective factors to offset these risks.⁸⁵ Thus, most youth in the juvenile justice system are at

Table 6. Six-Month Prevalence of *DSM-III-R* Diagnosis for Females by Age*

Disorder	Prevalence, % (95% CI)			Overall Significance	Protected Tests†
	Age ≤13 Years (n = 56)	Age 14 and 15 Years (n = 353)	Age ≥16 Years (n = 247)		
Any of the listed disorders	66.7 (53.2-77.9)	72.2 (67.2-76.7)	77.6 (71.6-82.7)	.18	
Any except conduct disorder	64.7 (51.2-76.2)	67.4 (62.2-72.1)	74.8 (68.6-80.2)	.13	
Any affective disorder	20.7 (12.0-33.3)	27.9 (23.5-32.9)	28.8 (21.2-37.8)	.50	
Major depressive episode	13.0 (6.5-24.2)	21.6 (17.6-26.3)	23.4 (16.0-32.9)	.27	
Dysthymia	10.4 (4.7-21.4)	15.6 (12.2-19.8)	17.2 (12.8-22.6)	.46	
Manic episode	3.9 (1.0-14.4)	1.4 (0.6-3.3)	1.9 (0.8-4.7)	.45	
Psychotic disorders	0.0	0.6 (0.2-2.5)	1.8 (0.7-4.3)	.21‡	
Any anxiety disorder	26.6 (16.7-39.7)	32.6 (27.8-37.7)	29.2 (23.4-35.7)	.55	
Panic disorder	1.9 (0.3-12.4)	1.7 (0.8-3.6)	1.0 (0.3-3.2)	.75	
Separation anxiety disorder	18.1 (10.0-30.6)	19.7 (15.8-24.2)	17.2 (12.9-22.7)	.77	
Overanxious disorder	7.1 (2.7-17.7)	13.8 (10.5-17.8)	11.4 (7.9-16.1)	.34	
Generalized anxiety disorder	3.8 (1.0-14.1)	7.1 (4.9-10.3)	8.4 (5.5-12.7)	.51	
Obsessive-compulsive disorder	10.4 (4.7-21.5)	11.8 (8.8-15.7)	8.8 (5.8-13.1)	.51	
Attention-deficit/hyperactivity disorder§	26.6 (16.6-39.7)	22.7 (18.6-27.4)	18.5 (14.0-24.0)	.30	
Any disruptive behavior disorder	44.7 (32.2-57.9)	50.0 (44.8-55.2)	39.6 (32.0-47.8)	.11	
Oppositional-defiant disorder	30.5 (19.9-43.6)	20.2 (16.4-24.7)	10.7 (7.3-15.2)	<.001	13 years and younger > 16 years and older; 14 and 15 years > 16 years and older
Conduct disorder	33.0 (22.0-46.3)	45.3 (40.2-50.5)	35.7 (28.1-44.2)	.06	
Any substance use disorder	30.5 (19.9-43.7)	45.8 (40.6-51.2)	52.0 (44.5-59.4)	.02	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Alcohol use disorder	16.7 (9.1-28.6)	25.4 (21.1-30.3)	30.3 (22.7-39.2)	.16	
Marijuana use disorder	24.8 (15.3-37.5)	41.3 (36.2-46.6)	43.3 (37.1-49.7)	.04	14 and 15 years > 13 years and younger; 16 years and older > 13 years and younger
Other substance use disorder	5.9 (2.2-14.9)	5.3 (3.5-7.8)	9.5 (3.7-22.2)	.52	
Both alcohol and other drug use disorders	11.5 (5.5-22.5)	21.8 (17.7-26.4)	22.0 (17.2-27.6)	.20	

*CI indicates confidence interval.

†Protected tests are performed only if the α for the overall test is less than .05.

‡Test computed with 1 *df* because of empty cells.

§Attention-deficit/hyperactivity disorder is reported without the criterion of onset before the age of 7 years because caretaker information is not available and self-report of symptoms before the age of 7 years is unreliable.

great risk for psychiatric disorders, problem behaviors, and even early death.^{86,87} Longitudinal studies are needed to examine why some delinquent youth develop new psychiatric disorders and others do not, to investigate protective factors, and to determine how vulnerability and risk differ by key variables such as sex and race/ethnicity. We are now collecting longitudinal data on our participants. Future articles will address persistence and change in psychiatric disorders (including onset, remission, and recurrence), comorbidity, associated functional impairments, and the risk and protective factors related to these disorders and impairments.

IMPLICATIONS FOR MENTAL HEALTH POLICY

Advocacy groups, researchers, and public policy experts believe that the juvenile justice system has become the only alternative for many poor and minority youth with psychiatric disorders.⁸⁸⁻⁹² Many states have imposed more severe sanctions for delinquent youth and transfer increasing numbers of juveniles to adult court,⁹³⁻⁹⁵ policies that disproportionately affect minority youth.^{94,96} In addition, 2 recent changes in public health policy may

have inadvertently contributed to the criminalization of youth with mental disorders.

1. *Welfare Reform.* Welfare reform has disrupted Medicaid benefits for millions of children who need treatment.^{97,98} Medicaid enables many youth to receive psychiatric treatment.⁹⁹ Many parents who left welfare to go to work found their new jobs did not provide insurance or, when available, they could not afford copayments.^{100,101} The State Children's Health Insurance Program, designed to offset the loss of Medicaid, did not fulfill its intended purpose.^{98,102} Moreover, welfare reform has not substantially decreased poverty¹⁰³; many poor children have become even poorer.¹⁰⁴ Poor children are vulnerable to poor outcomes,¹⁰⁵ including involvement with the juvenile justice system.
2. *Managed Care.* Managed care now dominates the private insurance industry⁹² and increasingly controls public insurance benefits, such as Medicaid.^{106,107} Many disorders prevalent among delinquent youth, such as conduct disorder, ADHD, and substance use disorders, are often not covered or have restrictive treatment guidelines.¹⁰⁸ As the public health system re-

duces services, youth with psychiatric disorders may increasingly fall through the cracks into the juvenile justice system.¹⁰⁹

These changes (welfare reform and managed care) have the most serious consequences for poor and minority children, groups overrepresented in the juvenile justice system. Our findings are even more sobering because the prevalence of psychosocial problems among youth seems to be increasing.^{110,111} The US Surgeon General reports that the unmet need for services is as high now as it was 20 years ago.¹¹² Even youth who are insured often cannot obtain treatment because few child and adolescent psychiatrists practice in poor and minority neighborhoods.^{113,114}

The juvenile justice system is not equipped to provide adequate mental health services for the large numbers of detainees with psychiatric disorders.^{115,116} Although the mental health needs of youth in the juvenile justice system have been given much attention recently,^{10,117,118} there are still few empirical studies of the effectiveness of treatment and outcomes.¹⁰ This omission is critical. We need research to guide mental health policy and to understand the complex interplay among the many systems—primary care, mental health, education, child welfare, and justice—that treat delinquent youth.

Submitted for publication October 11, 2001; final revision received March 7, 2002; accepted March 12, 2002.

This work was supported by National Institute of Mental Health (NIMH) grants R01MH54197 and R01MH59463, and grant 1999-JE-FX-1001 from the Office of Juvenile Justice and Delinquency Prevention. Major funding was also provided by the National Institute on Drug Abuse (Bethesda, Md), the Center for Mental Health Services (Rockville, Md), the Centers for Disease Control and Prevention (CDC) National Center for HIV, STD, and TB Prevention (Atlanta, Ga), the National Institute on Alcohol Abuse and Alcoholism (Bethesda), the National Institutes of Health (NIH) Office of Research on Women's Health (Bethesda), the Center for Substance Abuse Prevention (Rockville, Md), the NIH Center on Minority Health and Health Disparities (Bethesda), NIH Office of Rare Diseases (Bethesda), CDC National Center on Injury Prevention and Control (Atlanta), the Center for Substance Abuse Treatment, The William T. Grant Foundation (New York, NY), and The Robert Wood Johnson Foundation (Princeton, NJ). Additional funds were provided by The John D. and Catherine T. MacArthur Foundation (Chicago), the Open Society Institute (New York), and The Chicago Community Trust. We thank all our agencies for their collaborative spirit and steadfast support.

Many more people than the authors contributed to this project. From the NIMH, Ann Hohmann, PhD, and Kimberly Hoagwood, PhD, provided technical assistance and moral support that went beyond the call of duty; Eve Moscicki, ScD, and Heather Ringeisen, PhD, critiqued earlier versions of the article; Grayson Norquist, MD, and Delores Parron, PhD (now at NIH), provided steadfast support throughout. Celia Fisher, PhD, guided our human subjects procedures. We thank all project staff, especially Amy E. Lansing, PhD, for supervising the data collection. We also

thank Laura Coats, our expert editor and research assistant, and Kate Elkington for her meticulous library work. We also greatly appreciate the cooperation of everyone working in the Cook County systems, especially David H. Lux, our project liaison. Without the county's cooperation, this study would not have been possible. Finally, we thank our participants for their time and willingness to participate.

Corresponding author and reprints: Linda A. Teplin, PhD, Psycho-Legal Studies Program, Department of Psychiatry and Behavioral Sciences, The Feinberg School of Medicine, Northwestern University, 710 N Lake Shore Dr, Suite 900, Chicago, IL 60611 (e-mail: psycho-legal@northwestern.edu).

REFERENCES

1. Snyder HN. *Juvenile Arrests 1999*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 2000.
2. Puzanhera C, Stahl AL, Finnegan TA, Snyder HN, Poole RS, Tierney N. *Juvenile Court Statistics 1997*. Washington DC: Office of Juvenile Justice and Delinquency Prevention; 2000.
3. Snyder HN, Sickmund M. *Juvenile Offenders and Victims: 1999 National Report*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 1999.
4. Porter G. *Detention in Delinquency Cases, 1988-1997*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 2000.
5. Costello JC, Jameson EJ. Legal and ethical duties of health care professionals to incarcerated children. *J Leg Med*. 1987;8:191-263.
6. Dembo R, Schmeidler J, Pacheco K, Cooper S, Williams LW. The relationships between youth's identified substance use, mental health or other problems at a juvenile assessment center and their referrals to needed services. *J Child Adolesc Subst Abuse*. 1997;6:23-54.
7. Timmons-Mitchell J, Brown C, Schulz SC, Webster SE, Underwood LA, Semple WE. Comparing the mental health needs of female and male incarcerated juvenile delinquents. *Behav Sci Law*. 1997;15:195-202.
8. McCord J, Widom CS, Crowell NA, eds. *Juvenile Crime, Juvenile Justice*. Washington, DC: National Academy Press; 2001.
9. General Accounting Office. *Mentally Ill Inmates: Better Data Would Help Determine Protection and Advocacy Needs*. Washington, DC: General Accounting Office; 1991.
10. Cocozza JJ, Skowrya KR. Youth with mental health disorders: issues and emerging responses. *Juvenile Justice*. 2000;7:3-13.
11. Chiles JA, Miller ML, Cox GB. Depression in an adolescent delinquent population. *Arch Gen Psychiatry*. 1980;37:1179-1184.
12. Miller ML, Chiles JA, Barnes VE. Suicide attempters within a delinquent population. *J Consult Clin Psychol*. 1982;50:491-498.
13. McManus M, Alessi NE, Grapentine WL, Brickman A. Psychiatric disturbance in serious delinquents. *J Am Acad Child Adolesc Psychiatry*. 1984;23:602-615.
14. McManus M, Brickman A, Alessi NE, Grapentine WL. Borderline personality in serious delinquents. *Comp Psychiatry*. 1984;25:446-454.
15. Cocozza JJ, Ingalls RP. *Out of Home Care*. Albany: New York State Council on Children and Families; 1984.
16. Hollander HE, Turner FD. Characteristics of incarcerated delinquents: relationship between development disorders, environmental and family factors, and patterns of offense and recidivism. *J Am Acad Child Adolesc Psychiatry*. 1985;24:221-226.
17. Friedman RM, Kutash K. *Mad, Bad, Sad, Can't Add: Florida Adolescent and Child Treatment Study (FACTS)*. Tampa: University of South Florida, Florida Mental Health Institute; 1986.
18. Lewis DO, Pincus JH, Lovely R, Spitzer E, Moy E. Biopsychosocial characteristics of matched samples of delinquents and nondelinquents. *J Am Acad Child Adolesc Psychiatry*. 1987;26:744-752.
19. Davis DL, Bean GJ, Schumacher JE, Stringer TL. Prevalence of emotional disorders in a juvenile justice institutional population. *Am J Forensic Psychol*. 1991;9:5-17.
20. Eppright TD, Kashani JH, Robison BD, Reid JC. Comorbidity of conduct disorder and personality disorders in an incarcerated juvenile population. *Am J Psychiatry*. 1993;150:1233-1236.
21. Steiner H, Garcia IG, Mathews Z. Posttraumatic stress disorder in incarcerated juvenile delinquents. *J Am Acad Child Adolesc Psychiatry*. 1997;36:357-365.
22. Duclos CW, Beals J, Novins DK, Martin C, Jewett CS, Manson SM. Prevalence

- of common psychiatric disorders among American Indian adolescent detainees. *J Am Acad Child Adolesc Psychiatry*. 1998;37:866-873.
23. Gray TA, Wish ED. *Substance Abuse Need for Treatment Among Arrestees (SANTA) in Maryland: Youth in the Juvenile Justice System*. College Park, Md: Center for Substance Abuse Research; 1998.
 24. Cauffman E, Feldman S, Waterman J, Steiner H. Posttraumatic stress disorder among female juvenile offenders. *J Am Acad Child Adolesc Psychiatry*. 1998;37:1209-1216.
 25. Atkins DL, Pumariega AJ, Rogers K, Montgomery L, Nybro C, Jeffers G, Sease F. Mental health and incarcerated youth, I: prevalence and nature of psychopathology. *J Child Fam Stud*. 1999;8:193-204.
 26. Pliszka SR, Sherman JO, Barrow MV, Irick S. Affective disorder in juvenile offenders: a preliminary study. *Am J Psychiatry*. 2000;157:130-132.
 27. Aarons GA, Brown SA, Hough RL, Garland AF, Wood PA. Prevalence of adolescent substance use disorders across five sectors of care. *J Am Acad Child Adolesc Psychiatry*. 2001;40:419-426.
 28. Garland AF, Hough RL, McCabe KM, Yeh M, Wood PA, Aarons GA. Prevalence of psychiatric disorders in youth across five sectors of care. *J Am Acad Child Adolesc Psychiatry*. 2001;40:409-418.
 29. Whitaker A, Johnson J, Shaffer D, Rapoport JL, Kalikow K, Walsh BT, Davies M, Braiman S, Dolinsky A. Uncommon troubles in young people: prevalence estimates of selected psychiatric disorders in a nonreferred adolescent population. *Arch Gen Psychiatry*. 1990;47:487-496.
 30. Christie KA, Burke JD, Regier DA, Rae DS, Boyd JH, Locke BZ. Epidemiologic evidence for early onset of mental disorders and higher risk of drug abuse in young adults. *Am J Psychiatry*. 1988;145:971-975.
 31. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Earlbaum Associates; 1988.
 32. Shaffer D, Fisher P, Dulcan MK, Davies M, Piacentini J, Schwab-Stone ME, Lahey BB, Bourdon K, Jensen PS, Bird HR, Canino G, Regier DA. The NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3): description, acceptability, prevalence rates, and performance in the MECA Study. *J Am Acad Child Adolesc Psychiatry*. 1996;35:865-877.
 33. Pastore AL, Maguire K. *Sourcebook of Criminal Justice Statistics—1999*. Washington, DC: US Dept of Justice; 2000.
 34. US Bureau of the Census. *The Hispanic Population*. Washington, DC: US Dept of Commerce; 2001.
 35. US Bureau of the Census. *Population by Race and Hispanic or Latino Origin for the United States: 1990 and 2000*. Washington, DC: US Dept of Commerce; 2001: table 1. Available at: <http://www.census.gov/population/www/cen2000/phc-t1.html>. Accessed June 6, 2002.
 36. Grisso T, Tomkins A, Casey P. Psychosocial concepts in juvenile law. *Law Hum Behav*. 1988;12:403-437.
 37. Illinois Criminal Justice Information Authority. *Trends and Issues 1997*. Chicago: Illinois Criminal Justice Information Authority; 1997.
 38. Federal Policy for the Protection of Human Subjects: Notices and Rules, part 2, Vol 56, No. 117 (18 June 1991), 56 *Federal Register* 28002-32.
 39. Shaffer D. Use of passive consent in child/adolescent mental health research—effect of letter from Dr Charles R. McCarthy, Director of the Office for Protection from Research Risks, NIH [editorial]. *Res Notes Child Adolesc Psychiatry*. Summer 1992;10.
 40. Fisher CB. Integrating science and ethics in research with high-risk children and youth. *Soc Res Child Dev*. 1993;7:1-27.
 41. Nolan K. Ethical issues: assent, consent, and behavioral research with adolescents. *Res Notes Child Psychiatry*. Summer 1992;7-10.
 42. Bravo M, Woodbury-Farina M, Canino GJ, Rubio-Stipec M. The Spanish translation and cultural adaptation of the Diagnostic Interview Schedule for Children (DISC) in Puerto Rico. *Cult Med Psychiatry*. 1993;17:329-344.
 43. Fisher PW, Shaffer D, Piacentini JC, Lapkin J, Kafantaris V, Leonard H, Herzog DB. Sensitivity of the Diagnostic Interview Schedule for Children, 2nd edition (DISC-2.1), for specific diagnoses of children and adolescents. *J Am Acad Child Adolesc Psychiatry*. 1993;32:666-673.
 44. Piacentini J, Shaffer D, Fisher PW, Schwab-Stone ME, Davies M, Gioia P. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), III: concurrent criterion validity. *J Am Acad Child Adolesc Psychiatry*. 1993;32:658-665.
 45. Schwab-Stone M, Fisher PW, Piacentini J, Shaffer D, Davies M, Briggs M. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), II: test-retest reliability. *J Am Acad Child Adolesc Psychiatry*. 1993;32:651-657.
 46. Shaffer D, Schwab-Stone ME, Fisher PW, Cohen P, Piacentini J, Davies M, Conners CK, Regier D. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), I: preparation, field testing, interrater reliability, and acceptability. *J Am Acad Child Adolesc Psychiatry*. 1993;32:643-650.
 47. Schwab-Stone ME, Shaffer D, Dulcan MK, Jensen PS, Fisher P, Bird HR, Goodman SH, Lahey BB, Lichtman JH, Canino G, Rubio-Stipec M, Rae DS. Criterion validity of the NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3). *J Am Acad Child Adolesc Psychiatry*. 1996;35:878-888.
 48. Thompson LL, Riggs PD, Mikulich SK, Crowley TJ. Contribution of ADHD symptoms to substance problems and delinquency in conduct-disordered adolescents. *J Abnorm Child Psychol*. 1996;24:325-347.
 49. Bird HR, Davies M, Fisher P, Narrow WE, Jensen PS, Hoven C, Cohen P, Dulcan MK. How specific is specific impairment? *J Am Acad Child Adolesc Psychiatry*. 2000;39:1182-1189.
 50. Roberts RE, Attkisson C, Rosenblatt A. Prevalence of psychopathology among children and adolescents. *Am J Psychiatry*. 1998;155:715-725.
 51. Costello EJ, Angold A, Burns BJ, Erkanli A, Stangle DK, Tweed DL. The Great Smoky Mountains Study of Youth: functional impairment and serious emotional disturbance. *Arch Gen Psychiatry*. 1996;53:1137-1143.
 52. Bird HR, Canino G, Rubio-Stipec M, Ribera J. Further measures of the psychometric properties of the Children's Global Assessment Scale. *Arch Gen Psychiatry*. 1987;44:821-824.
 53. Shaffer D, Gould MS, Brasic J, Ambrosini P, Fisher P, Bird HR, Aluwahlia S. A Children's Global Assessment Scale (CGAS). *Arch Gen Psychiatry*. 1983;40:1228-1231.
 54. Cochran WG. *Sampling Techniques*. New York, NY: John Wiley & Sons; 1997.
 55. Levy PS, Lemeshow S. *Sampling of Populations: Methods and Applications*. New York, NY: John Wiley & Sons; 1999.
 56. Costello EJ, Angold A, Burns BJ, Stangle DK, Tweed DL, Erkanli A, Worthman CM. The Great Smoky Mountains Study of Youth: goals, design, methods, and the prevalence of DSM-III-R disorders. *Arch Gen Psychiatry*. 1996;53:1129-1136.
 57. Simonoff E, Pickles A, Meyer JM, Silberg JL, Maes HH, Loeber R, Rutter M, Hewitt JK, Eaves LJ. The Virginia Twin Study of Adolescent Behavioral Development: influences of age, sex, and impairment on rates of disorder. *Arch Gen Psychiatry*. 1997;54:801-808.
 58. Turner RJ, Gil AG. Psychiatric and substance use disorders in South Florida: racial/ethnic and gender contrasts in a young adult cohort. *Arch Gen Psychiatry*. 2002;59:43-50.
 59. Kessler RC, Walters EE. Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Comorbidity Survey. *Depress Anxiety*. 1998;7:3-14.
 60. McGee R, Feehan M, Williams S, Anderson J. DSM-III disorders from age 11 to age 15 years. *J Am Acad Child Adolesc Psychiatry*. 1992;31:50-59.
 61. Garrison CZ, Waller JL, Cuffe SP, McKeown RE, Addy CL, Jackson KL. Incidence of major depressive disorder and dysthymia in young adolescents. *J Am Acad Child Adolesc Psychiatry*. 1997;36:458-465.
 62. Feitel B, Margetson N, Chamas J, Lipman C. Psychosocial background and behavioral and emotional disorders of homeless and runaway youth. *Hosp Comm Psychiatry*. 1992;43:155-159.
 63. Famularo R, Kinscherff R, Fenton T. Psychiatric diagnoses of maltreated children: preliminary findings. *J Am Acad Child Adolesc Psychiatry*. 1992;31:863-867.
 64. Lewis DO, Yeager CA, Cobham-Portorreal CS, Klein N, Showalter C, Anthony A. A follow-up of female delinquents: maternal contributions to the perpetuation of deviance. *J Am Acad Child Adolesc Psychiatry*. 1991;30:197-201.
 65. Teplin LA, Abram KM, McClelland GM. Prevalence of psychiatric disorders among incarcerated women, I: pretrial jail detainees. *Arch Gen Psychiatry*. 1996;53:505-512.
 66. Cohen P, Cohen J, Brook JS. An epidemiological study of disorders in late childhood and adolescence, II: persistence of disorders. *J Child Psychol Psychiatry*. 1993;34:869-877.
 67. Kandel DB, Johnson JG, Bird HR, Canino G, Goodman SH, Lahey BB, Regier DA, Schwab-Stone M. Psychiatric disorders associated with substance use among children and adolescents: findings from the Methods for the Epidemiology of Child and Adolescent Mental Disorders (MECA) Study. *J Abnorm Child Psychol*. 1997;25:121-132.
 68. Newman DL, Moffitt TE, Caspi A, Magdol L, Silva PA, Stanton WR. Psychiatric disorder in a birth cohort of young adults: prevalence, comorbidity, clinical significance, and new case incidence from ages 11 to 21. *J Consult Clin Psychol*. 1996;64:552-562.
 69. Angold A, Costello EJ. Depressive comorbidity in children and adolescents: empirical, theoretical, and methodological issues. *Am J Psychiatry*. 1993;150:1779-1791.
 70. Bukstein OG, Brent DA, Kaminer Y. Comorbidity of substance abuse and other psychiatric disorders in adolescents. *Am J Psychiatry*. 1989;146:1131-1141.
 71. Abram KM, Teplin LA. Co-occurring disorders among mentally ill jail detainees: implications for public policy. *Am Psychol*. 1991;46:1036-1045.
 72. Hesselbrock MN, Meyer RE, Keener JJ. Psychopathology in hospitalized alcoholics. *Arch Gen Psychiatry*. 1985;42:1050-1055.
 73. Caton CLM, Gralnick A, Bender S, Simon R. Young chronic patients and substance abuse. *Hosp Comm Psychiatry*. 1989;40:1037-1040.

74. Breakey WR, Fischer PJ, Kramer M, Nestadt G, Romanoski AJ, Ross A, Royal RM, Stine OC. Health and mental health problems of homeless men and women. *JAMA*. 1989;262:1352-1357.
75. Poe-Yamagata E, Butts JA. *Female Offenders in the Juvenile Justice System*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 1996.
76. Zoccolillo M. Co-occurrence of conduct disorder and its adult outcomes with depressive and anxiety disorders: a review. *J Am Acad Child Adolesc Psychiatry*. 1992;31:547-556.
77. Loeber R, Stouthamer-Loeber M. Development of juvenile aggression and violence. *Am J Psychol*. 1998;53:242-259.
78. Gender-Specific Programming for Girls Advisory Committee. *Guiding Principles for Promising Female Programming*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 1998. Available at: <http://www.ojjdp.ncjrs.org/pubs/principles/contents.html>. Accessed July 30, 2001.
79. Werner EE. High-risk children in young adulthood: a longitudinal study from birth to 32 years. *Am J Orthopsychiatry*. 1989;59:72-81.
80. National Research Council. *Losing Generations: Adolescents in High-Risk Settings*. Washington, DC: National Academy Press; 1993.
81. Lewis DO, Yeager CA, Lovely R, Stein A, Cobham-Portorreal CS. A clinical follow-up of delinquent males: ignored vulnerabilities, unmet needs, and the perpetuation of violence. *J Am Acad Child Adolesc Psychiatry*. 1994;33:518-528.
82. Leventhal T, Brooks-Gunn J. The neighborhoods they live in: the effects of neighborhood residence on child and adolescent outcomes. *Psychol Bull*. 2000;126:309-337.
83. Buckner JC, Bassuk EL. Mental disorders and service utilization among youths from homeless and low-income housed families. *J Am Acad Child Adolesc Psychiatry*. 1997;36:890-900.
84. Dembo R, Williams L, Schmeidler J. Gender differences in mental health service needs among youths entering a juvenile detention center. *J Prison Jail Health*. 1993;12:73-101.
85. Cocozza JJ. *Responding to the Mental Health Needs of Youth in the Juvenile Justice System*. Seattle, Wash: National Coalition for the Mentally Ill in the Criminal Justice System; 1992.
86. Lattimore PK, Linster RL, MacDonald JM. Risk of death among serious young offenders. *J Res Crime Delinq*. 1997;34:187-209.
87. Loeber R, DeLamatre M, Tita G, Cohen J, Stouthamer-Loeber M, Farrington DP. Gun injury and mortality: the delinquent backgrounds of juvenile victims. *Violence Vict*. 1999;14:339-352.
88. National Alliance for the Mentally Ill. *Families on the Brink: The Impact of Ignoring Children With Serious Mental Illness: Results of a National Survey of Parents and Other Caregivers*. Arlington, Va: National Alliance for the Mentally Ill; 1999.
89. Knitzer J. Children's mental health: changing paradigms and policies. In: Zigler EF, Kagan SL, Hall NW, eds. *Children, Families, and Government: Preparing for the Twenty-first Century*. New York, NY: Cambridge University Press; 1996:207-232.
90. Srebnik D, Cauce AM, Baydar N. Help-seeking pathways for children and adolescents. *J Emotional Behav Dis*. 1996;4:210-220.
91. Redding RE. Juvenile offenders in criminal court and adult prison: legal, psychological, and behavioral outcomes. *Juvenile Fam Court J*. Winter 1999;1-20.
92. US Department of Health and Human Services. *Mental Health: A Report of the Surgeon General*. Rockville, Md: US Dept of Health and Human Services; 1999.
93. Grisso T. What we know about youths' capacities as trial defendants. In: Grisso T, Schwartz RG, eds. *Youth on Trial: A Developmental Perspective on Juvenile Justice*. Chicago, Ill: University of Chicago Press; 2000:139-171.
94. Bishop DM. Juvenile offenders in the adult criminal justice system. *Crime Justice*. 2000;27:81-167.
95. Mears DP. Getting tough with juvenile offenders: explaining support for sanctioning youths as adults. *Criminal Justice Behav*. 2001;28:206-226.
96. Hamparian DM, Estep LK, Muntean SM, Priestino RR, Swisher RG, Wallace PL, White TL. *Major Issues in Juvenile Justice Information and Training*. Washington, DC: US Dept of Justice; 1982.
97. Heymann SJ, Earle A. The impact of welfare reform on parents' ability to care for their children's health. *Am J Public Health*. 1999;89:502-505.
98. Center for Mental Health Services. *Mental Health and Substance Abuse Services Under the State Children's Health Insurance Program*. Rockville, Md: US Dept of Health and Human Services, Substance Abuse and Mental Health Services Administration; 2000.
99. Burns BJ, Costello EJ, Erkanli A, Tweed DL, Farmer EMZ, Angold A. Insurance coverage and mental health service use by adolescents with serious emotional disturbance. *J Child Fam Studies*. 1997;6:89-111.
100. Children's Defense Fund. *Families Struggling to Make It in the Workforce: A Post Welfare Report*. Washington, DC: Children's Defense Fund; 2000. Available at: <http://www.childrensdefense.org/pdf/CMPReport.pdf>. Accessed June 5, 2002.
101. Sherman A, Amey C, Duffield B, Ebb N, Weinstein D. *Welfare to What: Early Findings on Family Hardship and Well-Being*. Washington, DC: Children's Defense Fund and National Coalition for the Homeless; 1998. Available at: <http://www.childrensdefense.org/pdf/wlfwhat.pdf>. Accessed June 5, 2002.
102. General Accounting Office. *Medicaid and SCHIP: Comparisons of Outreach, Enrollment Practices, and Benefits*. Washington, DC: General Accounting Office; 2000.
103. Zuckerman DM. The evolution of welfare reform: policy changes and current knowledge. *J Soc Issues*. 2000;56:811-820.
104. Porter K, Primus W. *Recent Changes in the Impact of the Safety Net on Child Poverty*. Washington, DC: Center on Budget and Policy Priorities; 1999. Available at: <http://www.cbpp.org/12-23-99wel-es.htm>. Accessed June 5, 2002.
105. Knitzer J, Yoshikawa H, Cauthen NK, Aber JL. Welfare reform, family support, and child development: perspectives from policy analysis and developmental psychopathology. *Dev Psychopathol*. 2000;12:619-632.
106. Iglehart JK. Managed care and mental health. *N Engl J Med*. 1996;334:131-135.
107. Frank RG. The creation of Medicare and Medicaid: the emergence of insurance and markets for mental health services. *Psychiatr Serv*. 2000;51:465-468.
108. Shirk S, Talmi A, Olds D. A developmental psychopathology perspective on child and adolescent treatment policy. *Dev Psychopathol*. 2000;12:835-855.
109. Mechanic D. Topics for our times: managed care and public health opportunities. *Am J Public Health*. 1998;88:874-875.
110. Kelleher KJ, McInerney TK, Gardner WP, Childs GE, Wasserman RC. Increasing identification of psychosocial problems: 1979-1996. *Pediatrics*. 2000;105:1313-1321.
111. Burns BJ. Mental health service use by adolescents in the 1970s and 1980s. *J Am Acad Child Adolesc Psychiatry*. 1991;30:144-150.
112. US Department of Health and Human Services. *Report of the Surgeon General's Conference on Children's Mental Health: A National Action Agenda*. Washington, DC: US Dept of Health and Human Services; 2000.
113. Thomas CR, Holzer CE III. National distribution of child and adolescent psychiatrists. *J Am Acad Child Adolesc Psychiatry*. 1999;38:9-16.
114. Coalition for Juvenile Justice. *Handle With Care: Serving the Mental Health Needs of Young Offenders*. Washington, DC: Coalition for Juvenile Justice; 2000.
115. Redding RE. Barriers to meeting the mental health needs of juvenile offenders. *Dev Mental Health Law*. 1999;19:1-23.
116. Faenza M, Siegfried C, Wood J. *Community Perspectives on the Mental Health and Substance Abuse Treatment Needs of Youth Involved in the Juvenile Justice System*. Alexandria, Va: National Mental Health Association and the Office of Juvenile Justice and Delinquency Prevention; 2000.
117. Grisso T. Juvenile offenders and mental illness. *Psychiatry Psychol Law*. 1999;6:143-151.
118. Ulzen TPM, Hamilton H. The nature and characteristics of psychiatric comorbidity in incarcerated adolescents. *Can J Psychiatry*. 1998;43:57-63.