

Bile Duct Injury During Laparoscopic Cholecystectomy

Results of an Italian National Survey on 56 591 Cholecystectomies

Gennaro Nuzzo, MD; Felice Giuliani, MD; Ivo Giovannini, MD; Francesco Ardito, MD; Fabrizio D'Acapito, MD; Maria Vellone, MD; Marino Murazio, MD; Giovanni Capelli, MD

Hypothesis: Bile duct injury (BDI) remains the most serious complication of cholecystectomy. With laparoscopic cholecystectomy (LC), the incidence has become more frequent. This study verifies the current incidence, mechanism, presentation, and treatment of BDI occurring during LC in general surgical practice.

Design: Anonymous retrospective multicenter survey.

Setting: Department of surgery at a university referral center, collecting data from general surgical units.

Patients: Data from 56 591 patients who underwent LC between January 1, 1998, and December 31, 2000, in 184 hospitals in Italy were analyzed.

Main Outcome Measures: Current incidence, mechanism, presentation, and treatment of BDI occurring during LC in general surgical practice.

Results: Two hundred thirty-five BDIs were reported, with an overall incidence of 0.42%. There were no risk factors in 80.0% of the patients. Poor identification of the

anatomical features of the hepatic pedicle was the most frequently reported cause (36.8%), and technical problems accounted for 27.0% of causes. The incidence of BDI was higher during cholecystitis ($P < .001$) and decreased with increasing number of LCs performed by the surgical teams ($P < .01$). There was no difference in incidence according to technique (French or US) or to routine or selective intraoperative cholangiography. One hundred eight BDIs (46.0%) were recognized intraoperatively and immediately repaired in 89.8% of patients. One hundred twenty-seven BDIs (54.0%) were diagnosed postoperatively, the dominant manifestation being biliary fistula (44.1%).

Conclusions: This study confirms a higher incidence of BDI during LC. It highlights the relevance of the number of previously performed LCs and of the correct surgical technique to avoid BDI. The need for correct procedures, adequate expertise of the repairing surgeon in BDI repairs, and a multidisciplinary approach in the management of BDI is emphasized.

Arch Surg. 2005;140:986-992

LAPAROSCOPIC CHOLECYSTECTOMY (LC), after rapidly substituting traditional cholecystectomy, represents the gold standard for surgical treatment of cholelithiasis. Despite the progress achieved, bile duct injuries (BDIs) still represent an important complication, and have become more frequent than in the past.¹⁻⁸ Since the introduction of LC, more than 10 years ago, the rate of BDI does not seem to be substantially changed, even if some researchers report a trend toward a decrease.⁹ Moreover, evaluation of the literature on treatment of BDI suggests indirectly that the actual rate of BDI may be higher than that commonly estimated.

We report the results of a recently performed multicenter national survey to provide an up-to-date assessment of the rate,

causes, clinical manifestations, and treatment of BDI in those who underwent LC in Italy between January 1, 1998, and December 31, 2000.

METHODS

The national survey was based on anonymous questionnaires sent to 316 heads of surgical units, all members of the Italian Society of Surgery.

CME course available at
www.archsurg.com

Questions were divided into 4 main sections. The first section included general information on the number of LCs performed in the 3-year period (1998-2000) in each unit, use of the French technique (ie, surgeon between the

Author Affiliations:
Department of Surgical Sciences, Hepato-Biliary Surgery Unit, Catholic University, School of Medicine, Rome (Drs Nuzzo, Giuliani, Giovannini, Ardito, D'Acapito, Vellone, and Murazio); and Department of Science and Society, Institute of Hygiene, University of Cassino, Cassino (Dr Capelli), Italy.

legs of the patient, umbilical telescope, operating port in left upper quadrant, liver retractor in epigastrium, and grasper in right upper quadrant) or the US technique (ie, surgeon to the left of the patient, umbilical telescope, operating port in epigastrium, and liver retractor and grasper in the right upper quadrant), use of intraoperative cholangiography and abdominal drainage, and number of BDIs that occurred during the 3-year period. The second section included preoperative data on BDI: indications for cholecystectomy, patient risk factors (obesity, previous abdominal surgery, and liver cirrhosis), and number of LCs performed by the surgical team (teams in Italy are usually composed of permanent members of hospital staff with prolonged collective shared experience). The third section included intraoperative data on BDI: technical complexity of LC, identification of the cause of injury, recognition of the injury (by the presence of bile in the operative field, at cholangiography, or by other modalities), immediate treatment, and immediate results. The fourth section included postoperative data on BDI: clinical presentation of the injury and decisions on treatment. The sources of data were in-depth medical record reviews, operative dictations, and detailed written records of the units. Statistical analysis was performed using Stata 7.0 software (Stata Corp, College Station, Tex), with data expressed as means, medians, and ranges and with the use of χ^2 and Fisher exact tests.

RESULTS

The survey collected 56 591 LCs from 184 Italian surgical units (response rate, 58.2%) between January 1, 1998, and December 31, 2000. The geographical distribution throughout the country was homogeneous.

INCIDENCE OF BDI

A total of 235 BDIs were reported (overall incidence, 0.42%). These were major injuries (involving the common bile duct [CBD], main biliary confluence, or main bile ducts) in 178 patients (75.7%) and minor injuries (involving the cystic duct or small peripheral or Luschka ducts) in 57 patients (24.3%).

The incidence of major injuries was 0.31%, ranging from 0% to 3.75% in individual units. At least 1 injury was reported by 68.5% (126/184) of the units.

CAUSES AND MECHANISMS OF BDI

These were identified in 163 patients (69.4% of cases). The most frequently reported cause was poor identification of the anatomical features of the hepatic pedicle (36.8% of cases), followed by inflammatory changes in the gallbladder (23.3%), anatomical anomalies (12.9%), improper use of monopolar coagulation (12.3%), an unspecified technical mistake (9.8%), and a problem during the control of intraoperative hemorrhage (4.9%). Thus, technical mistakes, reflected by the latter 3 factors, accounted cumulatively for 27.0% of the recognized causes of injury.

With regard to the technical complexity of the operation during which the injury had occurred, the procedure was described as easy in 46.8% and difficult in 53.2% of cases, a rate that did not change significantly with the number of LCs performed.




Figure 1. Mean rate of major bile duct injuries (BDIs) for surgical teams with different volumes of laparoscopic cholecystectomy (LC) in the 3-year period. The rate significantly decreased ($P=.02$) with the increasing number of LCs. The number of teams in each group was similar.




Figure 2. Mean rate of cholecystitis in patients with bile duct injury reported by surgical teams with different volumes of laparoscopic cholecystectomy (LC) in the 3-year period. The change in rate was significant ($P=.04$).

RISK FACTORS FOR BDI

In 188 cases (80.0%), no risk factors related to the patient were reported. These were reported in only 47 cases (20.0%): obesity in 33, previous abdominal surgery in 9, and cirrhosis in 5. In 33 of these 47 cases, cholecystitis was an associated risk factor.

In 112 (47.7%) of the 235 BDIs, LC was performed for simple cholelithiasis; and in 123 (52.3%), for cholecystitis.

Laparoscopic cholecystectomies were performed for simple cholelithiasis in 61.1% of patients, and for cholecystitis in 38.9% of patients; the incidence of injuries was 0.32% in the former and 0.56% in the latter group of patients ($P<.001$).

The incidence of BDI significantly decreased with increasing number of LCs performed by the teams, ranging from 0.9% for teams who had performed fewer than 150 LCs in the 3-year period to 0.3% for those who had performed more than 450 LCs ($P<.01$). A similar result was observed for major injuries (**Figure 1**).

Interestingly, BDIs reported by teams performing more LCs were more frequently associated with cholecystitis (**Figure 2**), while those reported by teams performing

Table 1. Treatment in 100 Patients With Intraoperatively Recognized Major BDI

Treatment	No. (%) of Patients*
Laparotomy	91 (91.0)
Repair over the T tube	37 (40.7)
Roux-en-Y hepaticojejunostomy	27 (29.7)
End-to-end reconstruction over the T tube	15 (16.5)
Bridging of biliary stumps with the T tube	11 (12.1)
Drainage of the abdomen	1 (1.1)
Laparoscopy	9 (9.0)
Drainage of the abdomen	6 (66.7)
Repair over the T tube	2 (22.2)
Suture of the CBD	1 (11.1)

Abbreviations: BDI, bile duct injury; CBD, common bile duct.
*Percentages within each section may not total 100 because of rounding.

fewer LCs were more frequently associated with simple cholelithiasis.

SURGICAL TECHNIQUE

The French technique was used in 123 units (66.8%) and the US technique in 58 units (31.5%). Three units (1.6%) did not answer this question. (Percentages do not total 100 because of rounding.)

One hundred sixty-six BDIs occurred in 85 units using the French technique (85/123=69.1%). For a total of 41 256 LCs, the incidence of BDI was 0.40% (and 0.29% of major injuries).

Sixty-eight BDIs occurred in 40 units using the US technique (40/58=69.0%). For a total of 15 239 LCs, the incidence of BDI was 0.45% (and 0.36% of major injuries).

The incidence of injuries was, thus, similar for units using different techniques.

INTRAOPERATIVE CHOLANGIOGRAPHY

Routine cholangiography was performed in 10.3% of the units, and a total of 25 BDIs occurred. Fourteen injuries (56.0%) were recognized during LC: 7 before performing cholangiography, by the presence of bile in the operative field, and 7 during cholangiography (in one case, the injury was caused by cannulation of the cystic duct). The incidence of BDI in these units was 0.32%.

Selective cholangiography was used in 89.7% of the units, and a total of 210 BDIs occurred. Ninety-four injuries (44.8%) were diagnosed during LC: 72 by the presence of bile, 14 at cholangiography, and 8 by the presence of a double biliary stump. The incidence of BDI in these units was 0.43%. Of the 94 patients who had a BDI diagnosed intraoperatively, only 43 (45.7%) underwent cholangiography after diagnosis.

The overall incidences of BDI (0.32% vs 0.43%) and of intraoperative diagnosis of BDI (56.0% vs 44.8%) were not significantly different in units using routine vs selective cholangiography ($P=.25$ and $.38$, respectively).

DIAGNOSIS OF BDI AND TREATMENT

A total of 108 BDIs (46.0%) were diagnosed during LC; these were 100 major and 8 minor injuries. The remain-

ing 127 (54.0%) were recognized postoperatively; these were 78 major and 49 minor injuries.

Intraoperative Diagnosis of BDI

Intraoperative diagnosis occurred in 79 cases (73.1%) by the presence of bile in the operative field, in 21 (19.4%) by cholangiography, and in 8 (7.4%) by the presence of a double biliary stump. (Percentages do not total 100 because of rounding.)

Major BDI occurred in 100 patients. Immediate repair of BDI was performed in 93 patients (93.0%), in 90 after laparotomy and in 3 during laparoscopy. The most frequent repair was suture or reconstruction of the CBD with positioning of a T tube (65.0%) (**Table 1**).

Immediate failure of repair was reported in 15 patients (15/93=16.1%): 7 required a subsequent operation within a week, 5 were treated endoscopically, 1 was treated percutaneously, and 2 were treated with a combination endoscopic/percutaneous method.

In 7 patients (7.0%), simple abdominal drainage was performed, in 6 during laparoscopy and in 1 after laparotomy (Table 1). Three of these patients needed a subsequent operation early, and 4 underwent endoscopic or percutaneous treatment.

Minor BDI occurred in 8 patients. Immediate repair of BDI was performed in 3 patients during laparoscopy and in 1 after laparotomy (followed in this patient by endoscopic treatment). In the other 4 patients, simple abdominal drainage was performed, in 3 during laparotomy and in 1 during laparoscopy.

Postoperative Diagnosis of BDI

The clinical manifestation was biliary fistula (bilious drainage from an operatively placed drain) in 44.1% of cases, bile peritonitis in 37.8%, and jaundice in 18.1%. When 2 or more manifestations coexisted, the dominant one determining management priorities was considered.

Biliary fistula tended to be more frequently associated with a minor than a major injury (58.9% vs 41.1%; $P=.30$). Bile peritonitis was associated more frequently with a major injury (66.7% vs 33.3%; $P<.005$), and jaundice was always associated with a major injury (100.0% vs 0%; $P<.005$).

The clinical manifestation of a BDI was only partly conditioned by the presence of an abdominal drainage. Abdominal drainage was used routinely in 45.7% (84/184) of units. In these units, 55 BDIs were diagnosed postoperatively, and the most frequent clinical presentation (60.0% of cases) was an external biliary fistula; this was associated with bile peritonitis in an additional 27.3% of cases. In the 54.3% (100/184) of units in which drainage was not used routinely, 72 BDIs were diagnosed postoperatively, and the most frequent clinical presentation was bile peritonitis (45.8% of cases). The manifestation through a biliary fistula was more frequent in units using routine abdominal drainage ($P=.002$), whereas bile peritonitis was more frequent in units not using routine drainage ($P=.03$).

Table 2. Treatment in 127 Patients With Postoperatively Recognized BDI

Treatment	No. (%) of Patients	No. of Major Injuries	No. of Minor Injuries
Surgery	77 (60.6)	59	18
Endoscopy/radiology	25 (19.7)	5	20
Spontaneous healing	11 (8.7)	0	11
Referral to another center	8 (6.3)	8	0
Not evaluable	6 (4.7)	6	0
Total	127 (100.0)	78	49

Abbreviation: See Table 1.

Table 3. Treatment in 23 Patients With a Biliary Fistula From a Major BDI

Treatment	No. (%) of Patients	Those Who Received Treatment Within 15 d	Those Who Received Treatment Between 16 and 45 d
Surgery	18 (78.3)	15	3
Roux-en-Y hepaticojejunostomy	8	6	2
Repair over the T tube	7	6	1
End-to-end reconstruction over the T tube	2	2	0
Suture of the CBD	1	1	0
Endoscopy	3 (13.0)	3	0
Referral to another center	2 (8.7)	NA	NA
Total*	23 (100.0)	18	3

Abbreviations: BDI, bile duct injury; CBD, common bile duct; NA, data not applicable.

*The total in column 2 differs from the sum of columns 3 and 4 because the patients referred to another center did not have data for receipt of treatment.

Treatment of Postoperatively Diagnosed BDI

Surgical treatment was performed in 60.6% of patients, and endoscopic or radiologic treatment in 19.7% of patients (**Table 2**).

There were 56 patients with a biliary fistula: 23 (41.1%) had a major injury and 33 (58.9%) had a minor injury.

For the 23 major injuries, the most common treatment was surgical repair (**Table 3**). For the 33 minor injuries, the most common treatment was endoscopy (13 patients, or 39.4% of cases). In 11 patients (33.3%), the fistula closed spontaneously. Of the remaining 9 patients (27.3%), 7 underwent a new laparoscopy with closure of the cystic stump or of a minor bile duct and 2 underwent laparotomy (in one case, combined with endoscopy).

There were 48 patients with bile peritonitis: 32 (66.7%) had a major injury and 16 (33.3%) had a minor injury.

For the 32 major injuries, the most common decision was to reoperate immediately on the patient to drain the bile peritonitis and at the same time repair the injury (**Table 4**). For the 16 minor injuries, the most frequent treatment was also surgical drainage and simultaneous repair of the injury (9 patients [56.2%]). In the other 7 patients (43.8%), treatment was by endoscopy, in 5 after surgical or percutaneous drainage of the abdomen.

Table 4. Treatment in 32 Patients With Bile Peritonitis From a Major BDI

Treatment	No. (%) of Patients
Drainage of bile peritonitis with simultaneous surgical repair	24 (75.0)
Repair over the T tube	10
Roux-en-Y hepaticojejunostomy	6
Suture of the CBD	4
Suture of the hepatic duct	
Right	1
Left	1
Removal of the surgical clip from the CBD	1
Drainage of microscopic injury of the CBD	1
Drainage of bile peritonitis with postponement of surgical repair	6 (18.8)
Surgical	3
Percutaneous	3
Referral to another center	2 (6.2)
Total	32 (100.0)

Abbreviations: See Table 1.

Table 5. Treatment in 23 Patients With Jaundice From a Major BDI

Treatment	No. (%) of Patients	Those Who Received Treatment Within 15 d	Those Who Received Treatment Between 16 d and 6 mo
Surgery	17 (73.9)	12	5
Roux-en-Y hepaticojejunostomy	12	7	5
Repair over the T tube	3*	3*	0
End-to-end reconstruction over the T tube	1	1	0
Removal of the clip from the main bile duct	1	1	0
Endoscopy	3 (8.7)*†	1*	2
Referral to another center	4 (17.4)	NA	NA
Total‡	23 (100.0)*	12	7

Abbreviations: See Table 3.

*In 1 patient, there were 2 treatments, first surgery and then endoscopy.

†The numerator used was 2 (vs 3) because this was the second treatment for one patient.

‡The total in column 2 differs from the sum of columns 3 and 4 because the patients referred to another center did not have data for receipt of treatment.

There were 23 patients with jaundice, all with a major injury. Surgical treatment was most common, and in 12 patients (52.2%), was performed within 15 days of LC (**Table 5**).

COMMENT

These data provide the most recent and comprehensive information on BDI associated with LC in Italy.

The relevance of this survey is supported by the high rate of response to the questionnaire (58.2%), higher than the mean rate for similar national surveys pub-

Table 6. Rate of Reply to National Surveys on BDI During LC

Source	Country	No. of Mailed Questionnaires	No. of Answered Questionnaires	Rate of Reply, %*
Deziel et al, ¹⁰ 1993	United States	4292	1750	40.8
Gouma and Go, ¹¹ 1994	The Netherlands	138	122	88.4
Gigot et al, ³ 1997	Belgium	98	75	76.5
Windsor and Pong, ¹² 1998	New Zealand	184	111	60.3
Torkington et al, ¹³ 1998	United Kingdom-Ireland	1100	362	32.9
Regöly-Mérei et al, ¹⁴ 1998	Hungary	119	105	88.2
Archer et al, ¹⁵ 2001	United States	3657	1661	45.4
Present study, 2002	Italy	316	184	58.2

Abbreviations: BDI, bile duct injury; LC, laparoscopic cholecystectomy.

*The average rate of reply for all evaluable surveys, excluding the last one (the Italian survey), was 43.7% (calculated as $100 \times [\text{total number of replies}/\text{total number of questionnaires}]$).

lished during the past 10 years (43.7%) (**Table 6**). Furthermore, the geographical distribution of participating surgical units throughout the country was homogeneous, as was the distribution of surgical teams with different expertise.

The survey was limited to a brief period (3 years) to facilitate verification of data, and was performed long enough (10 years) after the introduction of LC in Italy to grant stabilization of the learning curve in the interviewed units.

The reported incidence of major BDI during LC ranges from 0.25% to 0.74%, and that of minor BDI from 0.28% to 1.7%.²⁻⁸ In our survey, the overall incidences of injuries and major injuries were 0.42% and 0.31%, respectively, and were, thus, similar to those reported in other Western national surveys.²⁻⁸ Contrary to the results of another recent report,⁹ this confirms that the incidence of BDI during LC remains higher than during traditional cholecystectomy.¹⁻⁸

Since the early days of LC, a higher incidence of injuries has been related to the learning curve of the surgeon³; also in our survey, the incidence has increased significantly with decreasing volume of LCs performed. However, this explanation alone is insufficient, and analysis of BDI occurring after stabilization of the learning curve has shown that one third of injuries may still be related to technical mistakes. Therefore, a critical component is also the correct operative technique.^{12,15,16}

Misidentification of the CBD for the cystic duct (the classic injury)³ is reported most commonly as the cause of BDI. The cause may also remain unrecognized, and this occurred in 30.6% of cases in our survey. In about a quarter of the patients in whom the cause was recognized, this was related to the surgical technique, and misidentification of the CBD for the cystic duct accounted for most injuries (36.8%).

Our results confirm that no cholecystectomy can be considered as a simple routine procedure, immune to the risk of BDI.¹⁷ Indeed, about half of BDIs occurred during LCs described as "technically easy"; furthermore, in 80.0% of injuries, no risk factors related to the patient were present.

Among risk factors, the survey confirms the importance of cholecystitis^{18,19} by showing a higher incidence of BDI in the presence of cholecystitis compared with simple cholelithiasis (0.56% vs 0.32%; $P < .001$). Interestingly, BDIs were more often associated with cholecystitis in teams per-

forming more LCs, and with simple cholelithiasis in teams performing fewer LCs. This probably reflects a more cautious attitude of the latter in the presence of cholecystitis, with less frequent use of laparoscopy or more frequent conversion. This may highlight the importance of conversion to laparotomy in difficult cases.

With regard to the choice of the French or US technique, neither one, if correctly used, seems to be associated with an increased risk of injury,¹² and indeed, the risk was not significantly different in our survey. With either technique, the critical points are good exposure of the cystic duct and hepatic pedicle and adhesion to the same principles that apply to laparotomic cholecystectomy. This was emphasized recently by Strasberg et al,²⁰ who reported that in more than 80% of cases of a series of BDIs, the cystic duct had been interrupted prematurely before completing identification and preparation of anatomical structures.

Opinions on the advantages of routine vs selective use of intraoperative cholangiography are still diverging. Routine cholangiography should reduce the risk of BDI by providing a clear map of the biliary tree and of its variants and a more prompt diagnosis of unsuspected injuries, thus avoiding an increase in their complexity.²¹⁻²⁴ However, this has not yet been proved.^{16,21,25-28} Furthermore, in some instances, cholangiography may not prevent the injury or may carry its own risk of causing an injury. In our survey, 10.3% of units were using routine cholangiography, and these did not report a significantly different incidence of BDI or of intraoperatively recognized injuries. Whatever the explanation of these results (incorrect execution or interpretation of cholangiography or occurrence of the injury after cholangiography), routine cholangiography does not seem to entirely protect from BDI.

In units not using routine cholangiography, it was surprising that 54.3% of the patients with intraoperatively recognized BDIs (presence of bile in the operative field) did not undergo subsequent intraoperative cholangiography. Indeed, independently of the preference for routine or selective cholangiography, the cholangiographic assessment of an already recognized lesion is essential to the choice of appropriate treatment. This is also a reason for including intraoperative cholangiography in the training for LC.

The issue is also relevant for the appropriate timing of BDI repair, because a factor increasing the chances of a successful immediate repair is adequate intraoperative

cholangiographic assessment of the injury.²⁴ Another major component is the adequate expertise of the surgeon in BDI repair. These are important points because failure of immediate repair worsens the injury, increases the complexity of subsequent repairs, and impairs short- and long-term results.^{1,17,24,29,30} Also, the decision to convert to laparotomy depends on the balance between expertise of the surgeon and complexity of the injury. If the conditions for an optimal treatment are not available, even after laparotomy, an adequate choice is simple drainage of the abdomen and referral to a tertiary care center.

In our survey, immediate recognition of the injury took place in 46.0% of cases, which falls within the range (27%-73%) observed in previous multicenter surveys.^{2-8,15,31} There was a high rate of conversions (91.0%) and of immediate repair of major BDIs (93.0%), mostly with T-tube reconstructions, which are known to have the risks of dehiscence and long-term stricture.¹⁷ Although treatment of late strictures may be performed successfully by endoscopic stenting,³² repairs over a T tube should be reserved only for small side wall injuries, while major BDIs and total transections should be treated with hepaticojejunostomy.

With regard to postoperatively recognized BDI, after the introduction of LC, rates of occurrence of biliary fistula and bile peritonitis have increased, although biliary fistula is caused more frequently by a minor BDI.^{10,33} This was also the case in our survey, in which biliary fistula accounted for 44.1% of postoperative manifestations. There was a significantly more frequent association of bile peritonitis with major BDI, and jaundice was always due to a major BDI. A corollary of these findings is also the tendency of major injuries associated with LC to be more complex than those that were previously associated with open cholecystectomy.

The fact that 27.3% of patients undergoing drainage developed bile peritonitis despite the drainage may emphasize the importance of the correct positioning of drainages of adequate size (small sizes are usually used in LC) following the decision to drain the abdomen. However, the presence of drainage may not totally protect from the occurrence of bile peritonitis.

The treatment of postoperatively recognized injuries was surgical in 60.6% of patients, with a definite tendency to subsequently operate early on those with biliary fistula or jaundice. In patients with bile peritonitis, the most common decision was to reoperate quickly, with simultaneous repair of the lesion, and only 6.3% of patients were referred to a tertiary care center. Although our survey did not address long-term success (and results are, thus, unknown), these high percentages of early repair, and those of immediate repair in intraoperatively recognized BDI, do not correspond with the diffusion of expertise in complex hepatobiliary surgery and repairs, and are partly in contrast with the indications reported in the literature. For instance, the indication for simultaneous repair of a major injury during surgery for drainage of bile peritonitis is not largely accepted.³⁴⁻³⁷ Furthermore, in the case of a major injury with a biliary fistula, which is well drained and not associated with other complications (in particular with sepsis), urgent repair is not needed. Also, an accurate cholangiographic assessment may show that surgical treatment is not needed or,

conversely, that it requires a more expert surgeon. A recent large survey³⁸ has shown that outcome in patients with BDI after cholecystectomy is worse for those repaired by the same surgeon who caused the injury, and improves with the experience of the repairing surgeon.

The choice of type and timing of repair of a postoperatively recognized injury should take into account 2 important aspects. First, the recent advances in endoscopy and interventional radiology should be considered, which may quite often permit an appropriate nonsurgical treatment, especially in the case of minor BDI or partial injuries of the CBD.^{20,32,39,40} In our survey, this occurred in 19.7% of postoperatively recognized BDIs, a rate that probably underestimates the actual impact of nonsurgical treatment of BDIs. This greatly emphasizes the need for multidisciplinary care of these patients. Second, experience with surgical repairs has shown that the best results are generally obtained by repairs on a dilated biliary tree, performed by a surgeon expert in biliary repair, at a distance from cholecystectomy.^{1,17,30,37} This may be about 2 weeks after cholecystectomy in the case of jaundice without secondary complications or about 2 months after cholecystectomy in the case of bile peritonitis or biliary fistula, after complete control of sepsis and other secondary complications and total recovery of good health and nutritional state.

In conclusion, BDI during LC represents a risk for every general surgeon. This may be a severe complication, and, quite often, a young patient is involved. The frequency of BDI remains higher than that reported for open cholecystectomy. Our survey provides an overview of risk factors, mechanisms, type, severity, and patterns of detection and repair of BDI. It highlights the importance of surgical experience and correct technique to avoid BDI. In turn, the optimal management of BDI requires the use of correct procedures with regard to choice, timing of repair, or referral to a tertiary care center. A multidisciplinary approach in dedicated centers may also be needed, as more patients benefit from nonsurgical treatments.

Accepted for Publication: December 15, 2004.

Correspondence: Gennaro Nuzzo, MD, Department of Surgical Sciences, Hepato-Biliary Surgery Unit, Catholic University, School of Medicine, Largo Agostino Gemelli, 8-00168 Rome, Italy (chirucsc@rm.unicatt.it).

Funding/Support: This study was supported by Catholic University; and by 2003 D.1 funds from the Italian Ministry of University and Scientific Research, Rome.

Acknowledgment: We thank Helen Raiswell, Catholic University, for her assistance. We also thank all the surgeons who contributed to the study by returning the questionnaire, including 26 who returned it anonymously. These surgeons include the following: C. Allegri (Roma); E. Ancona (Padova); L. Angelini (Roma); L. Angelini (Rieti); A. Arullani (Roma); G. Azzena (Ferrara); E. Banchini (Piacenza); M. Basile (Messina); G. Bassi (Napoli); N. Bassi (Treviso); N. Basso (Roma); G. Belli (Napoli); S. Berardi (Isernia); G. Biliotti (Firenze); F. Borghi (Cuneo); D. Borrelli (Firenze); V. Bosco (Siracusa); U. Braccioni (Ancona); L. Capussotti (Torino); F. Caracciolo (Viterbo); A. Caracino (Pescara); M. Carboni (Roma); G. Cardi (Formia); A. Carli (Siena); P. Cavaliere (Savona); A. Cavallari (Bologna); A.

Cavallaro (Roma); E. Cavina (Pisa); A. Cennamo (Napoli); G. Centonze (Brindisi); G. Cerone (Tagliacozzo); G. Chiara (Pordenone); O. Ciccarelli (Popoli); E. Cirino (Catania); G. Citone (L'Aquila); L. Civello (Roma); L. Cogolo (Genova); G. Colecchia (Pescara); G. Collini (Udine); C. Cordiano (Verona); A. Corsini (Padova); C. Cortesini (Firenze); G. Costamagna (Roma); E. Croce (Milano); G. Cucchiara (Roma); D. D'Amico (Padova); A. D'Angelo (Vallo della Lucania); G. M. Daniele (Cagliari); D. De Anna (Udine); G. De Bernardinis (Modena); U. De Petris (Roma); C. De Werra (Napoli); G. Dettori (Sassari); V. Di Carlo (Milano); R. Dionigi (Varese); G. Doglietto (Roma); I. Donini (Ferrara); C. Eccher (Trento); E. Fedeli (Roma); F. Feliciotti (Ancona); A. Filippini (Roma); P. Forestieri (Napoli); E. Forni (Pavia); D. Forti (Milano); G. R. Fronda (Torino); M. Garavoglia (Novara); D. Garcea (Forlì); A. Garofalo (Roma); M. Gazzaniga (Genova); G. Giovaninetti (Verbania); P. C. Giulianotti (Grosseto); S. M. Giuliani (Brescia); G. B. Grassi (Roma); M. Guarasci (Corigliano Calabro); C. Huscher (Roma); R. Iovino (Napoli); G. Jacono (Napoli); E. Landi (Ancona); A. Liboni (Ferrara); G. Liguori (Trieste); M. Lise (Padova); A. Maffei Faccioli (Padova); P. Magistrelli (Roma); S. Malan (Salerno); C. Manca (Lecce); G. Mantovani (Milano); F. Maratia (Melfi); L. Mariani (Arezzo); D. Marrano (Bologna); F. Martin (Bolzano); P. Melita (Messina); G. Melotti (Modena); V. Memeo (Bari); G. Mensa (Asti); U. Mercati (Perugia); D. Migliori (Castrovillari); P. Mioli (Torino); D. Molino (Napoli); C. Montesani (Roma); A. Monti (Monferrato); A. Montori (Roma); M. Montorsi (Milano); A. Moraldi (Roma); R. Moretti (Firenze); M. Morino (Torino); M. Mulieri (Roma); G. Nanni (Vercelli); C. Natale (Foggia); E. Opocher (Bergamo); N. Palasciano (Matera); S. Panella (Napoli); G. Pappalardo (Roma); U. Parini (Aosta); S. Pedrazzoli (Padova); R. Pellicci (Genova); A. Peracchia (Milano); O. Petrillo (Vallo della Lucania); A. Picciocchi (Roma); M. Pisani (Piacenza); R. Porzio (Roma); F. Prete (Bari); A. Principe (Bologna); K. Reiner (Merano); F. Rendano (Caserta); E. Restini (Bari); A. Robecchi (Torino); E. Rocca (Catanzaro); L. Roncoroni (Parma); V. Saba (Ancona); B. Salerni (Brescia); M. Saviano (Modena); R. Scorza (Milano); F. Scutari (Roma); G. Serio (Verona); M. Sianesi (Parma); M. Simi (Roma); M. Spangaro (Bologna); G. Spinoglio (Alessandria); V. Stancanelli (Ravenna); G. Sturniolo (Messina); G. Tanzini (Siena); B. Tardio (San Giovanni Rotondo); A. M. Taschieri (Milano); P. Tenchini (Milano); O. Terranova (Padova); R. Tersigni (Roma); B. Thoman (Gorizia); F. Tonetti (Verbania); G. Torre (Genova); G. Tosolini (Pordenone); N. Tricarico (Napoli); G. Uggeri (Massa Carrara); T. Verrienti (Tricase); R. Vincenti (Benevento); P. Volterrani (Torino); A. Wiel Marin (Roma); M. Zanghì (Catania); V. Ziparo (Roma); and E. Zotti (Padova).

REFERENCES

- Bismuth H, Lazorthes F. *Les Traumatismes Opératoires de la Voie Biliaire Principale: Monographies de l'Association Française de Chirurgie: Rapport Présenté au 83^e Congrès Français de Chirurgie*. Paris, France: Masson; 1981.
- Russell JC, Walsh SJ, Mattie AS, et al. Bile duct injuries, 1989-1993: a statewide experience. *Arch Surg*. 1996;131:382-388.
- Gigot J, Etienne J, Aerts R, et al. The dramatic reality of biliary tract injury during laparoscopic cholecystectomy: an anonymous multicenter Belgian survey of 65 patients. *Surg Endosc*. 1997;11:1171-1178.
- Wherry DC, Marohn MR, Malanoski MP, et al. An external audit of laparoscopic cholecystectomy in the steady state performed in medical treatment facilities of the Department of Defense. *Ann Surg*. 1996;224:145-154.
- Richardson MC, Bell G, Fullarton GM. Incidence and nature of bile duct injuries following laparoscopic cholecystectomy: an audit of 5913 cases. *Br J Surg*. 1996;83:1356-1360.
- MacFadyen BV, Vecchio R, Ricardo AE, et al. Bile duct injury after laparoscopic cholecystectomy. *Surg Endosc*. 1998;12:315-321.
- Zgraggen K, Wehrli H, Metzger A, Buehler M, Frei E, Klaiber C; Swiss Association of Laparoscopic and Thoracoscopic Surgery. Complications of laparoscopic cholecystectomy in Switzerland: a prospective 3-year study of 10,174 patients. *Surg Endosc*. 1998;12:1303-1310.
- Adamsen S, Hart Hansen O, Funch-Jensen P, et al. Bile duct injury during laparoscopic cholecystectomy: a prospective nationwide series. *J Am Coll Surg*. 1997;184:571-578.
- Buanes T, Mjaland O, Waage A, Langeggen H, Holmboe J. A population-based survey of biliary surgery in Norway. *Surg Endosc*. 1998;12:852-855.
- Deziel DJ, Millikan KW, Economou SG, et al. Complications of laparoscopic cholecystectomy: a national survey of 4,292 hospitals and an analysis of 77,604 cases. *Am J Surg*. 1993;165:9-14.
- Gouma DJ, Go PM. Bile duct injury during laparoscopic and conventional cholecystectomy. *J Am Coll Surg*. 1994;178:229-233.
- Windsor JA, Pong J. Laparoscopic biliary injury: more than a learning curve problem. *Aust N Z J Surg*. 1998;68:186-189.
- Torkington J, Pereira J, Chalmers RT, et al. Laparoscopic cholecystectomy, bile duct injury and the British and Irish surgeon. *Ann R Coll Surg Engl*. 1998;80:119-121.
- Reguly-Mérei J, Ihasz M, Szeberin Z, et al. Biliary tract complications in laparoscopic cholecystectomy: a multicenter study of 148 biliary tract injuries in 26,440 operations. *Surg Endosc*. 1998;12:294-300.
- Archer SB, Brown DW, Smith D, et al. Bile duct injury during laparoscopic cholecystectomy. *Ann Surg*. 2001;234:549-559.
- Strasberg SM, Hertl M, Soper JN. An analysis of the problem of biliary injury during laparoscopic cholecystectomy. *J Am Coll Surg*. 1995;180:101-125.
- Jarnagin WR, Blumgart LH. Benign biliary strictures. In: Blumgart LH, Fong Y, eds. *Surgery of the Liver and Biliary Tract*. Edinburgh, Scotland: Harcourt-Saunders; 2000:895-933.
- Navez B, Mutter D, Ruzier Y, et al. Safety of laparoscopic approach for acute cholecystitis: retrospective study of 609 cases. *World J Surg*. 2001;25:1352-1356.
- Habib FA, Kolachalam RB, Khilanani R. Role of laparoscopic cholecystectomy in the management of gangrenous cholecystitis. *Am J Surg*. 2001;181:71-75.
- Strasberg SM, Eagon CJ, Drebin JA. The "hidden cystic duct" syndrome and the infundibular technique of laparoscopic cholecystectomy: the danger of the false infundibulum. *J Am Coll Surg*. 2000;191:661-667.
- Woods MS, Traverso LW, Kozarek RA, et al. Characteristics of biliary tract complications during laparoscopic cholecystectomy: a multi-institutional study. *Am J Surg*. 1994;167:27-33.
- Lezoche E, Paganini A, Carle F. Routine versus selective intra-operative cholangiography during laparoscopic cholecystectomy. *World J Surg*. 1993;17:686-687.
- Fletcher DR. Biliary injury at laparoscopic cholecystectomy: recognition and prevention. *Aust N Z J Surg*. 1993;63:673-677.
- Stewart L, Way LW. Bile duct injuries during laparoscopic cholecystectomy: factors that influence the results of treatment. *Arch Surg*. 1995;130:1123-1129.
- Phillips EH. Routine versus selective intraoperative cholangiography. *Am J Surg*. 1993;165:505-507.
- Branum G, Schmitt C, Baillie J, et al. Management of major biliary complications after laparoscopic cholecystectomy. *Ann Surg*. 1993;217:532-541.
- Soper NJ, Dunnegan DL. Routine versus selective intra-operative cholangiography during laparoscopic cholecystectomy. *World J Surg*. 1992;16:1133-1140.
- Lorimer JW, Fairfull-Smith RJ. Intraoperative cholangiography is not essential to avoid duct injuries during laparoscopic cholecystectomy. *Am J Surg*. 1995;169:344-347.
- Jarnagin WR, Blumgart LH. Operative repair of bile duct injuries involving the hepatic duct confluence. *Arch Surg*. 1999;134:769-775.
- Lillemoe KD, Pitt HA, Cameron JL. Current management of benign bile duct strictures. *Adv Surg*. 1992;25:119-174.
- Olsen D. Bile duct injuries during laparoscopic cholecystectomy. *Surg Endosc*. 1997;11:133-138.
- Costamagna G, Pandolfi M, Mutignani M, et al. Long-term results of endoscopic management of postoperative biliary strictures with increasing number of stents. *Gastrointest Endosc*. 2001;54:162-168.
- Belghiti J. Résultats de la cholécystectomie par coelioscopie. *Gastroenterol Clin Biol*. 1994;18:1000-1004.
- Lillemoe KD, Martin SA, Cameron JL, et al. Major bile duct injuries during laparoscopic cholecystectomy: follow-up after combined surgical and radiologic management. *Ann Surg*. 1997;225:459-468.
- Lillemoe KD, Melton GB, Cameron JL, et al. Postoperative bile duct strictures: management and outcome in the 1990s. *Ann Surg*. 2000;232:430-441.
- Murr MM, Gigot JF, Nagorney DM, et al. Long-term results of biliary reconstruction after laparoscopic bile duct injuries. *Arch Surg*. 1999;134:604-609.
- Bismuth H, Majno PE. Hepatobiliary surgery. *J Hepatol*. 2000;32(suppl):208-224.
- Flum DR, Cheadle A, Prela C, et al. Bile duct injury during cholecystectomy and survival in Medicare beneficiaries. *JAMA*. 2003;290:2168-2173.
- Traverso LW, Kozarek RA, Ball TJ, et al. Endoscopic retrograde cholangiopancreatography after laparoscopic cholecystectomy. *Am J Surg*. 1993;165:581-586.
- Bergman JJ, van der Brink GR, Rauws EAJ, et al. Treatment of bile duct lesions after laparoscopic cholecystectomy. *Gut*. 1996;38:141-147.