

Polygenic Scores for Major Depressive Disorder and Risk of Alcohol Dependence

Allan M. Andersen, MD; Robert H. Pietrzak, PhD, MPH; Henry R. Kranzler, MD; Li Ma, PhD; Hang Zhou, PhD; Xiaoming Liu, PhD; John Kramer, PhD; Samuel Kuperman, MD; Howard J. Edenberg, PhD; John I. Nurnberger Jr, MD, PhD; John P. Rice, PhD; Jay A. Tischfield, PhD; Alison Goate, PhD; Tatiana M. Foroud, PhD; Jacquelyn L. Meyers, PhD; Bernice Porjesz, PhD; Danielle M. Dick, PhD; Victor Hesselbrock, PhD; Eric Boerwinkle, PhD; Steven M. Southwick, MD; John H. Krystal, MD; Myrna M. Weissman, PhD; Douglas F. Levinson, MD; James B. Potash, MD, MPH; Joel Gelernter, MD; Shizhong Han, PhD

[+ Supplemental content](#)

IMPORTANCE Major depressive disorder (MDD) and alcohol dependence (AD) are heritable disorders with significant public health burdens, and they are frequently comorbid. Common genetic factors that influence the co-occurrence of MDD and AD have been sought in family, twin, and adoption studies, and results to date have been promising but inconclusive.

OBJECTIVE To examine whether AD and MDD overlap genetically, using a polygenic score approach.

DESIGN, SETTINGS, AND PARTICIPANTS Association analyses were conducted between MDD polygenic risk score (PRS) and AD case-control status in European ancestry samples from 4 independent genome-wide association study (GWAS) data sets: the Collaborative Study on the Genetics of Alcoholism (COGA); the Study of Addiction, Genetics, and Environment (SAGE); the Yale-Penn genetic study of substance dependence; and the National Health and Resilience in Veterans Study (NHRVS). Results from a meta-analysis of MDD (9240 patients with MDD and 9519 controls) from the Psychiatric Genomics Consortium were applied to calculate PRS at thresholds from $P < .05$ to $P \leq .99$ in each AD GWAS data set.

MAIN OUTCOMES AND MEASURES Association between MDD PRS and AD.

RESULTS Participants analyzed included 788 cases (548 [69.5%] men; mean [SD] age, 38.2 [10.8] years) and 522 controls (151 [28.9%] men; age [SD], 43.9 [11.6] years) from COGA; 631 cases (333 [52.8%] men; age [SD], 35.0 [7.7] years) and 756 controls (260 [34.4%] male; age [SD] 36.1 [7.7] years) from SAGE; 2135 cases (1375 [64.4%] men; age [SD], 39.4 [11.5] years) and 350 controls (126 [36.0%] men; age [SD], 43.5 [13.9] years) from Yale-Penn; and 317 cases (295 [93.1%] men; age [SD], 59.1 [13.1] years) and 1719 controls (1545 [89.9%] men; age [SD], 64.5 [13.3] years) from NHRVS. Higher MDD PRS was associated with a significantly increased risk of AD in all samples (COGA: best $P = 1.7 \times 10^{-6}$, $R^2 = 0.026$; SAGE: best $P = .001$, $R^2 = 0.01$; Yale-Penn: best $P = .035$, $R^2 = 0.0018$; and NHRVS: best $P = .004$, $R^2 = 0.0074$), with stronger evidence for association after meta-analysis of the 4 samples (best $P = 3.3 \times 10^{-9}$). In analyses adjusted for MDD status in 3 AD GWAS data sets, similar patterns of association were observed (COGA: best $P = 7.6 \times 10^{-6}$, $R^2 = 0.023$; Yale-Penn: best $P = .08$, $R^2 = 0.0013$; and NHRVS: best $P = .006$, $R^2 = 0.0072$). After recalculating MDD PRS using MDD GWAS data sets without comorbid MDD-AD cases, significant evidence was observed for an association between the MDD PRS and AD in the meta-analysis of 3 GWAS AD samples without MDD cases (best $P = .007$).

CONCLUSIONS AND RELEVANCE These results suggest that shared genetic susceptibility contributes modestly to MDD and AD comorbidity. Individuals with elevated polygenic risk for MDD may also be at risk for AD.

JAMA Psychiatry. 2017;74(11):1153-1160. doi:10.1001/jamapsychiatry.2017.2269
Published online August 16, 2017.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Shizhong Han, PhD, Department of Psychiatry and Behavioral Sciences, Johns Hopkins School of Medicine, 600 N Wolfe St, Baltimore, MD 21287 (han0312@gmail.com).

The association between alcohol dependence (AD) and major depressive disorder (MDD) has been studied by clinicians, epidemiologists, and geneticists for more than 100 years. Bleuler and Brill¹ described “alcoholic melancholia,” characterized by brief periods of depression after alcohol binges, which could precipitate a suicide attempt. States of depressed mood associated with both alcohol use and withdrawal were documented by later researchers,^{2,3} while others noted that some individuals who increased their drinking during periods of depression tended to progress to AD.⁴

More recently, the rate of AD and MDD comorbidity was assessed in cross-sectional epidemiologic studies. The National Comorbidity Study linked AD to 3.7-fold higher odds of experiencing a major depressive episode in the prior year.⁵ In the 2012-2013 National Epidemiologic Survey on Alcoholism and Related Conditions, individuals with a lifetime diagnosis of MDD had a 1.3-fold increased risk of an alcohol use disorder.⁶ Furthermore, both the National Comorbidity Study and Epidemiological Catchment Area study found that AD worsened the symptoms of MDD and vice versa,^{7,8} which may explain why people with comorbid AD and MDD are at greater risk for suicide.^{3,4,9-11}

This pattern of comorbidity has prompted hypotheses regarding causal relationships between AD and MDD.⁸ One hypothesis is that AD causes MDD either directly, through alcohol’s pharmacologic effects on the brain,¹²⁻¹⁴ or indirectly, through negative social consequences that increase the risk of MDD. A second hypothesis is that MDD can lead to AD. Longitudinal studies¹⁵⁻¹⁷ support an internalizing pathway from MDD to AD, with some individuals using alcohol to self-medicate negative mood states.

A third hypothesis, that shared genetic factors predispose to both AD and MDD, has had mixed support.^{4,8,18-20} A longitudinal study of Finnish twins showed a shared genetic liability between alcohol use and depressive symptoms across adolescence.²¹ Researchers from the Collaborative Study on the Genetics of Alcoholism (COGA)²² reported aggregation of MDD in relatives of probands with AD, whether or not the proband had MDD or the relative had AD. Other family studies have also suggested a genetic link between AD and MDD, including both sex-specific effects and environmental influences on their comorbidity.²³⁻²⁵

Molecular genetic studies have implicated specific genomic regions and genes for both AD and MDD risk. Linkage analysis by COGA identified a region on chromosome 1 associated with both MDD and AD.²⁶ Candidate gene studies reported a number of genes possibly underlying both disorders, including *CHRM2*,^{27,28} *SLC6A4*,^{29,30} *COMT*,^{31,32} and *DRD2*,^{33,34} although other studies failed to replicate these associations.³⁵⁻³⁸ Genome-wide association studies (GWASs) have produced significant findings for both AD³⁹⁻⁴¹ and MDD,^{42,43} although not for comorbid AD-MDD.⁴⁴

In the present study, we examined whether AD and MDD overlap genetically using polygenic risk scores (PRSs),^{45,46} which are quantitative measures of the cumulative effects of common genetic variations across the genome on risk for a disorder. Using PRSs derived from a Psychiatric Genomics Consortium (PGC) meta-analysis of MDD,⁴⁷ we tested whether PRSs

Key Points

Question Do major depressive disorder and alcohol dependence share common genetic risk variants?

Findings In this cohort study of 4 independent samples of 3871 individuals with alcohol dependence and 3347 individuals serving as controls, elevated polygenic risk for major depressive disorder also conveyed a significant increase in risk for alcohol dependence.

Meaning Common polygenic risk contributes to susceptibility to both major depressive disorder and alcohol dependence.

for MDD were associated with AD case-control status in samples from 4 independent GWAS data sets: (1) COGA; (2) the Study of Addiction, Genetics, and Environment (SAGE); (3) the Yale-Penn genetic study of substance dependence; and (4) the National Health and Resilience in Veterans Study (NHRVS).

Methods

AD GWAS Data Sets

Basic information on the AD GWAS data sets is summarized in **Table 1**. All participants in the COGA, Yale-Penn, and NHRVS studies gave informed consent for participation, and all procedures and protocols used in each sample were approved by the institutional review board at the relevant institution. Participants received financial compensation. COGA involved multiple centers in the United States: Indiana University; State University of New York Health Science Center, Brooklyn; University of Connecticut; University of Iowa; University of California, San Diego; Washington University; and Howard University. The institutional review boards of all of these participating institutions approved the study. The Yale-Penn study was approved by the Yale University Human Investigation Committee. The NHRVS procedures were approved by the human subjects subcommittee of the Veterans Affairs (VA) Connecticut Health-care System and the VA Office of Research and Development.

COGA

Samples were drawn from an ongoing, family-based study of AD including participants from 6 sites across the United States.⁴⁸ Patient cases had a lifetime diagnosis of AD by *DSM-IV* criteria. Individuals serving as controls had consumed alcohol but did not meet criteria for AD, alcohol abuse, or other substance use disorders in their lifetime. A sample of 1945 participants from COGA was genotyped (Illumina Human1M microarray; Illumina). After data quality control, 1310 European American individuals (788 with AD and 522 controls) were included.

SAGE

Samples were selected from 3 studies: COGA, the Family Study of Cocaine Dependence, and the Collaborative Genetic Study of Nicotine Dependence. Full details are available elsewhere.⁴⁹ Alcohol dependence was defined based on *DSM-IV* criteria. Controls were individuals without a lifetime diagnosis of dependence on any substance. A total of 4316 samples were geno-

Table 1. Sample Demographics

Characteristic	No. (%)			
	COGA	SAGE	Yale-Penn	NHRVS
AD	788	631	2135	317
Men	548 (69.5)	333 (52.8)	1375 (64.4)	295 (93.1)
Age, mean (SD), y	38.2 (10.8)	35.0 (7.7)	39.4 (11.5)	59.1 (13.1)
MDD	149 (18.9)	NA	414 (19.4)	111 (35.0)
Controls	522	756	350	1719
Men	151 (28.9)	260 (34.4)	126 (36.0)	1545 (89.9)
Age, mean (SD), y	43.9 (11.6)	36.1 (7.7)	43.5 (13.9)	64.5 (13.3)
MDD	65 (12.5)	NA	22 (6.3)	180 (10.5)

Abbreviations: AD, alcohol dependence; COGA, Collaborative Study on the Genetics of Alcoholism; MDD, major depressive disorder; NA, not available; NHRVS, National Health and Resilience in Veterans Study; SAGE, Study of Addiction, Genetics, and Environment.

typed (Illumina Human1M microarray). After quality control and excluding participants overlapping with COGA, 1387 European American individuals (631 with AD and 756 controls) were retained for analysis.

Yale-Penn

The Yale-Penn samples included individuals recruited for genetic studies of cocaine dependence, opioid dependence, and AD, as described previously.^{50,51} Alcohol dependence was defined using *DSM-IV* criteria. Control participants did not meet the *DSM-IV* criteria for lifetime AD or other substance use disorders. DNA samples from 9459 individuals were genotyped using 1 of 2 platforms: (1) the Illumina HumanOmni1 Quad, version 1.0 microarray containing 988 306 autosomal single-nucleotide polymorphisms (SNPs) or (2) the Illumina Infinium Human Core Exome microarray, containing 265 919 exome-focused SNPs and approximately 240 000 tagging SNPs. Genome-wide association studies quality control and imputation were conducted for the merged GWAS samples based on a common set of autosomal SNPs (230 447). After quality control, the final analysis included 2485 European American individuals (2135 with AD and 350 controls).

NHRVS

The NHRVS data set included US military veterans who participated in a nationally representative cohort study in 2011.^{52,53} Alcohol dependence was assessed according to *DSM-IV* diagnostic criteria. Controls were defined as individuals without lifetime AD or other substance dependence. A total of 2825 samples were genotyped with the Infinium PsychArray, version 1.1. After quality control, 2036 European American individuals (317 AD and 1719 controls) were retained for analysis.

GWAS Quality Control

Each GWAS data set was cleaned via PLINK, version 1.07 before analysis.⁵⁴ Samples and SNPs were excluded based on predetermined quality control metrics, including sample call rate of 95% or less, SNP call rate of 95% or less, minor allele frequency of 0.01 or less in controls, and *P* values for Hardy-Weinberg equilibrium tests $\leq 10^{-6}$ for controls. We estimated the genome-wide identity by descent sharing between all pairs of participants to detect duplicates (individuals genotyped twice within 1 sample or individuals in 1 sample also included in another) and then randomly removed 1 duplicate from each pair. We used EIGENSOFT, version 6.1.3⁵⁵ to compute princi-

pal components for each GWAS data set using pruned SNPs in low linkage disequilibrium ($r^2 < 0.2$). Outliers, defined as participants with ancestry at least 3 SDs from the mean on 1 of the 2 largest principal components, were removed. After quality control, there were 843 188 (COGA), 837 142 (SAGE), 220 554 (Yale-Penn), and 287 923 (NHRVS) autosomal SNPs retained. To ensure that sample overlap did not contribute to inflated estimates of genetic overlap between AD and MDD, λ meta statistics were calculated. The λ meta is a statistic that uses effect size concordance to detect sample overlap or heterogeneity.⁵⁶ All λ meta values were larger than 1 ($1.02 < \lambda$ meta < 1.06), indicating no significant overlap between AD and PGC MDD samples.

Imputation

Following the best practice guidelines of IMPUTE2, version 2.3.2,⁵⁷ we imputed 1000 Genomes variants into each GWAS sample. Prephasing was first performed with SHAPEIT, version 2.790⁵⁸ to infer haplotypes for samples based on autosomal SNPs with minor allele frequency greater than 0.01. Imputation was carried out on prephased haplotypes using IMPUTE2 against reference data from the 1000 Genomes Phase 1 integrated variant set for all GWAS data except for the NHRVS and Yale-Penn samples, which were imputed against the 1000 Genomes Phase 3 haplotype reference. After postimputation (SNP missing rate < 0.05 , minor allele frequency > 0.05 , imputation quality score > 0.5 , and Hardy-Weinberg equilibrium $> 10^{-6}$), 5 960 232 (COGA), 5 955 937 (SAGE), 7 049 271 (Yale-Penn), and 7 055 726 (NHRVS) autosomal variants were retained. Imputed SNPs were used for calculating the PRS, as described below.

Polygenic Score Analysis

Each PRS was calculated as a weighted sum of reference alleles across independent SNPs on a genome-wide scale. Weights were \log_{10} (odds ratio) of SNPs estimated from a meta-analysis of MDD GWASs (9240 MDD cases and 9519 controls of European ancestry) from the PGC.⁴⁷ Summary results for 1 235 109 autosomal SNPs were downloaded from the PGC website (<https://www.med.unc.edu/pgc/>). Downloaded SNPs were pruned using *P* value-informed clumping in PLINK, with a cutoff of $r^2 = 0.2$ within a 200-kilobase window. Polygenic risk scores were calculated for each AD GWAS sample using the PLINK score option, based on sets of pruned SNPs at successive *P* value thresholds ($< .05$ to $\leq .99$).

Table 2. Meta-analysis of Associations Between MDD Polygenic Risk Scores and AD

Thresholds	Adjusting for MDD in AD Samples					
	No			Yes		
	β	SE	P Value	β	SE	P Value
$P < .05$	0.075	0.016	2.8×10^{-6}	0.064	0.017	1.3×10^{-4}
$P < .10$	0.075	0.016	3.1×10^{-6}	0.061	0.017	2.5×10^{-4}
$P < .20$	0.084	0.016	1.0×10^{-7}	0.070	0.017	2.0×10^{-5}
$P < .30$	0.088	0.016	2.6×10^{-8}	0.074	0.017	8.4×10^{-6}
$P < .40$	0.095	0.016	3.3×10^{-9}	0.079	0.017	3.3×10^{-6}
$P < .50$	0.094	0.016	3.5×10^{-9}	0.080	0.017	2.5×10^{-6}
$P < .60$	0.094	0.016	4.2×10^{-9}	0.079	0.017	3.4×10^{-6}
$P < .70$	0.095	0.016	3.8×10^{-9}	0.079	0.017	3.2×10^{-6}
$P < .80$	0.093	0.016	6.5×10^{-9}	0.078	0.017	4.4×10^{-6}
$P < .90$	0.094	0.016	5.3×10^{-9}	0.079	0.017	3.7×10^{-6}
$P \leq .99$	0.094	0.016	5.3×10^{-9}	0.079	0.017	3.7×10^{-6}

Abbreviations: AD, alcohol dependence; MDD, major depressive disorder.

We used GEMMA, version 0.96 software⁵⁹ to evaluate associations between the PRS and AD in a mixed-model framework accounting for related samples and cryptic relationships. Association analyses were corrected for age, sex, and the top 10 principal components to control for population stratification. Each PRS was scaled from 0 to 2 so that association strength was comparable across GWAS data sets and *P* value thresholds. To combine association evidence across samples, we carried out meta-analysis based on effect size and SE using METAL.⁶⁰ The portion of the variance in AD case-control status explained by the PRS was assessed by the Nagelkerke pseudo R^2 ,^{45,46} which was derived from the difference between the maximum likelihood of the null model (principal components + covariates) and the full model (principal components + covariates + PRSs).

We conducted 2 additional analyses to further evaluate bidirectional relationships between MDD and AD: (1) to determine whether the association of the PRS with AD was driven by comorbid MDD cases in AD GWAS data sets, we repeated the analyses adjusting for MDD status in the 3 AD GWAS data sets (COGA, NHRVS, and Yale-Penn) for which MDD status was available; and (2) to examine whether the association of the PRS with AD was driven by comorbid AD cases in MDD PGC data sets, we recalculated the MDD PRS for AD GWAS data sets using weights from meta-analysis of 2 MDD GWAS data sets (Genetics of Recurrent Early-Onset Depression [GenRED]-I, 860 MDD cases and 1636 controls; and GenRED-II, 1145 MDD cases and 930 controls) for which comorbid MDD-AD cases could be excluded. Details of GenRED GWASs are available elsewhere.⁶¹ Both data sets were imputed into 1000 Genomes Phase 3 variants using imputation procedures described above. GEMMA was used for association testing of MDD with each SNP, followed by meta-analysis of the 2 imputed data sets using METAL. Meta-analysis results were used to recalculate the PRS for each AD GWAS data set. We then performed association tests between recalculated PRS and AD case-control status in 3 AD GWAS samples (COGA, Yale-Penn, and NHRVS) excluding MDD cases.

Positive and Negative Controls

We tested the ability of the PGC-derived MDD PRS to predict MDD status in 3 AD GWAS data sets (COGA, NHRVS, and Yale-Penn) for which MDD status was available as a positive control. Height was similarly tested as a negative control in the same samples.


Results

Basic sample demographics are reported in Table 1. In the COGA sample, 18.9% of individuals with AD had MDD and 12.5% of controls had MDD. In the Yale-Penn sample, 19.4% of those with AD had MDD and 6.3% of controls had MDD. The NHRVS sample had the highest rate of comorbid MDD, with 35.0% in AD cases; the MDD rate in controls was 10.5%. Major depressive disorder status was unavailable for SAGE.

We observed significant associations between the MDD PRS and AD case-control status across all AD samples at various PRS *P* value thresholds (COGA: best $P = 1.7 \times 10^{-6}$, $R^2 = 0.026$; SAGE: best $P = .001$, $R^2 = 0.01$; Yale-Penn: best $P = .035$, $R^2 = 0.0018$; NHRVS: best $P = .004$, $R^2 = 0.0074$) (eTable 1 in the [Supplement](#)). Meta-analysis of the 4 AD samples showed stronger evidence for association with the smallest value ($P = 3.3 \times 10^{-9}$; *P* value threshold = .4) (Table 2). Regression coefficients (β) were positive at each PRS *P* value threshold, suggesting that an elevated MDD PRS is linked to an increased risk of AD. However, the proportion of variance in AD explained by the MDD PRS was small, ranging from $R^2 = 0.0018$ in the Yale-Penn sample to $R^2 = 0.026$ in the COGA sample (eTable 1 in the [Supplement](#) and [Figure](#)). To investigate potential sex-specific effects, the same analyses were repeated stratified by sex. No consistent difference in the strength or significance of association between MDD PRSs and AD status by sex was observed across the 4 samples (eTable 2 and eFigure in the [Supplement](#)).

We observed similar patterns of association between the MDD PRSs and AD case-control status when MDD status was included as a covariate (COGA: best $P = 7.6 \times 10^{-6}$, $R^2 = 0.023$;

Figure. Association of the Major Depressive Disorder Polygenic Risk Score With Alcohol Dependence (AD) in 4 Independent Samples



Yale-Penn: best $P = .08$, $R^2 = 0.0013$; and NHRVS: best $P = .006$, $R^2 = 0.0072$ (eTable 3 in the Supplement). Meta-analysis of the 3 samples strengthened evidence of association with the smallest $P = 2.5 \times 10^{-6}$ (P value threshold = .50) (Table 2). The proportion of variance in AD explained by the MDD PRS was nearly identical to that of the unadjusted analyses in the corresponding samples.

We recalculated the MDD PRSs for AD GWAS samples using weights derived from 2 MDD GWAS data sets in which comorbid MDD-AD cases were excluded. We then tested the new MDD PRSs with AD case-control status in 3 AD GWAS samples without MDD cases. The new MDD PRS was nominally significantly associated with AD in the COGA sample at a PRS P value threshold of 0.05 ($P = .04$) (eTable 4 in the Supplement). We did not observe significant evidence for association in the Yale-Penn and NHRVS samples, although the new MDD PRSs tended to increase AD risk across both samples. Meta-analysis of the 3 AD samples yielded significant evidence for association between the MDD PRS and AD with the smallest $P = .007$ (PRS P value threshold = .20 or .05) (eTable 5 in the Supplement).

We tested whether the MDD PRS would predict MDD status as a positive control. A significant association was seen in the Yale-Penn sample ($7.6 \times 10^{-4} \leq P \leq .004$) but not in either COGA or NHRVS, although a trend toward increased MDD risk was observed in both samples (eTable 6 in the Supplement). In a meta-analysis, significant associations were seen at all thresholds ($2.5 \times 10^{-4} \leq P \leq .05$) (eTable 7 in the Supplement). Height, our negative control, showed no significant association with the MDD PRS in any sample at any P value threshold (eTable 8 in the Supplement).

Discussion

In each independent AD GWAS sample, we found that a higher MDD PRS was associated with an increased risk for AD, and this result was even more pronounced in a meta-analysis of the 4 data sets. The same effect was seen in analyses with MDD status as a covariate in 3 AD GWAS samples. We also observed meta-analytic evidence for an association between MDD PRSs

and AD, using a PRS derived from MDD GWAS data sets without comorbid MDD-AD cases. Our results suggest that the bidirectional associations between AD and MDD may be partially accounted for by shared genetic factors.

Our study methodology is similar to that of other published studies examining the association between AD genetic risk and AD age of onset⁶²; between childhood trauma, MDD genetic risk, and adult MDD⁶³; and between schizophrenia genetic risk and cannabis use.⁶⁴ Consistent with those studies, the proportion of variance in AD explained by the MDD PRS in our samples was small, on the order of 10^{-2} to 10^{-3} . Similar effect sizes were seen in our positive control, MDD status.

Although our results do not illuminate the mechanism by which this shared genetic liability for MDD and AD acts, several possibilities are worth investigating in future studies. Anxiety disorders and MDD share genetic liability,^{24,65} and anxiety has been proposed as a significant risk factor linking AD and MDD via the internalizing pathway.⁸ More broadly, personality traits, such as neuroticism, disinhibition, and sensation-seeking, are associated with a range of psychiatric disorders, including anxiety, MDD, and AD, and are potential mechanisms through which shared AD-MDD polygenic risk may exert its effects.⁶⁵ Inclusion of comorbid disorders and personality trait measures in future studies may help to clarify these pathways.

The PGC MDD GWAS samples are likely enriched for individuals with AD, which could drive the associations initially observed between MDD PRSs and AD. To examine this possibility, we recalculated the MDD PRS using 2 MDD GWAS data sets without comorbid MDD-AD cases and tested the association of the new PRS with AD in AD GWAS samples without MDD cases. We observed a significant association between this recalculated MDD PRS and AD status only in the COGA sample at a threshold of $P < .05$ ($P = .04$), potentially due to the smaller sample size of this training data set. Nonetheless, meta-analysis of all samples showed significant associations between the recalculated MDD PRS and AD, suggesting that the observed associations are not driven solely by comorbid AD cases in the PGC MDD GWAS and providing further support for genetic overlap between MDD and AD.

Prior family and twin studies^{20,23} suggested a role for sex-specific transmission of shared AD-MDD risk. In post hoc analyses, we examined sex-specific patterns of association between MDD PRSs and AD risk. We did not find consistent evidence of sex-specific transmission of shared AD-MDD liability in sex-stratified analyses, although our study was likely underpowered to detect such an effect given the reduced sample sizes in sex-specific subgroups.

Limitations

This work should be viewed in light of 2 limitations. First, MDD GWASs with larger sample sizes will likely improve the predictive ability of the MDD PRS and thus further refine the associations that we observed. Second, further studies are needed to investigate whether the MDD PRS association is specific to AD or whether it generalizes to substance dependence more broadly. There is evidence that a large proportion of genetic risk for sub-

stance use disorders involves predisposition to addiction generally,⁶⁶ and some data have demonstrated that PRSs for psychopathology predict general substance involvement.⁶⁷

Conclusions

From a public health standpoint, MDD and AD are individually and jointly associated with significant morbidity and mortality, so understanding their association is a priority. Our findings suggest that common genetic factors contribute to MDD-AD comorbidity and that some individuals carry a genetic predisposition for both disorders. The consistency of our findings across 4 independent samples suggests the feasibility and value of a meta-analysis of AD and MDD GWAS data sets to identify specific genetic variations underlying this shared predisposition.

ARTICLE INFORMATION

Published Online: August 16, 2017.

doi:10.1001/jamapsychiatry.2017.2269

Author Affiliations: Department of Psychiatry, Carver College of Medicine, University of Iowa, Iowa City (Andersen, Kramer, Kuperman, Potash, Han); US Department of Veterans Affairs (VA) National Center for Posttraumatic Stress Disorder, Clinical Neurosciences Division, VA Connecticut Healthcare System, West Haven (Pietrzak, Southwick, Krystal, Gelernter); Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut (Pietrzak, Zhou, Southwick, Krystal, Gelernter); Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia (Kranzler); Mental Illness, Research, Education and Clinical Center of Veterans Integrated Service Network 4, Corporal Michael J. Crescenz VA Medical Center, Philadelphia, Pennsylvania (Kranzler); Department of Animal and Avian Sciences, University of Maryland, College Park (Ma); Human Genetics Center, University of Texas Health Science Center at Houston (Liu, Boerwinkle); Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis (Edenberg); Department of Psychiatry, Indiana University School of Medicine, Indianapolis (Nurnberger); Department of Psychiatry, Washington University School of Medicine, St Louis, Missouri (Rice); Department of Genetics and the Human Genetics Institute of New Jersey, Rutgers University, Piscataway (Tischfield); Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, New York (Goate); Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis (Foroud); Department of Psychiatry and Behavioral Sciences, State University of New York Downstate Medical Center, Brooklyn (Meyers, Porjesz); Departments of Psychology and Human and Molecular Genetics, Virginia Commonwealth University, Richmond (Dick); Department of Psychiatry, University of Connecticut School of Medicine, Farmington (Hesselbrock); Division of Epidemiology, New York State Psychiatric Institute, New York (Weissman); Department of Psychiatry, Columbia University, College of Physicians and Surgeons, New York, New York (Weissman); Columbia University, Mailman School of Public

Health, New York, New York (Weissman); Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, California (Levinson); Interdisciplinary Graduate Program in Genetics, University of Iowa, Iowa City (Potash); now with the Department of Psychiatry and Behavioral Sciences, Johns Hopkins School of Medicine, Baltimore, Maryland (Potash, Han).

Author Contributions: Dr Han had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Kramer, Kuperman, Edenberg, Tischfield, Hesselbrock, Potash, Gelernter, Han.

Acquisition, analysis, or interpretation of data: Andersen, Pietrzak, Kranzler, Ma, Zhou, Liu, Kramer, Kuperman, Edenberg, Nurnberger, Rice, Goate, Foroud, Meyers, Porjesz, Dick, Hesselbrock, Boerwinkle, Southwick, Krystal, Weissman, Levinson, Potash, Gelernter, Han.

Drafting of the manuscript: Andersen, Kuperman, Han.

Critical revision of the manuscript for important intellectual content: Pietrzak, Kranzler, Ma, Zhou, Liu, Kramer, Kuperman, Edenberg, Nurnberger, Rice, Tischfield, Goate, Foroud, Meyers, Porjesz, Dick, Hesselbrock, Boerwinkle, Southwick, Krystal, Weissman, Levinson, Potash, Gelernter.

Statistical analysis: Ma, Zhou, Rice, Foroud, Han.

Obtained funding: Andersen, Pietrzak, Kuperman, Edenberg, Nurnberger, Tischfield, Porjesz, Dick, Hesselbrock, Boerwinkle, Southwick, Weissman, Levinson, Potash, Han.

Administrative, technical, or material support: Pietrzak, Kranzler, Liu, Kuperman, Edenberg, Tischfield, Hesselbrock, Southwick.

Study supervision: Potash.

Conflict of Interest Disclosures: Dr Kranzler has been a paid consultant, advisory board member, or continuing medical education speaker for Indivior and Lundbeck. He is also a member of the American Society of Clinical Psychopharmacology's Alcohol Clinical Trials Initiative, which was supported in the past 3 years by AbbVie, Alkermes, Ethypharm, Indivior, Lilly, Lundbeck, Otsuka, Pfizer, and Xenoport. No other conflicts were reported.

Funding/Support: Dr Andersen was funded by the American Academy of Child and Adolescent

Psychiatry Physician Scientist Program in Substance Abuse K12 award (K12DA000357) and received departmental support from the University of Iowa Carver College of Medicine, Department of Psychiatry. This study was supported by National Institutes of Health (NIH) grants R01AA022994, R01AA024486, and R01AA11330. The US Department of Veterans Affairs National Center for Posttraumatic Stress Disorder provided funding for the National Health and Resilience in Veterans Study.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We acknowledge the Psychiatric Genomics Consortium for making the major depression disorder genome-wide association study (GWAS) results publicly available. Genotyping services for part of the Yale-Penn GWAS were provided by the Center for Inherited Disease Research (CIDR) and Yale University (Center for Genome Analysis). The CIDR is fully funded through federal contract N01-HG-65403 from the NIH to The Johns Hopkins University. The Collaborative Study on the Genetics of Alcoholism (COGA) (principal investigators Bernice Porjesz, PhD, Victor Hesselbrock, PhD, Howard J. Edenberg, PhD, and Laura Bierut, MD) includes 12 different centers: University of Connecticut (Victor Hesselbrock, PhD); Indiana University (Howard J. Edenberg, PhD, John I. Nurnberger Jr, MD, PhD, and Tatiana M. Foroud, PhD); University of Iowa (Samuel Kuperman, MD, and John Kramer, PhD); State University of New York (SUNY) Downstate (Bernice Porjesz, PhD); Washington University in St. Louis (Laura Bierut, MD, John P. Rice, PhD, Kathleen Bucholz, PhD, and Arpana Agrawal, PhD); University of California at San Diego (Marc Schuckit, MD); Rutgers University (Jay A. Tischfield, PhD, and Andrew Brooks, PhD); Department of Biomedical and Health Informatics, The Children's Hospital of Philadelphia; Department of Genetics, Perelman School of Medicine, University of Pennsylvania (Laura Almasy, PhD); Virginia Commonwealth University (Danielle M. Dick, PhD); Icahn School of Medicine at Mount Sinai (Alison Goate, PhD); and

Howard University (Robert Taylor, PhD). Other COGA collaborators include Lance Bauer, PhD (University of Connecticut); Jeanette McClintick, PhD, Leah Wetherill, PhD, Xiaoling Xuei, PhD, Yunlong Liu, PhD, Dongbing Lai, PhD, Sean O'Connor, MD, Martin Plawecki, MD, PhD, and Spencer Lourens, PhD (Indiana University); Grace Chan, PhD (University of Iowa; University of Connecticut); Jacquelyn Meyers, PhD, David Chorlian, PhD, Chella Kamarajan, PhD, Ashwini Pandey, PhD, and Jian Zhang, PhD (SUNY Downstate); Jen Wang, PhD, Manav Kapoor, PhD, and Sarah Bertelsen, JD (Icahn School of Medicine at Mount Sinai); Andrey Anokhin, PhD, Vivian McCutcheon, PhD, and Scott Saccone, PhD (Washington University); Jessica Salvatore, PhD, Fazil Aliev, PhD, and Seung Cho, PhD (Virginia Commonwealth University); and Mark Kos, PhD (University of Texas Rio Grande Valley). Abbas Parsian, PhD, and Matthew Reilly, PhD, are the National Institute on Alcohol Abuse and Alcoholism (NIAAA) staff collaborators. We continue to be inspired by our memories of Henri Begleiter and Theodore Reich, founding principal investigator and co-principal investigator of COGA, and also owe a debt of gratitude to other past organizers of COGA, including Ting-Kai Li, P. Michael Connolly, Raymond Crowe, and Wendy Reich, for their critical contributions. This national collaborative study is supported by NIH grant U01AA008401 from the NIAAA and the National Institute on Drug Abuse (NIDA). The Study of Addiction: Genetics and Environment (SAGE) and COGA GWAS data sets in this study were obtained from the Database of Genotypes and Phenotypes (dbGaP). Funding support for SAGE was provided through the NIH Genes, Environment and Health Initiative (GEI) grant U01HG004422. SAGE is one of the GWASs funded as part of the Gene Environment Association Studies (GENEVA) under GEI. Assistance with phenotype harmonization and genotype cleaning, as well as with general study coordination, was provided by the GENEVA Coordinating Center grant U01HG004446. Assistance with data cleaning was provided by the National Center for Biotechnology Information. Support for collection of data sets and samples was provided by COGA grant U01AA008401, the Collaborative Genetic Study of Nicotine Dependence grant P01CA089392, and the Family Study of Cocaine Dependence grant R01DA013423. Funding support for genotyping, which was performed at the Johns Hopkins University Center for Inherited Disease Research, was provided by NIH GEI U01HG004438, NIAAA, NIDA, and NIH contract "high throughput genotyping for studying the genetic contributions to human disease" (HHSN268200782096C). The COGA case-control GWAS data set used for the analyses described in this article was obtained from dbGaP (accession number phs000125.v1.p1). Genotyping was performed at the Johns Hopkins University Center for Inherited Disease Research. Funding for data collection, phenotype harmonization, genotype cleaning, and analysis was through COGA grant U01AA008401.

REFERENCES

- Bleuler E, Brill AA. *Textbook of Psychiatry*. New York: Macmillan; 1924.
- Woodruff RA Jr, Guze SB, Clayton PJ, Carr D. Alcoholism and depression. *Arch Gen Psychiatry*. 1973;28(1):97-100.
- Whiters AC, Cadoret RJ, Widmer RB. Factors associated with suicide attempts in alcohol abusers. *J Affect Disord*. 1985;9(1):19-23.
- Pitts FN, Winokur G. Affective disorder—VII: alcoholism and affective disorder. *J Psychiatr Res*. 1966;4(1):37-50.
- Kessler RC, Nelson CB, McGonagle KA, Edlund MJ, Frank RG, Leaf PJ. The epidemiology of co-occurring addictive and mental disorders: implications for prevention and service utilization. *Am J Orthopsychiatry*. 1996;66(1):17-31.
- Grant BF, Goldstein RB, Saha TD, et al. Epidemiology of DSM-5 alcohol use disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry*. 2015;72(8):757-766.
- Regier DA, Farmer ME, Rae DS, et al. Comorbidity of mental disorders with alcohol and other drug abuse: results from the Epidemiologic Catchment Area (ECA) Study. *JAMA*. 1990;264(19):2511-2518.
- Swendsen JD, Merikangas KR. The comorbidity of depression and substance use disorders. *Clin Psychol Rev*. 2000;20(2):173-189.
- Cornelius JR, Salloum IM, Mezzich J, et al. Disproportionate suicidality in patients with comorbid major depression and alcoholism. *Am J Psychiatry*. 1995;152(3):358-364.
- Wilcox HC, Conner KR, Caine ED. Association of alcohol and drug use disorders and completed suicide: an empirical review of cohort studies. *Drug Alcohol Depend*. 2004;76(suppl):S11-S19.
- Angst F, Stassen HH, Clayton PJ, Angst J. Mortality of patients with mood disorders: follow-up over 34-38 years. *J Affect Disord*. 2002;68(2-3):167-181.
- Brujnzeel AW, Gold MS. The role of corticotropin-releasing factor-like peptides in cannabis, nicotine, and alcohol dependence. *Brain Res Brain Res Rev*. 2005;49(3):505-528.
- Feldstein Ewing SW, Filbey FM, Chandler LD, Hutchison KE. Exploring the relationship between depressive and anxiety symptoms and neuronal response to alcohol cues. *Alcohol Clin Exp Res*. 2010;34(3):396-403.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed, text revision. Washington, DC: American Psychiatric Association; 2000.
- Crum RM, Green KM, Storr CL, et al. Depressed mood in childhood and subsequent alcohol use through adolescence and young adulthood. *Arch Gen Psychiatry*. 2008;65(6):702-712.
- Crum RM, Mojtabai R, Lazareck S, et al. A prospective assessment of reports of drinking to self-medicate mood symptoms with the incidence and persistence of alcohol dependence. *JAMA Psychiatry*. 2013;70(7):718-726.
- Del Boca FK, Hesselbrock MN. Gender and alcoholic subtypes. *Alcohol Res Health*. 1996;20(1):56.
- Winokur G, Cadoret R, Dorzab J, Baker M. Depressive disease: a genetic study. *Arch Gen Psychiatry*. 1971;24(2):135-144.
- Winokur G, Coryell W. Familial alcoholism in primary unipolar major depressive disorder. *Am J Psychiatry*. 1991;148(2):184-188.
- Cadoret RJ, Winokur G, Langbehn D, Troughton E, Yates WR, Stewart MA. Depression spectrum disease, I: the role of gene-environment interaction. *Am J Psychiatry*. 1996;153(7):892-899.
- Edwards AC, Sihvola E, Korhonen T, et al. Depressive symptoms and alcohol use are genetically and environmentally correlated across adolescence. *Behav Genet*. 2011;41(4):476-487.
- Nurnberger JI Jr, Wiegand R, Bucholz K, et al. A family study of alcohol dependence: coaggregation of multiple disorders in relatives of alcohol-dependent probands. *Arch Gen Psychiatry*. 2004;61(12):1246-1256.
- Kendler KS, Heath AC, Neale MC, Kessler RC, Eaves LJ. Alcoholism and major depression in women: a twin study of the causes of comorbidity. *Arch Gen Psychiatry*. 1993;50(9):690-698.
- Kendler KS, Walters EE, Neale MC, Kessler RC, Heath AC, Eaves LJ. The structure of the genetic and environmental risk factors for six major psychiatric disorders in women: phobia, generalized anxiety disorder, panic disorder, bulimia, major depression, and alcoholism. *Arch Gen Psychiatry*. 1995;52(5):374-383.
- Prescott CA, Aggen SH, Kendler KS. Sex-specific genetic influences on the comorbidity of alcoholism and major depression in a population-based sample of US twins. *Arch Gen Psychiatry*. 2000;57(8):803-811.
- Nurnberger JI Jr, Foroud T, Flury L, et al. Evidence for a locus on chromosome 1 that influences vulnerability to alcoholism and affective disorder. *Am J Psychiatry*. 2001;158(5):718-724.
- Edenberg HJ, Foroud T. The genetics of alcoholism: identifying specific genes through family studies. *Addict Biol*. 2006;11(3-4):386-396.
- Wang JC, Hinrichs AL, Stock H, et al. Evidence of common and specific genetic effects: association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Hum Mol Genet*. 2004;13(17):1903-1911.
- Dick DM, Plunkett J, Hamlin D, et al. Association analyses of the serotonin transporter gene with lifetime depression and alcohol dependence in the Collaborative Study on the Genetics of Alcoholism (COGA) sample. *Psychiatr Genet*. 2007;17(1):35-38.
- Gokturk C, Schultz S, Nilsson KW, von Knorring L, Orelund L, Hallman J. Serotonin transporter (5-HTTLPR) and monoamine oxidase (MAOA) promoter polymorphisms in women with severe alcoholism. *Arch Womens Ment Health*. 2008;11(5-6):347-355.
- Baekken PM, Skorpen F, Stordal E, Zwart JA, Hagen K. Depression and anxiety in relation to catechol-O-methyltransferase Val¹⁵⁸Met genotype in the general population: the Nord-Trøndelag Health Study (HUNT). *BMC Psychiatry*. 2008;8:48.
- Serý O, Didden W, Mikes V, Pitelevá R, Znojil V, Zvolský P. The association between high-activity COMT allele and alcoholism. *Neuro Endocrinol Lett*. 2006;27(1-2):231-235.
- Dick DM, Wang JC, Plunkett J, et al. Family-based association analyses of alcohol dependence phenotypes across DRD2 and neighboring gene ANKK1. *Alcohol Clin Exp Res*. 2007;31(10):1645-1653.

34. Kōks S, Nikopensus T, Koido K, et al. Analysis of SNP profiles in patients with major depressive disorder. *Int J Neuropsychopharmacol*. 2006;9(2):167-174.
35. Cohen-Woods S, Gaysina D, Craddock N, et al. Depression Case Control (DeCC) Study fails to support involvement of the muscarinic acetylcholine receptor M2 (*CHRM2*) gene in recurrent major depressive disorder. *Hum Mol Genet*. 2009;18(8):1504-1509.
36. Furlong RA, Coleman TA, Ho L, et al. No association of a functional polymorphism in the dopamine D2 receptor promoter region with bipolar or unipolar affective disorders. *Am J Med Genet*. 1998;81(5):385-387.
37. Gillespie NA, Whitfield JB, Williams B, Heath AC, Martin NG. The relationship between stressful life events, the serotonin transporter (5-HTTLPR) genotype and major depression. *Psychol Med*. 2005;35(1):101-111.
38. Serretti A, Rotondo A, Lorenzi C, Smeraldi E, Cassano GB. Catechol-O-methyltransferase gene variants in mood disorders in the Italian population. *Psychiatr Genet*. 2006;16(5):181-182.
39. Gelernter J, Kranzler HR, Sherva R, et al. Genome-wide association study of alcohol dependence: significant findings in African- and European-Americans including novel risk loci. *Mol Psychiatry*. 2014;19(1):41-49.
40. Zuo L, Tan Y, Zhang X, et al. A new genome-wide association meta-analysis of alcohol dependence. *Alcohol Clin Exp Res*. 2015;39(8):1388-1395.
41. Kapoor M, Wang JC, Wetherill L, et al. Genome-wide survival analysis of age at onset of alcohol dependence in extended high-risk COGA families. *Drug Alcohol Depend*. 2014;142:56-62.
42. Hyde CL, Nagle MW, Tian C, et al. Identification of 15 genetic loci associated with risk of major depression in individuals of European descent. *Nat Genet*. 2016;48(9):1031-1036.
43. Okbay A, Baselmans BM, De Neve JE, et al; LifeLines Cohort Study. Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nat Genet*. 2016;48(6):624-633.
44. Edwards AC, Aliev F, Bierut LJ, et al. Genome-wide association study of comorbid depressive syndrome and alcohol dependence. *Psychiatr Genet*. 2012;22(1):31-41.
45. Purcell SM, Wray NR, Stone JL, et al; International Schizophrenia Consortium. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460(7256):748-752.
46. Power RA, Steinberg S, Bjornsdottir G, et al. Polygenic risk scores for schizophrenia and bipolar disorder predict creativity. *Nat Neurosci*. 2015;18(7):953-955.
47. Ripke S, Wray NR, Lewis CM, et al; Major Depressive Disorder Working Group of the Psychiatric GWAS Consortium. A mega-analysis of genome-wide association studies for major depressive disorder. *Mol Psychiatry*. 2013;18(4):497-511.
48. Edenberg HJ, Koller DL, Xuei X, et al. Genome-wide association study of alcohol dependence implicates a region on chromosome 11. *Alcohol Clin Exp Res*. 2010;34(5):840-852.
49. Bierut LJ, Agrawal A, Bucholz KK, et al; Gene, Environment Association Studies Consortium. A genome-wide association study of alcohol dependence. *Proc Natl Acad Sci U S A*. 2010;107(11):5082-5087.
50. Gelernter J, Kranzler HR, Sherva R, et al. Genome-wide association study of opioid dependence: multiple associations mapped to calcium and potassium pathways. *Biol Psychiatry*. 2014;76(1):66-74.
51. Gelernter J, Sherva R, Koesterer R, et al. Genome-wide association study of cocaine dependence and related traits: *FAM53B* identified as a risk gene. *Mol Psychiatry*. 2014;19(6):717-723.
52. Fuehrlein BS, Mota N, Arias AJ, et al. The burden of alcohol use disorders in US military veterans: results from the National Health and Resilience in Veterans Study. *Addiction*. 2016;111(10):1786-1794.
53. Wisco BE, Marx BP, Wolf EJ, Miller MW, Southwick SM, Pietrzak RH. Posttraumatic stress disorder in the US veteran population: results from the National Health and Resilience in Veterans Study. *J Clin Psychiatry*. 2014;75(12):1338-1346.
54. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81(3):559-575.
55. Patterson N, Price AL, Reich D. Population structure and eigenanalysis. *PLoS Genet*. 2006;2(12):e190.
56. Chen GB, Lee SH, Robinson MR, et al; Genetic Investigation of Anthropometric Traits (GIANT) Consortium. Cross-cohort QC analyses of GWAS summary statistics from complex traits. *Eur J Hum Genet*. 2016;25(1):137-146.
57. Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet*. 2009;5(6):e1000529.
58. Delaneau O, Marchini J, Zagury JF. A linear complexity phasing method for thousands of genomes. *Nat Methods*. 2011;9(2):179-181.
59. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies. *Nat Genet*. 2012;44(7):821-824.
60. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26(17):2190-2191.
61. Shyn SI, Shi J, Kraft JB, et al. Novel loci for major depression identified by genome-wide association study of sequenced treatment alternatives to relieve depression and meta-analysis of three studies. *Mol Psychiatry*. 2011;16(2):202-215.
62. Kapoor M, Chou YL, Edenberg HJ, et al. Genome-wide polygenic scores for age at onset of alcohol dependence and association with alcohol-related measures. *Transl Psychiatry*. 2016;6(3):e761.
63. Peyrot WJ, Milaneschi Y, Abdellaoui A, et al. Effect of polygenic risk scores on depression in childhood trauma. *Br J Psychiatry*. 2014;205(2):113-119.
64. Power RA, Verweij KJ, Zuhair M, et al. Genetic predisposition to schizophrenia associated with increased use of cannabis. *Mol Psychiatry*. 2014;19(11):1201-1204.
65. Kendler KS, Aggen SH, Knudsen GP, Røysamb E, Neale MC, Reichborn-Kjennerud T. The structure of genetic and environmental risk factors for syndromal and subsyndromal common DSM-IV axis I and all axis II disorders. *Am J Psychiatry*. 2011;168(1):29-39.
66. Kendler KS, Chen X, Dick D, et al. Recent advances in the genetic epidemiology and molecular genetics of substance use disorders. *Nat Neurosci*. 2012;15(2):181-189.
67. Carey CE, Agrawal A, Bucholz KK, et al. Associations between polygenic risk for psychiatric disorders and substance involvement. *Front Genet*. 2016;7:149.